

Clinical Trial Pre-Screening Report

Generated: 2026-07-31 18:22 UTC

Patient Profile

Field	Value
Patient ID	P-5236
Age	45
Gender	male
Conditions	Hypertension, Prediabetes, Anemia (disorder), Diabetes, Acute bronchitis (disorder), Hypertriglyceridemia (disorder), Metabolic syndrome X (disorder), Chronic kidney disease stage 1 (disorder), Diabetic renal disease (disorder), Acute viral pharyngitis (disorder), Hyperglycemia (disorder)
Medications	Insulin glargine injection, Empagliflozin 10 MG Oral Tablet

Lab Results:

- hba1c: **8.4 %** (recorded 2026-05-08)
- egfr: **84.0 mL/min/1.73m2** (recorded 2026-05-06)
- bmi: **37.3 kg/m2** (recorded 2026-05-01)
- hba1c: **9.4 %** (recorded 2025-03-25)

Safety note: This report is a pre-screening aid only. It does not constitute a final eligibility decision, diagnosis, or clinical recommendation.

Executive Summary

Showing **3** top-ranked trial(s) (capped at 3).

Clinical Fit	Count
[OK] Supported	0
[NO] Not Supported	0
[?] Other	3
[R] Human Review Required	3

Trial Results

1. Assessing the Efficacy of Targeted Home Visits in the Management of Chronic Conditions

Trial ID: NCT05168605

Reason surfaced: 2/8 evaluated criteria supported; trial is actively recruiting.

Field	Status
Clinical Fit	[?] Insufficient data
Recruiting	[OPEN] Actively recruiting
Score	0.250
Human Review	[!] Required

Criterion Evaluation

Criterion	Status	Evaluator	Reasoning	Evidence
inc_0	[OK] SUPPORTED	rule_engine	Patient age 45 is within the required range 18-60.	--
inc_1	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_2	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Inclusion
inc_3	[OK] SUPPORTED	rule_engine	Patient HBA1C 8.4 meets criterion > 8.0.	HBA1C (2026-05-08)
inc_4	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_0	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_1	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_2	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	Age criterion text '* Language other than English or Spanish' could not be parsed automatically.	--

Note: Some criteria requiring LLM evaluation (inc_2) could not be automatically assessed because the LLM service was unavailable. Please review these criteria manually.

Unanswered Questions / Items for Clinical Review

- Please review this criterion manually with the study team.
- Automated evaluation unavailable -- please review manually.
- Clinical review required -- please evaluate manually.

2. GLUCOSE-MGH: Genetic Links Understood Through Challenge With Oral Semaglutide Exposure at MGH

Trial ID: NCT06003153

Reason surfaced: 2/13 evaluated criteria supported; trial is actively recruiting.

Field	Status
Clinical Fit	[?] Insufficient data
Recruiting	[OPEN] Actively recruiting
Score	0.154
Human Review	[!] Required

Criterion Evaluation

Criterion	Status	Evaluator	Reasoning	Evidence
inc_0	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_1	[OK] SUPPORTED	rule_engine	Patient age 45 is within the required range 18-65.	--
inc_2	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_3	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion
exc_0	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion
exc_1	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion
exc_2	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion
exc_3	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion
exc_4	[OK] SUPPORTED	rule_engine	Patient EGFR 84.0 does not meet criterion < 60.0.	EGFR (2026-05-06)
exc_5	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_6	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_7	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_8	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--

Note: Some criteria requiring LLM evaluation (inc_3, exc_0, exc_1, exc_2, exc_3) could not be automatically assessed because the LLM service was unavailable. Please review these criteria manually.

Unanswered Questions / Items for Clinical Review

- Please review this criterion manually with the study team.
- Automated evaluation unavailable -- please review manually.

3. Primary Care Pragmatic, Real World Experience for Automated Insulin Delivery

Trial ID: NCT07011147

Reason surfaced: 5/48 evaluated criteria supported; trial is actively recruiting.

Field	Status
Clinical Fit	[?] Insufficient data
Recruiting	[OPEN] Actively recruiting
Score	0.104
Human Review	[!] Required

Criterion Evaluation

Criterion	Status	Evaluator	Reasoning	Evidence
inc_0	[OK] SUPPORTED	rule_engine	Patient age 45 meets criterion age > 18.	--
inc_1	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_2	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent \>18 and \<89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

inc_4	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	Age criterion text 'Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents' coul...	--
inc_5	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	Age criterion text 'Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.' could not be parsed automatic...	--
inc_6	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent ≥ 18 and ≤ 89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or >2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

inc_7	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	Age criterion text 'Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage th...	--
inc_8	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_9	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_10	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent \>18 and \<89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonyleurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the

participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

inc_11	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_12	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_13	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_14	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_15	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_16	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_17	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_18	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_19	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_20	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
inc_21	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_0	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_1	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_2	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent \>18 and \<89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or ≥ 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or ≥ 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or ≥ 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or ≥ 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

exc_6	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_7	[OK] SUPPORTED	rule_engine	Trial recruiting status verified in the filtering stage.	--
exc_8	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent >18 and <89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).

ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).

iii. Placement of an intrauterine device or intrauterine hormone-releasing system.

iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).

v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.

vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within

the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

exc_9	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	HBA1C criterion text 'Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere...	--
exc_10	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_11	[OK] SUPPORTED	rule_engine	Trial recruiting status verified in the filtering stage.	--
exc_12	[R] REQUIRE S_CLINICAL_REVIEW	llm_engine_error	LLM service unavailable -- manual review required.	Demographics; Conditions; Medication: Insulin glargine injection; Medication: Empagliflozin 10 MG Oral Tablet; hba1c (2026-05-08); egfr (2026-05-06); bmi (2026-05-01); hba1c (2025-03-25); Exclusion; Medication: 1. Age at time of consent \>18 and \<89 years

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

- a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).
- ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).
- iii. Placement of an intrauterine device or intrauterine hormone-releasing system.
- iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).
- v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.
- vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

- a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses) b. Acceptable methods of contraception include: i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).
- ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).
- iii. Placement of an intrauterine device or intrauterine hormone-releasing system.
- iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).
- v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.
- vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or > 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

Either 2.a. or 2.b.:

Clinical diagnosis of type 1 diabetes for at least one year and using insulin for at least 1 year

Clinical diagnosis of type 2 diabetes, on current injected or infused insulin regimen for at least 3 months prior to screening (e.g., basal-bolus, basal only, or pre-mix)

Stable doses of glucose lowering medications over the preceding 4 weeks as determined by Investigator, including GLP-1 receptor agonists (GLP-1 RA) and GLP-1/GIP RA agents

Stable doses of weight loss medications (including GLP-1 RA and GLP-1/GIP RA agents) over the preceding 4 weeks as determined by the investigator.

For those using the iLet Bionic Pancreas (during the RCT arm or observational extension phase), willingness to stay on current doses of medications throughout the study that may affect glycemia directly and/or indirectly, except for a dose reduction or discontinuation.

Have a primary care clinician willing to refer them to the study, confirm their diabetes diagnosis (for example: type 1 diabetes or type 2 diabetes), and recommend and manage the iLet for the duration of the study

Willing to comply with all study procedures for the duration of the study

Willing to wear a Dexcom CGM device and iLet system for duration of time randomized to iLet use or OEP

Willing to use the following insulin: lispro (including non-branded lispro and Humalog) or aspart (including non-branded aspart, Fiasp, and Novolog)

Investigator has confidence that the participant has the cognitive ability and can successfully operate all study devices and can adhere to the protocol

Willing and able to sign and date the Informed Consent Form (ICF)

If capable of becoming pregnant, willing and able to have pregnancy testing and use an acceptable method of contraception during the study period

- a. Capable of becoming pregnant means that menstruation has started and the participant is not surgically sterile or post-menopausal (12 months without menses)
- b. Acceptable methods of contraception include:
 - i. Combined estrogen and progestogen containing hormonal contraception associated with inhibition of ovulation (oral, intravaginal, transdermal).
 - ii. Progestogen-only hormonal contraception associated with inhibition of ovulation (oral, injectable, implantable).
 - iii. Placement of an intrauterine device or intrauterine hormone-releasing system.
 - iv. Barrier methods of contraception (condom or occlusive cap with spermicidal foam/gel/film/cream/suppository).
 - v. Has a vasectomized or sterile partner (where partner is sole partner of participant) and where vasectomy has been confirmed by medical assessment.
 - vi. Exercises true sexual abstinence. Sexual abstinence is defined as refraining from heterosexual intercourse during the entire period of risk associated with the study.

Agreement to adhere to Lifestyle Considerations (see Section 5.3) throughout study duration

Have hardware and internet access capable of 2-way video and audio communication

Exclusion Criteria:

Unable to safely comply with study procedures and reporting requirements (e.g. impairment of vision or dexterity that prevents safe operation of the bionic pancreas, impaired memory)

Unable to speak and read English, as iLet BP support materials and device menus are currently available in English only

Diagnosis of maturity-onset diabetes of the young (MODY)

Plan to change usual diabetes regimen between screening and study randomization

This would include changing from MDI to pump or from pump to MDI, starting a new class of type 2 diabetes medication, or starting or increasing GLP-1 RA or GLP-1/GIP RA medication

This would NOT include changes to any insulin doses, including pump settings, short- and/or long-acting insulin doses and type of insulin; changing type 2 diabetes medication dosing (except GLP-1 RA or GLP-1/GIP RA); or changing type of type 2 diabetes medication within the same class

Weigh more than 255 kg (561 pounds) as this is the maximum weight that can be entered into the iLet user interface

History of bariatric surgery within 12 months prior to enrollment or plans for bariatric surgery within the period of study participation

Current use of a closed-loop or hybrid closed-loop insulin delivery system that is not FDA-cleared (e.g. "DIY Loop", "AAPS", "iAPS" or "Open APS")

Diagnosed blood disorder or dyscrasia associated with hemolysis, including for example: sickle cell disease and thalassemia, which in the Investigator's opinion could interfere with HbA1c accuracy

Planned use of hydroxyurea at any dose and/or of acetaminophen at doses exceeding 1 gram (1000 mg) every 6 hours.

Plans to receive a blood transfusion over the course of the study or has received a transfusion within 3 months prior to enrollment

Current participation in another diabetes-related clinical trial

History of diabetes due to cystic fibrosis, pancreatitis, or other pancreatic disease, including pancreatic tumor or insulinoma, or history of complete pancreatectomy

Have a history of intermittent oral or injectable glucocorticoid treatment within 8 weeks prior to screening or plans to take intermittent oral or injectable glucocorticoid during the study (chronic, stable treatment is acceptable, unplanned use is acceptable)

History of more than 1 episode of diabetic ketoacidosis (DKA) or hyperglycemic hyperosmolar syndrome (HHS) in the 6 months prior to screening, unrelated to an intercurrent illness or to a kinked, dislodged, or occluded cannula

Established history of allergy or severe reaction to adhesive or tape that must be used in the study

Treated currently or within the past 30 days prior to enrollment, or plan to begin treatment, with sulfonylurea, pramlintide, or SGLT-2 inhibitor medication

Any planned surgery during the study that would be considered major in the opinion of the investigator

Pregnant or lactating, or planning to become pregnant in the next 6 months

Renal failure on dialysis or chronic renal disease with a GFR or eGFR ≤ 30 mL/min (values within the last two years will be accepted; if none available or ≥ 2 years prior, participant will be instructed to obtain GFR or eGFR through their usual care provider and to make copy of result available to study team)

Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromise the results of the study c. Conditions to be considered by the investigator may include, but are not limited to, the following: i. Active clinical diagnosis of substance use disorder ii. Chronic use of opiates and/or benzodiazepines which, in the opinion of the investigator, might make it difficult for the participant to follow study procedures iii. Coronary artery disease that is not stable with medical management, including unstable angina, angina that prevents moderate exercise (e.g. exercise of intensity up to 6 METS) despite medical management, or within the last 12 months before screening, a history of myocardial infarction, percutaneous coronary intervention, enzymatic lysis of a presumed coronary occlusion, or coronary artery bypass grafting iv. Known history of prolonged QTc interval, malignant arrhythmia, or severe congenital heart disease v. Congestive heart failure with New York Heart Association (NYHA) Functional Classification III or IV vi. History of TIA or stroke in the last 12 months vii. Untreated or inadequately treated mental illness viii. History of untreated or inadequately treated eating disorder within the last 2 years, such as anorexia, bulimia, or diabulimia, or omission of insulin to manipulate weight ix. History of intentional, inappropriate administration of insulin leading to severe hypoglycemia requiring treatment

Plans to travel outside of the US and its territories for more than four weeks consecutively

Plans not to have internet/video/phone access for more than one week consecutively

Employed by, or having immediate family members employed by Beta Bionics, or being directly involved in conducting the clinical trial, or having a direct supervisor at place of employment who is also directly involved in conducting the clinical trial (as a study investigator, coordinator, etc.); or having a first-degree relative who is directly involved in conducting the clinical trial

Previous use of the iLet for more than four weeks (including use of the commercially available iLet or prior participation in a study involving wearing the iLet for more than four weeks) |

exc_15	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_16	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_17	[OK] SUPPORTED	rule_engine	Trial recruiting status verified in the filtering stage.	--
exc_18	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_19	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_20	[OK] SUPPORTED	rule_engine	Patient EGFR 84.0 does not meet criterion < 30.0.	EGFR (2026-05-06)
exc_21	[R] REQUIRE S_CLINICAL_REVIEW	rule_engine	Age criterion text 'Any condition or circumstance that, in the opinion of the site principal investigator, could interfere with the safe or effective completion of the study or which could compromi...	--
exc_22	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_23	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_24	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--
exc_25	[R] REQUIRE S_CLINICAL_REVIEW	clinical_review	Criterion type OTHER requires clinical judgement and cannot be automated.	--

Note: Some criteria requiring LLM evaluation (inc_2, inc_3, inc_6, inc_10, exc_2, exc_3, exc_4, exc_5, exc_8, exc_12, exc_13, exc_14) could not be automatically assessed because the LLM service was unavailable. Please review these criteria manually.

Unanswered Questions / Items for Clinical Review

- Please review this criterion manually with the study team.
- Automated evaluation unavailable -- please review manually.
- Clinical review required -- please evaluate manually.

Appendix -- Trials Excluded Before Evaluation

These trials were removed by deterministic filters:

Trial ID	Reason
NCT03301792	Status: ACTIVE_NOT_RECRUITING
NCT04892199	Status: ACTIVE_NOT_RECRUITING
NCT05066542	Status: ACTIVE_NOT_RECRUITING
NCT05181449	Status: ACTIVE_NOT_RECRUITING
NCT05766488	Status: ENROLLING_BY_INVITATION
NCT06094491	Status: ACTIVE_NOT_RECRUITING
NCT06254274	Status: ACTIVE_NOT_RECRUITING
NCT06649344	Status: ACTIVE_NOT_RECRUITING
NCT06682351	Status: NOT_YET_RECRUITING
NCT06760715	Status: NOT_YET_RECRUITING
NCT06770829	Status: NOT_YET_RECRUITING
NCT06775600	Status: NOT_YET_RECRUITING
NCT06973954	Status: ACTIVE_NOT_RECRUITING
NCT07047248	Status: NOT_YET_RECRUITING
NCT07215533	Age 45 outside criterion range
NCT07220759	Status: ACTIVE_NOT_RECRUITING
NCT07278531	Status: NOT_YET_RECRUITING
NCT07468188	Age 45 outside criterion range
NCT07509502	Status: NOT_YET_RECRUITING
NCT07510386	Status: NOT_YET_RECRUITING
NCT07628985	Status: NOT_YET_RECRUITING
NCT07702097	Status: NOT_YET_RECRUITING

Pre-screening report generated by the Clinical Trial Pre-Screening Assistant. This report is intended to support -- not replace -- clinical judgement. All recommendations must be verified by a qualified healthcare professional before any action is taken.