- c) B. Diandra, mentioned by Vss, which may be B. Diandra Linn. = B. Hirsuta Linn., or B. Diandra Burm.f. (see b);
- (2) B. VERTICILLATA POIR. (Dgm, p. 210) = B. REPANDA ROXB. (Ainslie 2, p. 207);
  - (3) B. REPANDA WILLD. (PA 1, 5, p. 9-26);
- (4) B. ALATA (Wise); the Index Kewensis records only B. ALATA S. Wats., an American species;
  - (5) B. PROCUMBENS BANKS EX ROXB. (HK?);
- (6) TRIANTHEMA PORTULACASTRUM LINN. (Bpn, p. 421: raktapunarnavā; Chopra: punarnavī; KB 2, p. 1180-1181: śvetapunarnavā; PA 1, 5, p. 9-26) = T. MONOGYNA LINN. (Chopra: punarnavī; Dgm, p. 210; DWH 2, p. 102-103: śvetapunarnavā; Ghosh: śvetapunarnavā; KB 2, p. 1180-1181; Nadk. 1, nr. 2491; PA 1, 5, p. 9-26; Sanyal) = T. OBCORDATA ROXB. (Nadk. 1, nr. 2491; PA 1, 5, p. 9-26);
  - (7) T. PENTANDRA LINN. (Nadk. 1, nr. 2491) = T. OBCORDATA WALL.;
- (8) T. DECANDRA LINN. (Chopra: punarnavī; Dgm, p. 210; KB 2, p. 1182: punarnavī; PA 1, 5, p. 9-26);
  - (9) T. CRYSTALLINA VAHL (Dgm, p. 210).

## $punn\bar{a}ga$

- (1) CALOPHYLLUM INOPHYLLUM LINN. (Dgv, nr. 40; Dgm, p. 235-236; Dutt; DWH 1, p. 173-176; HK; IRM 2, p. 18-19; KB 1, p. 270-271; MW; Na 1, nr. 45; Nadk. 1, nr. 425; V 6, p. 124; Vśs);
  - (2) C. TOMENTOSUM WIGHT (Dgm, p. 235-236);
  - (3) C. WIGHTIANUM WALL. (Dgm, p. 235-236);
  - (4) C. APETALUM WILLD. (Dgm, p. 235-236) = C. INOPHYLLUM?;
- (5) Ochrocarpus longifolius Benth. et Hook.f. ex T. Anders. (DWH 1, p. 172-173; KB 1, p. 269-270; Nadk. 1, nr. 731);
  - (6) [Callitris inophyllum Linn.] (Nadk. 1, nr. 422);
- (7) Mallotus Philippinensis Muell. Arg. (KB 3, p. 2267-2270) = Rottlera tinctoria Roxb. (KB 3, p. 2267; MW; PW; Roxb., p. 737);
- (8) nutmeg, i.e. the seed of Myristica fragrans Houtt. (jātīphala) (MW; PW; Vśs);
  - (9) white lotus (MW; PW; Vśs: śvetapadma).

## $prthv\bar{\imath}k\bar{a}$

- (1) Elettaria Cardamomum Maton (KB 4, p. 2442-2444; MW; PW; Vss);
  - (2) Amomum subulatum Roxb. (Gul; MW; PW; Vśs);