species of this name : a) I. ARGENTEA LINN., b) I. PAUCIFOLIA DELILE = I. ARGENTEA ROXB.;

- (5) I. oblongifolia Forsk. (KB 1, p. 711-712 : $n\bar{\imath}l\bar{a}$);
- (6) I. Angustifolia (Nadk. 1, nr. 1293); the Index Kewensis records three species: a) I. Angustifolia Linn., b) I. Hirsuta Linn. = I. Angustifolia Blanco, c) I. Stricta Linn. = I. Angustifolia Curt.

$n\bar{\imath}v\bar{a}ra$

- (1) a) the wild variety of rice (Dutt; HK; MW; PW; Vss);
 - b) ORYZA SATIVA LINN. (KB 4, p. 2651-2653);
- (2) Hygroryza aristata Nees (DWH 3, p. 620; Gul; KB 4, p. 2653-2654);
- (3) Setaria Italica Beauv. = Panicum Italicum Linn. (Bpn, p. 655).

nyagrodha

- (1) = vata;
- (2) = $\sin i$ (MW; PW).

paţola

- (1) TRICHOSANTHES DIOICA ROXB. (Avk; Bpn, p. 674; Chopra; Dgv, nr. 166; Dutt; DWH 2, p. 72-74; Dy, p. 343-344; Ghosh; Gul; HB; HK; Kav; KB 2, p. 1110-1111; MW; Nadk. 1, nr. 2508; PA 1, 7, p. 38-49; V 6, p. 36; Vés);
- (2) T. CUCUMERINA LINN. (Avk; Bpn, p. 674; Chopra; DWH 2, p. 72-74; Dy, p. 343-344; KB 2, p. 1112-1114; Na 1, nr. 227; Nadk. 1, nr. 2506; PA 1, 7, p. 38-49; Sanyal) = T. LACINIOSA KLEIN EX WILLD. (Ainslie 2, p. 296-297; PA 1, 7, p. 38) = T. LOBATA ROXB. (PA 1, 7, p. 38) = T. PILOSA BUCH.-HAM. EX WALL. (PA 1, 7, p. 38);
 - (3) T. CORDATA ROXB. (PA 1, 7, p. 38) = T. PALMATA WALL.;
 - (4) T. NERVIFOLIA LINN. (KB 2, p. 1111);
 - (5) T. BRACTEATA VOIGT (PA 1, 7, p. 38);
- (6) T. Pubera Blume = T. Palmata Roxb. (PA 1, 7, p. 38); acc. to PA 1, 7, p. 38 T. Bracteata Voigt is a synonym of T. Palmata Roxb.; this is not confirmed by the Index Kewensis.

patra

(1) CINNAMOMUM TAMALA T. NEES ET EBERM. (Bpn, p. $228 := tam\bar{a}la$