Nadk. 1, nr. 1034; PW; Vśs) = Urostigma Bengalense Gasp. (KB 3, p. 2312-2314); see Hooker 5, p. 499-500 about the nomenclature.

vanakārpāsī

- (1) Hibiscus vitifolius Linn. (Avk; Dutt; Kav and Vśs, = $bh\bar{a}radv\bar{a}\bar{j}\bar{i}$) = H. Truncatus Roxb. (Avk);
- (2) Thespesia macrophylla Blume = Th. Lampas Dalz. et Gibs. (Avk; Bpn, p. 375) = Hibiscus Lampas Cav. (Avk);
- (3) Gossypium nigrum Buch.-Ham. (Avk); this is an obsolete name, which covers two species acc. to the Index Kewensis:
 - a) G. BARBADENSE LINN.;
 - b) G. ARBOREUM LINN.

vanakodrava = kodrava

$vanabarbarikar{a}$

- (1) OCIMUM GRATISSIMUM LINN. (KB 3, p. 1964-1965);
- (2) O. BASILICUM LINN. = O. PILOSUM WILLD. (MW);
- (3) = kuthera (PW).

vanamudga

- (1) Phaseolus trilobus Ait. (Avk; KB 1, p. 794-795; Nadk. 1, nr. 1870, = mudgaparnī);
- (2) PH. ACONITIFOLIUS JACQ. (Bpn, p. 646 and KB 1, p. 798-799, = makuṣṭha; MW; Vśs, = makuṣṭhaka) == PH. TRILOBUS WALL.; unclear are MW and PW: PH. TRILOBUS.
- (3) Ph. Adenanthus G. F. W. Meyer (Chopra and KB 1, p. 799-800 : aranyamudga) = Ph. Rostratus Wall. (KB 1, p. 799-800).

$vanayam\bar{a}n\bar{\imath}$

(1) Seseli indicum Wight et Arn. (Avk; Chopra; Dutt; Kav; KB 2, p. 1209-1210; Nadk. 1, nr. 2289; V 1, p. 21; Vśs) = Cnidium diffusum DC. (Kav; MW) = Ligusticum diffusum Roxb. ex Sm. (KB 2, p. 1209-1210; Vśs).

$v\bar{a}r\bar{a}h\bar{i}kanda$

(1) Tacca integrifolia Ker-Gawl = T. aspera Roxb. (Bpn, p. 386-387; Chopra; DWH 3, p. 549-550; KB 4, p. 2476-2477);