

## Ordinary differential equations

A differential equation is an equation where the unknown (or unknowns) is a function  $f$ , and the equation **relates values of  $f$  at a point  $x$**  with values of derivatives of the function at **the same point  $x$** . If the function has one variable only (as is the case in this chapter), one speaks of ordinary differential equations.

**Theorem 2.1.6:**  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a differentiable function of two variables. Let  $x_0 \in \mathbb{R}$  and  $y_0 \in \mathbb{R}^2$ . Then the ODE  $y' = F(x, y)$  has an **unique** solution  $f$  defined on a *largest* open interval  $I$  containing  $x_0$  such that  $f(x_0) = y_0$ .

$\exists f : I : \mathbb{R} \text{ s.t. } \forall x \in I : f'(x) = F(x, y)$  and one cannot find a larger interval containing  $I$  with such a solution.

### Separation of variables:

If the ODE can be rewritten as  $\frac{dy}{dx} = f(x)g(y)$ , then

$$\int \frac{dy}{g(y)} = \int f(x)dx.$$

## Linear differential equations

**Definition 2.2.1:** Let  $I \subset \mathbb{R}$  open interval and  $k \geq 1$  an integer.

Homogeneous ODE of order  $k$ :  $y^{(k)} + a_{k-1}y^{(k-1)} + \dots + a_1y' + a_0y = 0$  where the coefficients  $a_0, \dots, a_{k-1}$  are complex-valued functions on  $I$ .

Inhomogeneous ODE of order  $k$ :  $y^{(k)} + a_{k-1}y^{(k-1)} + \dots + a_1y' + a_0y = b$  where  $b : I \rightarrow \mathbb{C}$ .

**Theorem 2.2.3:** Let  $I \subset \mathbb{R}$  an open interval and  $k \geq 1$  an integer,

$y^{(k)} + a_{k-1}y^{(k-1)} + \dots + a_1y' + a_0y = 0$  a linear ODE over  $I$  with continuous coefficients.

1. Set  $S$  of  $k$ -times differentiable solutions  $f : I \rightarrow \mathbb{C}$  is a **complex vector space** which is a subspace of complex-valued functions on  $I$ .

If the functions  $a_i$  are real-valued, the set of real-valued solutions is also a vector space.

2. The dimension of  $S$  is  $k$ , and for any choice of  $x_0 \in I$  and any  $(y_0, \dots, y_{k-1}) \in \mathbb{C}^k$ , there exists a unique  $f \in \mathbb{C}$  such that  $f(x_0) = y_0, f'(x_0) = y_1, \dots, f^{(k-1)}(x_0) = y_{k-1}$ .

3. Let  $b$  be a continuous function on  $I$ . There exists a solution  $f_0$  to the inhomogeneous equation and the set  $S_b$  is the set of functions  **$f + f_0$**  where  $f \in S$ .
4. For any  $x_0 \in I$  and any  $(y_0, \dots, y_{k-1}) \in \mathbb{C}^k$ , there exists a unique  $f \in S_b$  such that  $f(x_0) = y_0, f'(x_0) = y_1, \dots, f^{(k-1)}(x_0) = y_{k-1}$ .

**If  $b \neq 0$ , the set  $S_b$  is not a vector space!**

## Linear differential equations of order 1

Let  $I \subset \mathbb{R}$  be an open interval. We consider here the linear differential equation  **$y' + ay = b$**  when  $a$  and  $b$  are general continuous functions defined on  $I$ .

Steps to solve:

1. Solve the **homogeneous equation**:  $y' + ay = 0$  and obtain  $S$ .
2. Find solution  $f_0$  to the **inhomogeneous equation**.
3.  $S_b$  will contain  $f_0 + f$  where  $f \in S$  and if some  $f_1$  is a basis of  $S$  then the solutions are given by  $f_0 + zf_1$  where  $z \in \mathbb{C}$  are arbitrary.

If the initial value  $f(x_0) = y_0$  is given, then one must solve  $f_0(x_0) + zf_1(x_0) = y_0$  and determine the value of  $z$ .

**Proposition 2.3.1:** Any solution of  **$y' + ay = 0$**  is of the form  **$f(x) = z \exp(-A(x))$**  where  $A$  is a primitive of  $a$ . The unique solution with  $f(x_0) = y_0$  is  **$f(x) = y_0 \exp(A(x_0) - A(x))$** .

## Linear differential equations with constant coefficients

Now let  $k \geq 1$  an integer;  $a_0, \dots, a_{k-1}$  constant coefficients and  $b$  a continuous function. We consider the equation  $y^{(k)} + a_{k-1}y^{(k-1)} + \dots + a_1y' + a_0y = b$ .

### Solution of the homogeneous equation:

Let  **$P(\lambda) = \lambda^k + a_{k-1}\lambda^{k-1} + \dots + a_1\lambda + a_0$** . Find the **roots** of this polynomial. If the roots are not real but the coefficients are, express the solution in terms of sin and cos using  $e^{ix} = \cos x + i \sin x$ .

### Case 1: No multiple roots

Any solution  $f \in S$  of solutions of the homo equation is of the form  **$f(x) = z_1 e^{\alpha_1 x} + \dots + z_k e^{\alpha_k x}$**  for arbitrary  $z_1, \dots, z_k$ . To find an unique solution to  $f(x_0) = y_0, \dots, f^{(k-1)}(x_0) = y_{k-1}$  for given  $(y_0, \dots, y_{k-1})$  just view  $z_1, \dots, z_k$  as unknowns.

To obtain the **real valued solutions** if the coefficients are real:

$$f(x) = x_1 e^{\alpha_1 x} + \dots + x_m e^{\alpha_m x} + x_{m+1} e^{\alpha_{m+1} x} \cos(b_{m+1} x) + y_{m+1} e^{\alpha_{m+1} x} \sin(b_{m+1} x) + \dots + x_k e^{\alpha_k x} \cos(b_k x) + y_k e^{\alpha_k x} \sin(b_k x)$$

with  $\alpha_1, \dots, \alpha_m$  being the real solutions and  $\alpha_{m+1}, \dots, \alpha_k$  the complex solutions  $\alpha_j = a_j + ib_j$ .

### Case 2: Multiple roots

Assume  $\alpha$  is a multiple root of order  $j$  of  $P$  then the solutions look as follows:

$$f_{\alpha,0}(x) = e^{\alpha x}, \quad f_{\alpha,1}(x) = x e^{\alpha x}, \quad \dots, \quad f_{\alpha,j-1}(x) = x^{j-1} e^{\alpha x}$$

That is, multiply the original solution from *Case 1* with  $x^{i-1}$  for  $i = 1, \dots, j$ .

### Solution to the inhomogeneous equation:

1. **Ansatz method** (left side is  $b(x)$  and the right side is Ansatz)

$$a e^{\alpha x} \rightarrow b e^{\alpha x}$$

$$a \sin(\beta x) \text{ or } a \cos(\beta x) \rightarrow c \sin(\beta x) + d \cos(\beta x)$$

$$a e^{\sin(\beta x)} \text{ or } a e^{\cos(\beta x)} \rightarrow e^{\alpha x} [c \sin(\beta x) + d \cos(\beta x)]$$

$$P_n(x) e^{\alpha x} \rightarrow R_n(x) e^{\alpha x}$$

$$P_n e^{\alpha x} \sin(\beta x) \text{ or } P_n e^{\alpha x} \cos(\beta x) \rightarrow e^{\alpha x} [R_n \sin(\beta x) + S_n \cos(\beta x)]$$

With  $P_n(x), R_n(x), Q_n(x), S_n(x)$  being polynomials of degree  $n$ .

If  $b(x)$  is a linear combination of the above functions, then one should try the corresponding linear combination of the *Ansatz* functions.

If  $\lambda = \alpha + \beta i$  is a root of  $P(\lambda)$  of multiplicity  $m$ , then the *Ansatz* function should be multiplied by  $x^m$  (otherwise the *Ansatz* would solve the homo solution again)

### 2. Variation of constants

Assume  $(f_1, \dots, f_k)$  is the basis of the space  $S$  of solutions of the homogeneous equation. Now we search for a solution of the **inhomogeneous equation** of the form  $f(x) = z_1(x)f_1(x) + \dots + z_k(x)f_k(x)$  and impose the following **conditions**:

$$\begin{cases} z'_1(x)f_1(x) + \cdots + z'_k(x)f_k(x) = 0 \\ z'_1(x)f'_1(x) + \cdots + z'_k(x)f'_k(x) = 0 \\ \cdots \\ z'_1(x)f_1^{(k-2)}(x) + \cdots + z'_k(x)f_k^{(k-2)}(x) = 0 \end{cases}$$

## Differential calculus in $\mathbb{R}^n$

### Continuity in $\mathbb{R}^n$

The norm  $\|x\|$  satisfies  $\|x\| > 0$ ,  $\|x\| = 0 \Leftrightarrow x = 0$ ,  $\|tx\| = |t|\|x\|$  for all  $t \in \mathbb{R}$ , and  $\|x + y\| \leq \|x\| + \|y\|$  (triangle inequality).

**Definition 3.2.1:** Let  $(x_k)_{k \in \mathbb{N}}$  where  $x_k \in \mathbb{R}^n$ .  $x_k = (x_{k,1}, \dots, x_{k,n})$ . Let  $y = (y_1, \dots, y_n)$ . We say that the sequence  $(x_k)$  **converges to  $y$**  as  $y \rightarrow +\infty$  if for all  $\epsilon > 0$ , there exists  $N \geq 1$  such that for all  $n \geq N$  we have  $\|x_k - y\| < \epsilon$ .

**Lemma 3.2.2:** The sequence  $(x_k)$  **converges to  $y$**  as  $y \rightarrow +\infty \Leftrightarrow$  one of the following holds:

1. For each  $1 \leq i \leq n$ , the sequence  $(x_{k,i})$  of real numbers **converges to  $y_i$** .
2. The sequence of real numbers  $\|x_k - y\|$  **converges to 0** as  $y \rightarrow +\infty$ .

**Definition 3.2.3:** Let  $X \subset \mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}^m$

1. Let  $x_0 \in X$ .  $f$  is continuous at  $x_0$  if for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that, if  $x \in X$  satisfies  $\|x - x_0\| < \delta$ , then  $\|f(x) - f(x_0)\| < \epsilon$ .
2.  $f$  is continuous on  $X$  if it is continuous at all  $x_0 \in X$ .

**Proposition 3.2.4:** Let  $X \subset \mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}^m$ . Let  $x_0 \in X$ .  $f$  is **continuous at  $x_0$**   $\Leftrightarrow$  for **every sequence**  $(x_k)_{k \geq 1}$  in  $X$  such that  $x_k \rightarrow x_0$  as  $k \rightarrow +\infty$ , the sequence  $(f(x_k))_{k \geq 1}$  in  $\mathbb{R}^m$  **converges to  $f(x_0)$** .

**Definition 3.5.5:** Let  $X \subset \mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}^m$ . Let  $x_0 \in X$  and  $y \in \mathbb{R}^m$ .  $f$  has a limit  $y$  as  $x \rightarrow x_0$  with  $x \neq x_0$  if for every  $\epsilon > 0$ , there exists  $\delta > 0$ , such that for all  $x \in X$ ,  $x \neq x_0$ , such that  $\|x - x_0\| < \delta : \|f(x) - y\| < \epsilon$ . Then  $\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) = y$ .

**Proposition 3.2.7:** Let  $X \subset \mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}^m$ . Let  $x_0 \in X$  and  $y \in \mathbb{R}^m$ .

$\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) = y \Leftrightarrow$  for **every sequence**  $(x_k)$  in  $X$  such that  $x_k \rightarrow x$  as  $k \rightarrow +\infty$ , and  $x_k \neq x_0$ , the sequence  $(f(x_k))_{k \geq 1}$  in  $\mathbb{R}^m$  **converges to  $y$** .

**Proposition 2.3.9:** Let  $X \subset \mathbb{R}^n$ ,  $Y \subset \mathbb{R}^m$  and  $p \geq 1$  an integer. Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow \mathbb{R}^l$  be **continuous functions**. Then the composition  $f \circ g$  is **continuous**.

**Cartesian product, linear maps, multiplication and addition of continuous functions are continuous.**

If  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is continuous then so is the function  $g$  defined by  $g(x) = f(x, y_0)$  for a fixed  $y_0$ . The converse is not true.

**Definition 3.2.11:** A subset  $X \subset \mathbb{R}^n$  is

1. **Bounded** if the set of  $\|x\|$  for  $x \in X$  is bounded in  $\mathbb{R}$ .
2. **Closed** if for every sequence  $(x_k)$  in  $X$  that converges in  $\mathbb{R}^n$  to some vector  $y \in \mathbb{R}^n$ , we have  $y \in X$ .
3. **Compact** if it is bounded and closed.

$\{x \in \mathbb{R}^n : \|x - x_0\| = r, r \geq 0\}$  is closed (same for  $\mathbb{R}^3$ ),  $\{x \in \mathbb{R}^n : \|f(x)\| \leq r, r \geq 0\}$  is closed. **The union of open sets is open.**

**Proposition 3.2.13:** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a continuous map. For any closed set  $Y \subset \mathbb{R}^m$ , the set  $f^{-1}(Y) = \{x \in \mathbb{R}^n : f(x) \in Y\} \subset \mathbb{R}^n$  is closed.

**Theorem 3.2.15:** Let  $X \subset \mathbb{R}^n$ , **non empty compact set** and  $f : X \rightarrow Y$  a **continuous function**. Then  $f$  is **bounded and achieves its maximum and minimum** ( $\exists x_+$  and  $x_-$  in  $X$  such that  $f(x_+) = \sup_x \in X$  and  $f(x_-) = \inf_x \in X$ ).

### Partial derivatives

**Definition 3.3.1:** A subset  $X \subset \mathbb{R}^n$  is **open** if, for any  $x = (x_1, \dots, x_n) \in X$ ,  $\exists \delta > 0$  such that the set  $\{y = (y_1, \dots, y_n) \in \mathbb{R}^n : |x_i - y_i| < \delta \text{ for all } i\}$  is **contained in  $X$** .

**Proposition 3.3.2:** A set  $X \subset \mathbb{R}^n$  is **open**  $\Leftrightarrow$  the **complement**  $Y = \{x \in \mathbb{R}^n : x \notin X\}$  is **closed**.

**Corollary 3.3.3:** If  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is continuous and  $Y \subset \mathbb{R}^m$  is open, then  $f^{-1}(Y)$  is open in  $\mathbb{R}^n$ .

**Definition 3.3.5:** Let  $X \subset \mathbb{R}^n$  be an **open set**. Let  $f : X \rightarrow \mathbb{R}^m$ . Let  $1 \leq i \leq n$ .  $f$  has a **partial derivative on  $X$  with respect to the  $i$ -th variable** if for all  $x_0 = (x_{0,1}, \dots, x_{0,n}) \in X$ , the function defined by  $g(t) = f(x_{0,1}, \dots, x_{0,i-1}, t, x_{0,i+1}, \dots, x_{0,n})$  on the set  $I = \{t \in \mathbb{R} : (x_{0,1}, \dots, x_{0,i-1}, t, x_{0,i+1}, \dots, x_{0,n}) \in X\}$  is dif-

**ferentiable at  $t = x_{0,i}$** . Its derivative  $g'(x_{0,i})$  is denoted  $\frac{\partial f}{\partial x_i}(x_0)$ ,  $\partial_{x_i} f(x_0)$ ,  $\partial_i f(x_0)$ .  $g'(t) = (g'_1(t), \dots, g'_n(t))$ .

**Proposition 3.3.7:**  $X \subset \mathbb{R}^n$  **open** and  $f, g$  functions from  $X$  to  $\mathbb{R}^n$ .  $1 \leq i \leq n$ . If  $f$  and  $g$  have **partial derivatives with respect to the  $i$ -th coordinate on  $X$** , then:

1.  **$f + g$  also does**, and  $\partial_{x_i}(f + g) = \partial_{x_i}f + \partial_{x_i}g$
2. **if  $m = 1$ ,  $fg$  also does** and  $\partial_{x_i}(fg) = \partial_{x_i}f \cdot g + f \cdot \partial_{x_i}g$ . Furthermore, if  **$g(x) \neq 0$  for all  $x \in X$** , then  **$\frac{f}{g}$  has a partial derivative** with respect to the  $i$ -th coordinate on  $X$ , with  $\partial_{x_i}(\frac{f}{g}) = \frac{\partial_{x_i}f \cdot g - f \cdot \partial_{x_i}g}{g^2}$ .

**Definition 3.3.9:** Let  $X \subset \mathbb{R}^n$  be an **open set** and  $f : X \rightarrow \mathbb{R}^m$  with partial derivatives on  $X$ .  $f(x) = (f_1(x), \dots, f_m(x))$ . For any  $x \in X$ , the matrix  $J_f(x) = (\partial_{x_j} f_i(x))_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$  is the **Jacobi matrix** of  $f$  at  $x$ .

**Definition 3.3.11:** Let  $X \subset \mathbb{R}^n$  be an **open set** and  $f : X \rightarrow \mathbb{R}$ . If **all partial derivatives** of  $f$  exists at  $x_0$ , then the column vector  $\nabla f(x_0) = \begin{pmatrix} \partial_{x_1} f(x_0) \\ \vdots \\ \partial_{x_n} f(x_0) \end{pmatrix}$  is the **Gradient of  $f$  at  $x_0$** . The Jacobi matrix for  $m = 1$ .

### The differential

**Definition 3.4.3:** Let  $X \subset \mathbb{R}^n$  be an **open set** and  $f : X \rightarrow \mathbb{R}^m$ . Let  $u$  be a linear map  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $x_0 \in X$ .  $f$  is **differentiable at  $x_0$  with differential  $u$**  if

$$\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} \frac{f(x) - f(x_0) - u(x - x_0)}{\|x - x_0\|} = 0$$

where the limit is in  $\mathbb{R}^m$ . Denote  **$df(x_0) = u$** . If  $f$  is differentiable at every  $x \in X$  then  $f$  is differentiable on  $X$ . **The existence of partial derivatives or directional derivatives does not guarantee continuity (only the existence of all partial derivatives and their continuity does).**

**Proposition 3.4.4:** Let  $X \subset \mathbb{R}^n$  be an **open set** and  $f : X \rightarrow \mathbb{R}^m$  **differentiable on  $X$** . Then

1.  $f$  is **continuous on  $X$** .
2.  $f$  admits **partial derivatives** with respect to each variable.

3. If  $m = 1$ , let  $x_0 \in X$  and  $u(x_1, \dots, x_n) = a_1 x_1, \dots, a_n x_n$  the differential of  $f$  at  $x_0$ . Then  $\partial x_i f(x_0) = a_i$  for  $1 \leq i \leq n$ .

Let  $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h(x, y) = (f(x, y), g(x, y))$  and  $m(u, v) = uv$ , so that  $m \circ h(x, y) = f(x, y)g(x, y)$ . Therefore  $\frac{\partial(fg)}{\partial x} = x \partial_x f + u \partial_x g$ .

Let  $f : I \rightarrow \mathbb{R}^m$  and  $g : \mathbb{R}^m \rightarrow \mathbb{R}$ ,  $f$  differentiable on  $I$  and  $g$  differentiable on  $\mathbb{R}^m$ , then  $g \circ f$  is differentiable on  $I$  and  $(g \circ f)'(t) = dg(f(t))f'(t) = \nabla g(f(t)) \cdot f'(t)$ .

**Proposition 3.4.6:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$ ,  $g : X \rightarrow \mathbb{R}^m$  differentiable on  $X$ .

1.  $f + g$  is differentiable with differential  $d(f + g) = df + dg$  and if  $m = 1$  then  $f, g$  differentiable.
2. If  $m = 1$  and  $g(x) \neq 0$  for all  $x \in X$ , then  $\frac{f}{g}$  is differentiable.

**Proposition 3.4.7:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$ . If  $f$  has all partial derivatives on  $X$ , and if the partial derivatives are continuous on  $X$ , then  $f$  is differentiable on  $X$ . With  $df(x_0) = J_f(x_0)$ .

**Proposition 3.4.9:** Let  $X \subset \mathbb{R}^n$  and  $Y \subset \mathbb{R}^m$  be both open and  $f : X \rightarrow Y$ ,  $g : Y \rightarrow \mathbb{R}^p$  differentiable. Then  $g \circ f : X \rightarrow \mathbb{R}^p$  is differentiable on  $X$  and for any  $x_0 \in X$ , its differential is given by the composition  $d(g \circ f)(x_0) = dg(f(x_0)) \circ df(x_0)$ , in particular  $J_{g \circ f}(x_0) = J_g(f(x_0))J_f(x_0)$ .

**Definition 3.4.11:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$  differentiable. Let  $x_0 \in X$  and  $u = df(x_0)$ . The graph of the affine linear approximation  $g(x) = f(x_0) + u(x - x_0)$  from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  is called the **tangent space at  $x_0$  to the graph of  $f$** .

**Definition 3.4.13:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$ . Let  $v \in \mathbb{R}^n$  be non-zero vector and  $x_0 \in X$ . We say that  $f$  has a **directional derivative  $w \in \mathbb{R}^m$  in the direction  $v$** , if  $g$  defined on the set  $I = \{t \in \mathbb{R} : x_0 + tv \in X\}$  by  $g(t) = f(x_0 + tv)$  has a derivative at  $t = 0$ , and this is equal to  $w$ .

In other words:  $\lim_{\substack{t \rightarrow 0 \\ t \neq 0}} \frac{f(x_0 + tv) - f(x_0)}{t} = w$

**Computing:** Let  $\varphi(t) = f(x + tv)$ , then compute  $\varphi'(0)$ . If it doesn't exist, then the directional derivative does not exist.

**Proposition 3.4.15:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$  differentiable. Then for any  $x \in X$  and non-zero

$v \in \mathbb{R}^n$ ,  $f$  has a **directional derivative at  $x_0$  in the direction  $v$  equal to  $df(x_0)(v)$** . Suppose  $m = 1$ . Let  $f : X \rightarrow \mathbb{R}$  be differentiable and  $x_0 \in X$ . The **tangent space at  $x_0$**  to the graph of  $f$  is the set of  $(x, y) \in \mathbb{R}^n \times \mathbb{R}$  such that  $y = f(x_0) + \nabla f(x_0) \cdot (x - x_0)$ . The vector  $\nabla f(x_0)$  points to the direction of greatest increase.

The **gradient is perpendicular to the level sets** determined by an equation of the form  $f(x) = c$ .

$$0 = (f \circ \gamma)'(0) = \nabla f(x_0) \cdot \gamma'(0).$$

$$D_v f(\gamma(t)) = df(\gamma(t))v = \nabla f(\gamma(t)) \cdot v.$$

$\nabla f(x_0) \cdot v = \|\nabla f(x_0)\| \cos(\theta)$  where  $\theta$  is the angle between the gradient and the direction  $v$ .

## Higher derivatives

**Definition 3.5.1:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m$ .  $f$  is of class  $C^1$  if  $f$  is differentiable on  $X$  and all its partial derivatives are continuous. The set of functions of class  $C^1$  from  $X$  to  $\mathbb{R}^m$  is denoted  $C^1(X; \mathbb{R}^m)$ . Let  $k \geq 2$ , then  $f$  is of class  $C^k$  if it is differentiable and each partial derivative  $\partial x_i f : X \rightarrow \mathbb{R}^m$  is of class  $C^{k-1}$ . If  $f \in C^k(C; \mathbb{R}^m)$  for all  $k \geq 1$ , then  $f \in C^\infty$ .

**Proposition 3.5.4:** Let  $k \geq 2$ ,  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}^m \in C^k$ . Then the partial derivatives of order  $k$  are independent of the order in which the partial derivatives were taken: For any  $x$  and  $y$ ,  $\partial x_i y_j = \partial y_j x_i$ . Same for more variables.

**Definition 3.5.8:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R} \in C^2$ . For  $x \in X$ , the **Hessian matrix** of  $f$  at  $x$  is the **symmetric square matrix**  $\text{Hess}_f(x) = (\partial_{x_i, x_j}^2 f)_{\substack{1 \leq i \\ j \leq n}} = H_f(x)$ .

$$\text{Hess}_f(x) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \cdots & \frac{\partial^2 f}{\partial x_1 x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n x_1} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

## Change of variable

We have an open set  $U \subset \mathbb{R}^n$  with  $(y_1, \dots, y_n)$  the new variables and a change of variable  $g : U \rightarrow X$  which expresses the variables  $(x_1, \dots, x_n)$  in terms of the new variables. For a given function  $f : X \rightarrow \mathbb{R}$ , the composite  $h = f \circ g : U \rightarrow \mathbb{R}$  is  $f$  expressed in terms of the new variables  $y$ .

$$\partial_{y_1} h = \frac{\partial f}{\partial x_1} \frac{\partial g_1}{\partial y_1} + \cdots + \frac{\partial f}{\partial x_n} \frac{\partial g_n}{\partial y_1}$$

Some common notation:  $\partial_{y_1} h = \partial_{y_1} f$  since they are the same function with different coordinate systems.  $g_i$  is usually also replaced with  $x_i$ , s.t.  $\partial_{y_1} f = \frac{\partial f}{\partial x_1} \frac{\partial g_1}{\partial y_1} + \cdots + \frac{\partial f}{\partial x_n} \frac{\partial g_n}{\partial y_1} = \frac{\partial f}{\partial x_1} \frac{\partial x_1}{\partial y_1} + \cdots + \frac{\partial f}{\partial x_n} \frac{\partial x_n}{\partial y_1}$

## Taylor polynomials

**Definition 3.7.1:** Let  $k \geq 1$  be an integer and  $f : X \rightarrow \mathbb{R}$  a function of class  $C^k$  on  $X$  and some fix  $x_0 \in X$ . The  $k$ -th Taylor polynomial of  $f$  at  $x_0$  is the polynomial in  $n$  variables of degree  $\leq k$  given by

$$T_k f(y; x_0) = f(x_0) + \sum_{i=1}^n \frac{\partial f}{\partial x_i} y_i + \cdots + \sum_{m_1 + \cdots + m_n = k} \frac{1}{m_1! \cdots m_n!} \frac{\partial^k f}{\partial x_1^{m_1} \cdots \partial x_n^{m_n}}(x_0) y_1^{m_1} \cdots y_n^{m_n},$$

such that for  $y = x - x_0 : f(x) = T_k f(x - x_0; x_0) + (\text{remainder})$ .

$$\frac{1}{2} \sum_{i=1}^n \frac{\partial^2 f}{\partial x_i^2}(x_0) y_i^2 + \sum_{1 \leq i < j \leq n} \frac{\partial^2 f}{\partial x_i \partial x_j}(x_0) y_i y_j = \frac{1}{2} y^t \text{Hess}_f(x_0) y \text{ hence}$$

$$T_2 f(y; x_0) = f(x_0) + \nabla f(x_0) \cdot y + \frac{1}{2} y^t \text{Hess}_f(x_0) \cdot y,$$

for  $y \in \mathbb{R}$ .

**Proposition 3.7.3:** Let  $k \geq 1$  be an integer,  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}$  a function of class  $C^k$ . For  $x_0 \in X$ , if we define  $E_k f(x; x_0)$  by  $f(x) = T_k f(x - x_0; x_0) + E_k f(x; x_0)$  then  $\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} \frac{E_k f(x; x_0)}{\|x - x_0\|^k} = 0$ .

## Critical points

**Proposition 3.8.1:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}$  differentiable. If  $x_0 \in X$  s.t.

$f(y) \leq f(x_0)$  for all  $y$  close enough to  $x_0$  (local maximum at  $x_0$ )

or  $f(y) \geq f(x_0)$  for all  $y$  close enough to  $x_0$  (local minimum at  $x_0$ )

then  $df(x_0) = 0$ ,  $\nabla f(x_0) = 0$  or  $\partial_{x_i} f(x_0) = 0$  equivalently for  $1 \leq i \leq n$ .



**Definition 3.8.2:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}$  differentiable. A point  $x_0 \in X$  s.t.  $\nabla f(x_0) = \mathbf{0}$  is called a **critical point** of  $f$ .

When  $f$  is defined on a compact set  $\bar{X} = X \cup B$  with  $X$  an open set and  $B$  the boundary, then the maxima and minima should also be explicitly **computed at the boundary**.

**Definition 3.8.6:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}$  of class  $C^2$ . A critical point  $x_0 \in X$  of  $f$  is **non-degenerate** if the **Hessian matrix has non-zero determinant**.

**Corollary 3.8.7:** Let  $X \subset \mathbb{R}^n$  be open and  $f : X \rightarrow \mathbb{R}$  of class  $C^2$ ,  $x_0 \in X$  a non-degenerate critical point of  $f$ . Let  $p$  and  $q$  be the positive and negative eigenvalues of  $\text{Hess}_f(x_0)$ .

1. If  $p = n$ ,  $f$  has a local **minimum** at  $x_0$ .
2. If  $q = n$ ,  $f$  has a local **maximum** at  $x_0$ .
3. Otherwise,  $f$  has a **saddle point** at  $x_0$ .

$p = n \Leftrightarrow H_f(x_0)$  is positive definite.

$A = (a_{i,j})_{\substack{1 \leq i \\ j \leq n}}^1 \leq i$  is **positive definite**  $\Leftrightarrow \det(A_k) > 0$  for  $1 \leq k \leq n$ .  $A$  is **negative definite**  $\Leftrightarrow (-1)^k \det(A_k) > 0$  for  $1 \leq k \leq n$ .

## Integration in $\mathbb{R}^n$

### Line integrals

**Definition 4.1.1:**

1. Let  $I = [a, b]$  a closed and bounded interval in  $\mathbb{R}$ . Let  $f(t) = (f_1(t), \dots, f_n(t))$  continuous from  $I$  to  $\mathbb{R}^n$ . Then we define  $\int_a^b f(t) dt = (\int_a^b f_1(t) dt, \dots, \int_a^b f_n(t) dt) \in \mathbb{R}^n$ .
2. A **parameterised curve** in  $\mathbb{R}^n$  is a continuous map  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  that is piecewise  $C^1$ .
3. Let  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  be a parameterised curve. Let  $X \subset \mathbb{R}^n$  be a subset containing the image of  $\gamma$ , and let  $f : X \rightarrow \mathbb{R}^n$  continuous.

$$\int_a^b f(\gamma(t)) \cdot \gamma'(t) dt \in \mathbb{R}$$

is called the line integral of  $f$  along  $\gamma$ . It is denoted  $\int_\gamma f(s) ds$  or  $\int_\gamma f(s) d\vec{s}$ .

For  $X \subset \mathbb{R}^n$ ,  $f : X \rightarrow \mathbb{R}^n$  is usually called a **vector field**.

**Definition 4.1.4:** Let  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  be a parameterised curve. An oriented reparametrisation of  $\gamma$  is a parame-

terised curve  $\sigma : [c, d] \rightarrow \mathbb{R}^n$  such that  $\sigma = \gamma \circ \varphi$  where  $\varphi : [c, d] \rightarrow [a, b]$  is a continuous map, differentiable on  $]a, b[$ , that is strictly increasing and satisfies  $\varphi(a) = c$  and  $\varphi(b) = d$ .

**Proposition 4.1.5:** Let  $\gamma$  be a **parameterised curve** in  $\mathbb{R}^n$  and  $\sigma$  an **oriented reparametrisation** of  $\gamma$ . Let  $X$  be a set containing the image of  $\gamma$ , or equivalently the image of  $\sigma$ , and  $f : X \rightarrow \mathbb{R}^n$  continuous. Then we have  $\int_\gamma f(s) \cdot d\vec{s} = \int_\sigma f(s) \cdot d\vec{s}$ .

**Definition 4.1.8:** Let  $X \subset \mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}^n$  a **continuous vector field**. If for any  $x_1, x_2$  in  $X$ , the **line integral**  $\int_\gamma f(c) \cdot d\vec{s}$  is **independent of the choice of a parameterised curve**  $\gamma$  in  $X$  from  $x_1$  to  $x_2$ , then we say that the vector field is **conservative**.

Equivalently,  $f$  is conservative  $\Leftrightarrow \int_\gamma f(s) \cdot d\vec{s} = \mathbf{0}$  for any **closed** parameterised curve in  $X$ . Where a curve is closed if  $\gamma(a) = \gamma(b)$ .

**Theorem 4.1.10:** Let  $X$  be an open set and  $f$  a **conservative** vector field. Then there exists a  $C^1$  function  $g$  on  $X$  such that  $f = \nabla g$ . If any two points of  $X$  are conjoined by a parameterised curve, then  $g$  is unique up to addition of a constant: if  $\nabla g_1 = f$ , then  $g - g_1$  is constant on  $X$ .

Any two points  $x$  and  $y$  on  $X$  can be **joined by a parameterised curve**  $\Leftrightarrow \exists \gamma : [a, b] \rightarrow X$  such that  $\gamma(a) = x$  and  $\gamma(b) = y$ . Then  $X$  is **path connected**. This is also true whenever  $X$  is convex. If  $f$  is conservative, then a function  $g$  such that  $\nabla g = f$  is called a **potential for  $f$** .

$$\int_a^b f(\gamma(t)) \cdot \gamma'(t) dt \in \mathbb{R} = g(\gamma(b)) - g(\gamma(a))$$

A set  $X$  is **convex** if for any  $x, y \in X$ , the line segment joining  $x$  to  $y$  is contained in  $X$ .

**Proposition 4.1.13:** Let  $X \subset \mathbb{R}^n$  be an open set and  $f : X \rightarrow \mathbb{R}^n$  of class  $C^1$ .

$f(x) = (f_1(x), \dots, f_n(x))$ . If  $f$  is **conservative**, then we have  $\frac{\partial f_i}{\partial x_j} = \frac{\partial f_j}{\partial x_i}$  for any integers with  $1 \leq i \neq j \leq n$ .

**Definition 4.1.15:** A subset  $X \subset \mathbb{R}^n$  is **star-shaped** if there exists  $x_0 \in X$  such that, for all  $x \in X$ , the **line segment joining  $x_0$  to  $x$  is contained in  $X$** . We then say that  $X$  is star-shaped around  $x_0$ .

**Theorem 4.1.17:** Let  $X$  be a **star-shaped** open subset of  $\mathbb{R}^n$  and  $f$  a  $C^1$  vector field such that  $\frac{\partial f_i}{\partial x_j} = \frac{\partial f_j}{\partial x_i}$  on  $X$

for all  $i \neq j$  between 1 and  $n$ . Then the vector field  $f$  is **conservative**.

The requirement that  $X$  is star-shaped is not necessary.  $X$  should have no “hole” in the middle around which a circle can go without it being possible to contract it within  $X$ .

**Definition 4.1.20:** Let  $X \subset \mathbb{R}^3$  be an open set and  $f : X \rightarrow \mathbb{R}^3$  a  $C^1$  **vector field**. Then the **curl** of  $f$ , denoted  $\text{curl}(f)$ , is the continuous vector field on  $X$  defined by

$$\text{curl}(f) = \begin{pmatrix} \partial_y f_3 - \partial_z f_2 \\ \partial_z f_1 - \partial_x f_3 \\ \partial_x f_2 - \partial_y f_1 \end{pmatrix}$$

where  $f(x, y, z) = (f_1(x, y, z), f_2(x, y, z), f_3(x, y, z))$ .

$$\text{curl}(f) = \begin{vmatrix} e_1 & e_2 & e_3 \\ \partial_x & \partial_y & \partial_z \\ f_1 & f_2 & f_3 \end{vmatrix}$$

with the rule that  $\partial_x \cdot f_i = f_i \cdot \partial_x = \partial_x f_i$ .

**curl( $\nabla f$ ) = 0.**

### The Riemann integral in $\mathbb{R}^n$

The goal is to define the integral  $\int_X f(x_1, \dots, x_n) dx$  for a closed and bounded set  $X$  and continuous function  $f$ , so that it has analogue properties to the Riemann integral for  $n = 1$ .

For a bounded and closed subset  $X \subset \mathbb{R}^n$  and any continuous function  $f : X \rightarrow \mathbb{R}$ , one can define the integral of  $f$  over  $X$  denoted  $\int_X f(x) dx$ , which is a real number.

The integral satisfies the following properties:

1. **Compatibility:** If  $n = 1$  and  $X = [a, b]$  with  $a \leq b$ , then

$$\int_{[a,b]} f(x) dx = \int_a^b f(x) dx$$

2. **Linearity:** If  $f$  and  $g$  are continuous on  $X$  and  $a, b \in \mathbb{R}$ , then

$$\int_X (af_1(x) + bf_2(x)) dx = a \int_X f_1(x) dx + b \int_X f_2(x) dx$$

3. **Positivity:** If  $f \leq g$ , then

$$\int_X f(x)dx \leq \int_X g(x)dx$$

and especially, if  $f \geq 0$ , then

$$\int_X f(x)dx \geq 0.$$

Moreover, if  $Y \subset X$  is compact and  $f \geq 0$ , then

$$\int_Y f(x)dx \leq \int_X f(x)dx.$$

4. **Upper bound and triangle inequality:** In particular, since  $-|f| \leq f \leq |f|$ , we have

$$\left| \int_X f(x)dx \right| \leq \int_X |f(x)|dx,$$

and since  $|f+g| \leq |f|+|g|$ , we have

$$\left| \int_X (f(x)+g(x))dx \right| \leq \int_X |f(x)|dx + \int_X |g(x)|dx.$$

5. **Volume:** If  $f = 1$ , then the integral of  $f$  is the volume in  $\mathbb{R}^n$  of the set  $X$ , and if  $f \geq 0$  in general, the integral of  $f$  is the volume of the set  $\{(x,y) \in X \times \mathbb{R} : 0 \leq y \leq f(x)\} \subset \mathbb{R}^{n+1}$ . In particular, if  $X$  is a bounded rectangle, say  $X = [a_1, b_1] \times \dots \times [a_n, b_n] \subset \mathbb{R}^n$  and if  $f = 1$ , then

$$\int_X dx = (b_n - a_n) \cdots (b_1 - a_1).$$

6. **Multiple integral or Fubini's Theorem:** If  $n_1$  and  $n_2$  are integers  $\geq 1$  such that  $n = n_1 + n_2$ , then for  $x_1 \in \mathbb{R}^{n_1}$ , let

$$Y_{x_1} = \{x_2 \in \mathbb{R}^{n_2} : (x_1, x_2) \in X\} \subset \mathbb{R}^{n_2}.$$

Let  $X_1$  be the set of  $x_1 \in \mathbb{R}^{n_1}$  such that  $Y_{x_1}$  is not empty. Then  $X_1$  is compact in  $\mathbb{R}^{n_1}$  and  $Y_{x_1}$  is compact in  $\mathbb{R}^{n_2}$  for all  $x_1 \in X_1$ . If the function

$$g(x_1) = \int_{Y_{x_1}} f(x_1, x_2)dx_2$$

on  $X_1$  is continuous, then

$$\int_X f(x_1, x_2)dx = \int_{X_1} g(x_1)dx_1$$

$$= \int_{X_1} \left( \int_{Y_{x_1}} f(x_1, x_2)dx_2 \right) dx_1.$$

Similarly, exchanging the role of  $x_1$  and  $x_2$ , we have

$$\int_X f(x_1, x_2)dx = \int_{X_2} \left( \int_{Z_{x_2}} f(x_1, x_2)dx_1 \right) dx_2$$

where  $Z_{x_2} = \{x_1 : (x_1, x_2) \in X\}$ , if the integral over  $x_1$  is a continuous function.

If the variables are  $x_1, \dots, x_n$  we also write  $\int_X f(x_1, \dots, x_n)dx_1, \dots, dx_n$

**Definition 4.2.3:**

1. Let  $1 \leq m \leq n$  be an integer. A parameterised  $m$ -set in  $\mathbb{R}^n$  is a **continuous map**  $f : [a_1, b_1] \times \dots \times [a_m, b_m] \rightarrow \mathbb{R}^n$  which is  $C^1$  on  $]a_1, b_1[ \times \dots \times ]a_m, b_m[$ .
2. A subset  $B \subset \mathbb{R}^n$  is **negligible** if there exists an integer  $k \geq 0$  and parameterised  $m_i$ -sets  $f_i : X_1 \rightarrow \mathbb{R}^n$ , with  $1 \leq i \leq k$  and  $m_i < n$ , such that  $X \subset f_1(X_1) \cup \dots \cup f_k(X_k)$ .

The **image** of a parameterised  $m$ -set  $f$  might be of **dimension smaller than  $m$**  (e.g. if  $f$  is constant, in which case the image is a single point, which is an object of dimension 0)

**Proposition 4.2.5:** Let  $X \subset \mathbb{R}^n$  be compact. Assume that  $X$  is **negligible**. Then for any continuous function on  $X$ , we have  $\int_X f(x)dx = 0$ .

**Improper integrals**

Let  $I \subset \mathbb{R}$  be a **bounded interval**,  $J = [a, +\infty[$  for some  $a \in \mathbb{R}$  and  $f$  a **continuous function** on  $X = J \times I$ . We say that it is **Riemann-integrable** on  $X$  if the limit

$$\begin{aligned} \lim_{x \rightarrow +\infty} \int_{[a, x] \times I} f(x, y)dx dy &= \lim_{x \rightarrow +\infty} \int_a^x \left( \int_I f(x, y)dy \right) dx \\ &= \lim_{x \rightarrow +\infty} \int_I \left( \int_a^x f(x, y)dx \right) dy \end{aligned}$$

exists. Denote this limit by

$$\int_{J \times I} f(x, y)dx dy.$$

Similarly let  $f$  be continuous on  $\mathbb{R}^2$ . Assume that  $f \geq 0$ .  $f$  is Riemann-integrable on  $\mathbb{R}^2$ , if the limit

$$\lim_{R \rightarrow +\infty} \int_{[-R, R]^2} f(x, y)dx dy$$

exists, which is called the integral of  $f$  over  $\mathbb{R}^2$  and denoted

$$\int_{\mathbb{R}^2} f(x, y)dx dy.$$

This integral is also the limit of

$$\int_{D_R} f(x, y)dx dy$$

where  $D_R$  is the disc of radius  $R$  centred at 0.

The easiest way to prove that a certain **improper integral exists**: If  $|f| \leq g$  (resp  $0 \leq f \leq g$ ), and

$$\int_{J \times I} g(x, y)dx dy$$

or

$$\int_{\mathbb{R}^2} g(x, y)dx dy$$

exists, then so does

$$\int_{J \times I} f(x, y)dx dy \text{ or } \int_{\mathbb{R}^2} f(x, y)dx dy.$$

**The change of variable formula**

Let  $\bar{X} \subset \mathbb{R}^n$  and  $\bar{Y} \subset \mathbb{R}^n$  be **compact subsets**,  $\varphi : \bar{X} \rightarrow \bar{Y}$  a **continuous map**. We assume that we can write  $\bar{X} = X \cup B, \bar{Y} = Y \cup C$  where

1.  $X$  and  $Y$  are **open**
2.  $B$  and  $C$  are **negligible**
3. the restriction of  $\varphi$  to the open set  $X$  is a  $C^1$  **bijective map from  $X$  to  $Y$** .

In this situation, the Jacobian matrix  $J_\varphi(x)$  is **invertible at all  $x \in X$** .

**Theorem 4.4.2:** In the situation described above, for any continuous function  $f$  on  $\bar{Y}$ , we have  $\int_{\bar{X}} f(\varphi) |\det(J_\varphi(x))| dx = \int_{\bar{Y}} f(y) dy$ .

**Cartesian to polar coordinates:**  $x = r \cos(\theta), y = r \sin(\theta)$ , then calculate all partial derivatives with respect to the new variables:

$\partial_r x = \cos \theta, \partial_r y = \sin \theta, \partial_\theta x = -r \sin \theta, \partial_\theta y = r \cos \theta$ . Then  $J_{(r, \theta)} = r, dx dy \rightarrow r dr d\theta$ .

Geometric applications of integrals

1. Centre of mass:

Let  $X$  be a compact subset of  $\mathbb{R}^n$ , such that the volume of  $X$  is positive. The centre of mass of  $X$  is the point  $\bar{x} \in \mathbb{R}^n$  such that  $\bar{x} = (\bar{x}_1, \dots, \bar{x}_n)$  with

$$\bar{x}_i = \frac{1}{\text{vol}(X)} \int_X x_i dx.$$

Intuitively,  $\bar{x}_i$  is the average over  $X$  of the  $i$ -th co-ordinate, and  $\bar{x}$  is the point where  $X$  is perfectly balanced.

Note that  $\bar{x}$  is not necessarily in  $X$ .

2. Surface area:

Consider the continuous function  $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$  which is  $C^1$  on  $]a, b[ \times ]c, d[$ . Let

$$\Gamma = \{(x, y, z) \in \mathbb{R}^3 : (x, y) \in [a, b] \times [c, d], z = f(x, y)\}$$

$\subset \mathbb{R}^3$  be the graph of  $f$ . This is a surface and should have an area, which is given by

$$\int_a^b \int_c^d \sqrt{1 + (\partial_x f(x, y))^2 + (\partial_y f(x, y))^2} dx dy.$$

3. Length of a curve:

$$f : [a, b] \rightarrow \mathbb{R} : \int_a^b \sqrt{1 + f'(x)^2} dx.$$

The Green formula

The Green formula concerns the case of relating an integral over a subset  $X$  of  $\mathbb{R}^2$  with a line integral over its boundary.

**Definition 4.6.1:** A simple closed parameterised curve  $\gamma : [a, b] \rightarrow \mathbb{R}^2$  is a closed parameterised curve such that  $\gamma(t) \neq \gamma(s)$  unless  $t = s$  or  $\{s, t\} = \{a, b\}$ , and such that  $\gamma'(t) \neq 0$  for  $a < t < b$ . If  $\gamma$  is only piecewise  $C^1$  inside  $]a, b[$ , this condition only applies to where  $\gamma'(t)$  exists.

**Theorem 4.6.3:** Let  $X \subset \mathbb{R}^2$  compact with a boundary  $\partial X$  that is the union of finitely many simple closed parameterised curves  $\gamma_1, \dots, \gamma_k$ . Assume that  $\gamma_i : [a_i, b_i] \rightarrow \mathbb{R}^2$  has the property that  $X$  lies always to the left of the tangent vector  $\gamma_i'(t)$  based at  $\gamma_i(t)$ . Let  $f = f(x, y)$  be a

vector field of class  $C^1$  defined on some open set containing  $X$ . Then we have

$$\int_X \left( \frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) dx dy = \sum_{i=1}^k \int_{\gamma_i} f \cdot d\vec{s}.$$

**Corollary 4.6.5:** Let  $X \subset \mathbb{R}^2$  compact with a boundary  $\partial X$  that is the union of finitely many simple closed parameterised curves  $\gamma_1, \dots, \gamma_k$ . Assume that  $\gamma_i : [a_i, b_i] \rightarrow \mathbb{R}^2$  has the property that  $X$  lies always to the left of the tangent vector  $\gamma_i'(t)$  based at  $\gamma_i(t)$ . Then we have

$$\text{Vol}(X) = \sum_{i=1}^k \int_{\gamma_i} x \cdot d\vec{s} = \sum_{i=1}^k \int_{a_i}^{b_i} \gamma_{i,1}(t) \gamma'_{i,2}(t) dt.$$

The Gauss-Ostrogradski formula

This formula is the analogue of the Green formula in  $\mathbb{R}^3$ .

**Definition 4.7.1:** A parameterised surface  $\Sigma : [a, b] \times [c, d] \rightarrow \mathbb{R}^3$  is a 2-set in  $\mathbb{R}^3$  such that the rank of the Jacobian matrix is 2 at all  $(s, t) \in ]a, b[ \times ]c, d[$ .

**Definition 4.7.3:** Let  $x$  and  $y$  be two linearly independent vectors in  $\mathbb{R}^3$ . The vector product, or cross product  $z = x \times y$  is the unique vector in  $\mathbb{R}^3$  such that  $(x, y, z)$  is a basis of  $\mathbb{R}^3$  with  $\det(x, y, z) > 0$ , and  $\|z\| = \|x\| \|y\| \sin(\theta)$ , where  $\theta$  is the angle between  $x$  and  $y$ .

$$x \times y = \begin{pmatrix} x_2 y_3 - x_3 y_2 \\ x_3 y_1 - x_1 y_3 \\ x_1 y_2 - x_2 y_1 \end{pmatrix} = \det \begin{vmatrix} e_1 & e_2 & e_3 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{vmatrix}$$

$$e_1 \times e_2 = e_3, \quad e_2 \times e_3 = e_1, \quad e_3 \times e_1 = e_2 \quad \text{and} \quad y \times x = -x \times y$$

**Theorem 4.7.6:** Let  $X \subset \mathbb{R}^3$  compact with a boundary  $\partial X$  that is a parameterised surface  $\Sigma : [a, b] \times [c, d] \rightarrow \mathbb{R}^3$ . Assume that  $\Sigma$  is injective in  $]a, b[ \times ]c, d[$ , and that  $\Sigma$  has the property that the normal vector  $\vec{n}$  points away from  $\Sigma$  at all points. Let  $\vec{u} = \frac{\vec{n}}{\|\vec{n}\|}$  be the unit exterior normal vector. Let  $f = (f_1, f_2, f_3)$  be a vector field of class  $C^1$  defined on some open set containing  $X$ . Then we have

$$\int_X \text{div}(f) dx dy dz = \int_{\Sigma} (f \cdot \vec{u}) d\sigma.$$

For a vector field  $f = (f_1, f_2, f_3)$  on  $X \subset \mathbb{R}^3$ , we denote the divergence of the field  $\text{div}(f) = \partial_x f + \partial_y f + \partial_z f$ .

For a parameterised surface  $\Sigma : [a, b] \times [c, d] \rightarrow \mathbb{R}^3$  with exterior normal vector field  $\vec{n} = (n_1, n_2, n_3) = \partial_s \Sigma \times \partial_t \Sigma$

and a function  $g$  defined on the image of  $\Sigma$ , we define the surface integral

$$\int_{\Sigma} g \, d\sigma = \int_a^b \int_c^d g(\Sigma(s, t)) \sigma(s, t) ds dt$$

where  $\sigma(s, t) = \|\partial_s \Sigma \times \partial_t \Sigma\| = \|\vec{n}(s, t)\|$ .

Like the integral for a parameterised curve, the surface integral is independent of the chosen parameterisation of the surface.

For a  $C^1$  vector field  $f = (f_1, f_2, f_3)$  on  $\mathbb{R}^3$ , we define

$$\int_{\Sigma} (f \cdot \vec{n}) d\sigma = \int_{\Sigma} g \, d\sigma,$$

where

$$g(\Sigma(s, t)) = f(\Sigma(s, t)) \cdot \vec{u}(s, t) = \sum_{i=1}^3 u_i(s, t) f_i(\Sigma(s, t)).$$

This is called the flux of the vector field  $f$  through the surface  $\Sigma$ .

In the flux,  $\vec{u}(s, t) \sigma(s, t) = \vec{n}(s, t)$ .

Examples and other

Basic rules that you will forget :)

Complex Numbers

$$e^{i\varphi} = \cos(\varphi) + i \sin(\varphi), \quad e^{i\varphi} = \text{cis}(\varphi), \quad |e^{i\varphi}| = 1$$

$z$	$z = x + iy \begin{cases} \text{x: Real} \\ \text{y: Imaginary} \end{cases}$	$z = r \cdot e^{i\varphi} = r \cdot \text{cis}(\varphi)$
$\bar{z}$	$\bar{z} = x - iy$	$z = r \cdot e^{-i\varphi}$
$ z $	$ z  = \sqrt{z \cdot \bar{z}} = \sqrt{x^2 + y^2}$	$ z  = r = \sqrt{z \cdot \bar{z}}$
$z_1 + z_2$	$x_1 + x_2 + i(y_1 + y_2)$	
$z_1 - z_2$	$x_1 - x_2 + i(y_1 - y_2)$	
$z_1 \cdot z_2$	$(x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1)$	$r_1 \cdot r_2 \cdot e^{i(\varphi_1 + \varphi_2)}$
$\frac{z_1 \cdot \bar{z}_2}{ z_2 ^2}, z_2 \neq 0$	$\frac{(x_1 x_2 + y_1 y_2) + i(x_2 y_1 - x_1 y_2)}{x_2^2 + y_2^2}$	$\frac{r_1}{r_2} \cdot e^{i(\varphi_1 - \varphi_2)}$
$\frac{1}{z}, z \neq 0$	$\frac{\bar{z}}{ z ^2} = \frac{(x - iy)}{x^2 + y^2}$	$\frac{1}{r} \cdot e^{-i\varphi}$

$z^n$	$r^n \cdot (\cos(n \cdot \varphi) + i \sin(n \cdot \varphi)) = r^n \cdot e^{in\varphi}$
$\sqrt[n]{z}$	$\sqrt[n]{r} \cdot (\cos(\frac{\varphi + 2\pi k}{n}) + i \sin(\frac{\varphi + 2\pi k}{n})), \quad k = 0, 1, \dots, n - 1$

$i^0$	$i^1$	$i^2$	$i^3$
1	$i$	-1	$-i$

Abs:

$$|ab| = |a||b| \quad \left| \frac{a}{b} \right| = \frac{|a|}{|b|} \quad |a + b| \leq |a| + |b|$$

## Power rules:

$$\begin{aligned}x &= \sqrt[n]{a} \Leftrightarrow (x^n = a \text{ and } x \geq 0) & \sqrt[n]{-a} &= -\sqrt[n]{a}, \quad a \geq 0 \\a^{-n} &= \frac{1}{a^n} = \left(\frac{1}{a}\right)^n & a^{\frac{1}{n}} &= \sqrt[n]{a} & \sqrt{ab} &= \sqrt{a}\sqrt{b} & a^{\frac{m}{n}} &= \sqrt[n]{a^m} \\ \sqrt{\frac{a}{b}} &= \frac{\sqrt{a}}{\sqrt{b}} & a^x &= e^{x \cdot \ln a} & \sqrt[n]{a^{-m}} &= \frac{1}{\sqrt[n]{a^m}} & a^m a^n &= a^{m+n} \\ a^{m+n} & \sqrt[n]{a^m} = \sqrt[n]{a^k} & \frac{a^m}{a^n} &= a^{m-n} & \sqrt[n]{\sqrt[k]{a}} &= \sqrt[nk]{a} \\ (a^m)^n &= a^{mn} & \sqrt[n]{a} \sqrt[n]{b} &= \sqrt[n]{ab} & a^n b^n &= (ab)^n & \frac{\sqrt[n]{a}}{\sqrt[n]{b}} &= \sqrt[n]{\frac{a}{b}} \\ \frac{a^n}{b^n} &= \left(\frac{a}{b}\right)^n\end{aligned}$$

## Differentiation:

- **Sum**  $(f(x) + g(x))' = f'(x) + g'(x)$
- **Factor**  $(c \cdot f(x))' = c \cdot f'(x)$
- **Product**  $(f(x) \cdot g(x))' = f'(x)g(x) + f(x)g'(x)$
- **Quotient**  $\left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)} \quad (g \neq 0)$
- **Chain Rule**  $(f(g(x)))' = (f \circ g)' = f'(g(x))g'(x)$

## Integration:

- $\int_a^b (f(x) + / - g(x))dx = \int_a^b f(x)dx + / - \int_a^b g(x)dx$
- $\int_a^b c \cdot f(x)dx = c \cdot \int_a^b f(x)dx$
- $\int_a^b f'(x) \cdot g(x)dx = [f(x)g(x)]_a^b - \int_a^b f(x)g'(x)dx$
- $\int_a^{\phi(b)} f(x)dx = \int_a^b f(\phi(t))\phi'(t)dt$
- $a + c, b + c \in I \quad \int_a^b f(t + c)dt = \int_{a+c}^{b+c} f(x)dx$
- $ca, cb \in I: \quad \int_a^b f(ct)dt = \frac{1}{c} \int f(x)dx$
- $\int \frac{f'(t)}{f(t)}dt = \log(|f(x)|), \quad \text{bzw.} \quad \int_a^b \frac{f'(t)}{f(t)}dt = \log(f(|b|)) - \log(f(|a|))$
- **Partial fraction:**  
$$\frac{2x^6 - 4x^5 + 5x^4 - 3x^3 + x^2 + 3x}{(x-1)^3(x^2+1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{(x-1)^3} + \frac{Dx+E}{x^2+1} + \frac{Fx+G}{(x^2+1)^2}$$
- **Substitution:**  
$$\int_a^b f(x)dx = \int_{g^{-1}(a)}^{g^{-1}(b)} f(g(u))g'(u)du$$

$$\int_{\Omega} f(x, y) dx dy =$$

$$\int_{\Omega} f(g(u, v), h(u, v)) \left| \det \underbrace{\begin{pmatrix} \frac{\partial g}{\partial u} & \frac{\partial g}{\partial v} \\ \frac{\partial h}{\partial u} & \frac{\partial h}{\partial v} \end{pmatrix}}_{d\Phi = \nabla \Phi} \right| du dv$$

## Coordinate transformations:

### Polar Coordinates in $\mathbb{R}^2$

$$\begin{aligned}x &= r \cos \varphi & 0 \leq r < \infty & & dx dy &= r \cdot dr d\varphi \\ y &= r \sin \varphi & 0 \leq \varphi < 2\pi\end{aligned}$$

### Elliptic Coordinates $\mathbb{R}^2$

$$\begin{aligned}x &= r \cdot a \cos \varphi & 0 \leq r < \infty & & dx dy &= a \cdot b \cdot r \cdot dr d\varphi \\ y &= r \cdot b \sin \varphi & 0 \leq \varphi < 2\pi\end{aligned}$$

### Cylinder Coordinates $\mathbb{R}^3$

$$\begin{aligned}x &= r \cdot a \cos \varphi & 0 \leq r < \infty & & dx dy dz &= r \cdot dr d\varphi dz \\ y &= r \cdot b \sin \varphi & 0 \leq \varphi < 2\pi \\ z &= z & \infty \leq z < \infty\end{aligned}$$

### Sphere Coordinates $\mathbb{R}^3$

$$\begin{aligned}x &= r \cdot \sin \theta \cos \varphi & 0 \leq r < \infty & & dx dy dz &= r^2 \sin \theta \cdot dr d\theta d\varphi \\ y &= r \cdot \sin \theta \sin \varphi & 0 \leq \theta < \pi \\ z &= r \cos \theta & 0 \leq \varphi < 2\pi\end{aligned}$$

## Limits:

- $\lim_{x \rightarrow 0} \arctan(x) = 0, \quad \lim_{x \rightarrow \infty} \arctan(x) = \frac{\pi}{2}$
- $\lim_{x \rightarrow 0} \tan(x) = 0, \quad \lim_{x \rightarrow \infty} \tan(x) = \infty, \quad \lim_{x \rightarrow \frac{\pi}{2}} \tan(x) = \infty$
- $\lim_{x \rightarrow \infty} \cos(x) = [-1, 1], \quad \lim_{x \rightarrow \infty} \sin(x) = [-1, 1]$
- $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x \quad \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e \quad \lim_{n \rightarrow \infty} \left(1 + n\right)^{\frac{1}{n}} = e$
- $\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln(a) \quad \lim_{x \rightarrow 0} \frac{\log_a(1+x)}{x} = \frac{1}{\ln(a)} \quad \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = 0$
- $\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x^2} = \frac{1}{2} \quad \lim_{x \rightarrow 0} \frac{\tan(x)}{x} = 1 \quad \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1 = \frac{x}{\sin(x)}$
- $\lim_{x \rightarrow \infty} \frac{n!}{n^n} = 0 \quad \lim_{x \rightarrow 0} \frac{e^n - 1}{n} = 1 \quad \lim_{x \rightarrow \infty} \sqrt[n]{n!} = \infty \quad \lim_{x \rightarrow \infty} \sqrt[n]{n} = 1$

## Obvious cases:

$$\frac{1}{0} = \infty \quad \frac{1}{\infty} = 0 \quad \infty + \infty = \infty \quad 0 + \infty = \infty \quad 0^\infty = 0 \quad \infty^\infty = \infty$$

## Sums:

- $\sum_{k=0}^{\infty} aq^k = a + aq + aq^2 + \dots = \frac{a}{1-q}$  (Geometric Series)

- $\sum_{k=0}^{\infty} (k+1)q^k = 1 + 2q + 3q^2 + \dots = \frac{1}{(1-q)^2}, \quad |q| < 1$
- $\sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}$
- $\sum_{k=1}^{\infty} \frac{1}{k^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots = \frac{\pi^2}{6}$
- $\zeta(a) = \sum_{k=0}^{\infty} \frac{1}{k^a}$  ist konvergent  $\Leftrightarrow a > 1$
- $\sum_{k=0}^{\infty} \frac{1}{k!} = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots = e$
- $\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} = 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots = \frac{1}{e}$
- $\frac{1}{1 \pm x} = 1 \mp x + x^2 \mp x^3 + x^4 \mp \dots$
- $\frac{1}{(1 \pm x)^2} = 1 \mp 2x + 3x^2 \mp 4x^3 + 5x^4 \mp \dots$
- $\sqrt{1 \pm x} = 1 \pm \frac{x}{2} - \frac{1 \cdot 1}{2 \cdot 4} x^2 \pm \frac{1 \cdot 1 \cdot 3}{2 \cdot 4 \cdot 6} x^3 - \frac{1 \cdot 1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 8} x^4 \pm \dots$
- $\exp(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$
- $\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$
- $\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$
- $\sinh(x) = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$
- $\cosh(x) = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$
- $\tan(x) = 1 + \frac{\phi^3}{3} + \frac{2\phi^5}{15} + \dots$
- $\tanh(z) = 1 - \frac{z^3}{3} + \frac{2z^5}{15} - \dots$
- $\ln(1+z) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} z^k = z - \frac{z^2}{2} + \frac{z^3}{3} + \dots$
- $(1+z)^\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} z^k = 1 + \alpha z + \frac{\alpha(\alpha-1)}{2!} z^2 + \dots$

## Random:

- Circle formula:  $(x - x_0)^2 + (y - y_0)^2 = r^2$
- Ellipse formula:  $\frac{(x-x_0)^2}{a^2} + \frac{(y-y_0)^2}{b^2} = 1$
- Quadratic formula:  $x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
- Determinant:  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$
- Matrix is invertible iff. the Determinant  $\neq 0$ .
- Scalar Product:  $x \cdot y = \sum_{i=1}^n x_i y_i$
- Crossss Product:  $a \times b = (a_2 b_3 - a_3 b_2, \quad a_3 b_1 - a_1 b_3, \quad a_1 b_2 - a_2 b_1)^\top$

Let  $L$  be a **continuous linear function**, then if  $L$  is continuous at some point  $x_0$ , it is continuous at every point.


$$\lim_{x \rightarrow 0} ||L(a+x) - L(a)|| = \lim_{x \rightarrow 0} ||L(a) + L(x) - L(a)|| = \lim_{x \rightarrow 0} ||L(x)|| = \lim_{x \rightarrow 0} ||L(x_0 + x) - L(x_0)|| = 0.$$

## Finding limits:

1. Different paths
2. If  $f$  is continuous then  $\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$
3. Squeeze theorem
4. Taylor expansion
5. Polar coordinates:

$$x = r \cos(\phi), y = r \sin(\phi), \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{4x^2 + y^6} = \lim_{r \rightarrow 0} \frac{r^2}{4r^2 \cos^4(\phi) + r^6 \sin^6(\phi)}$$

## Sin and Cos



Degrees	0°	30°	45°	60°	90°	120°	135°	150°	180°
$\varphi$	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$
$\sin(\varphi)$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\cos(\varphi)$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1
$\tan(\varphi)$	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	$\pm\infty$	$-\sqrt{3}$	-1	$-\frac{\sqrt{3}}{3}$	0

$$\sin(x) = x + o(x)$$

$$\sin(x+y) \sin(x-y) = \cos^2(y) - \cos^2(x) = \sin^2(x) - \sin^2(y)$$

$$\cos(x+y) \cos(x-y) = \cos^2(y) - \sin^2(x) = \cos^2(x) - \sin^2(y)$$

$$\sin x \cos y = \frac{1}{2}(\sin(x+y) + \sin(x-y))$$

$$\cos x \cos y = \frac{1}{2}(\cos(x+y) + \cos(x-y))$$

$$\sin x \sin y = \frac{1}{2}(\cos(x-y) - \cos(x+y))$$

$$\cos(x)^2 + \sin(x)^2 = 1$$

$$\cos(\pi - x) = -\cos(x), \sin(\pi - x) = \sin(x)$$

$$\cos(x + \pi) = -\cos(x), \sin(x + \pi) = -\sin(x)$$

$$\cos(2x) = \cos^2(x) - \sin^2(x) = 1 - 2\sin^2(x) = 2\cos^2(x) - 1$$

$$\sin(2x) = 2\sin(x) \cos(x) \quad \tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$$

$$\sin\left(\frac{x}{2}\right) = \sqrt{\frac{1 - \cos(x)}{2}} \quad \cos\left(\frac{x}{2}\right) = \sqrt{\frac{1 + \cos(x)}{2}}$$

$$\tan\left(\frac{x}{2}\right) = \frac{1 - \cos(x)}{\sin(x)} = \frac{\sin(x)}{1 + \cos(x)} \quad \cot\left(\frac{x}{2}\right) = \frac{1 + \cos(x)}{\sin(x)} = \frac{\sin(x)}{1 - \cos(x)}$$

$$\sin^2(x) = \frac{1 - \cos(2x)}{2} \quad \cos^2(x) = \frac{1 + \cos(2x)}{2}$$

$$\tan(\pi + x) = \tan(x)$$

$$-\sin(-x) = \sin(x), \cos(-x) = \cos(x), \tan(-x) = -\tan(x)$$

$$\text{For all } (a, b) \in \mathbb{R}^2, \text{ such that } a^2 + b^2 = 1, \text{ there is } x \in \mathbb{R}, \text{ such that } a = \cos(x), b = \sin(x).$$

$$\sin(x) = \frac{2\tan(x/2)}{1 + \tan^2(x/2)} \quad \cos(x) = \frac{1 - \tan^2(x/2)}{1 + \tan^2(x/2)}$$

$$\sin(x) = \frac{e^{ix} - e^{-ix}}{2i} \quad \cos(x) = \frac{e^{ix} + e^{-ix}}{2}$$

$$\int_0^{2\pi} \sin(t) \cdot \cos(t) dt = \int_0^{2\pi} \sin(t) dt = \int_0^{2\pi} \cos(t) dt = 0$$

$$\int \sin^2(x) dx = \frac{1}{2}(x - \sin(x) \cos(x))$$

$$\int \cos^2(x) dx = \frac{1}{2}(x + \sin(x) \cos(x))$$

$$\int x \sin(x) dx = \sin(x) - x \cos(x)$$

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

$$\sin(\arccos(x)) = \sqrt{1 - x^2} \quad \sin(\arctan(x)) = \frac{x}{\sqrt{1 + x^2}}$$

$$\cos(\arctan(x)) = \frac{1}{\sqrt{1 + x^2}} \quad \cos(\arcsin(x)) = \sqrt{1 - x^2}$$

$$\tan(\arcsin(x)) = \frac{x}{\sqrt{1 - x^2}} \quad \tan(\arccos(x)) = \frac{\sqrt{1 - x^2}}{x}$$

$$\cosh(x) := \frac{e^x + e^{-x}}{2} \quad \sinh x := \frac{e^x - e^{-x}}{2} \quad \tanh x := \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

$$\sinh(z) = -\sinh(-z) \quad \cosh(z) = \cosh(-z)$$

$$\sinh(z) = \sinh(z + 2\pi i) \quad \cosh(z) = \cosh(z + 2\pi i)$$

$$\sinh(z_1 \pm z_2) = \sinh(z_1) \cdot \cosh(z_2) \pm \sinh(z_2) \cdot \cosh(z_1)$$

$$\sinh(z_1 \pm z_2) = \cosh(z_1) \cdot \cosh(z_2) \pm \sinh(z_2) \cdot \sinh(z_1)$$

$$\tanh(z_1 \pm z_2) = \frac{\tanh(z_1) \pm \tanh(z_2)}{1 \pm \tanh(z_1) \cdot \tanh(z_2)}$$

## Prove if $f$ is differentiable at $x_0$

$f$  continuous at  $x_0$ ? No  $\Rightarrow$  Not differentiable.

$\Downarrow$  Yes

Is  $f$  partially differentiable at  $x_0$ , does  $\partial_{x_i} f(x_0)$  exist for all  $i$ ? No  $\Rightarrow$  Not differentiable.

$\Downarrow$  Yes

Are all partial derivatives continuous at  $x_0$ ? Yes  $\Rightarrow f$  is differentiable at  $x_0$  with  $df(x_0) = J_f(x_0)$ .

$\Downarrow$  No

Does a linear mapping  $u$  exist, such that  $\lim_{\substack{x \rightarrow x_0 \\ x \neq 0}} \frac{f(x) - f(x_0) - u(x - x_0)}{\|x - x_0\|} = 0$ ?

Yes  $\Rightarrow f$  is differentiable at  $x_0$ .

No  $\Rightarrow f$  is not differentiable at  $x_0$  \*cries\*.

## Primitives and Derivatives

$$\int (ax + b)^s dx = \frac{1}{a(s+1)}(ax + b)^{s+1} + C, s \neq -1$$

$$\int \frac{1}{ax + b} dx = \frac{1}{a} \log|ax + b| + C$$

$$\int (ax^p + b)^s x^{p-1} dx = \frac{(ax^p + b)^{s+1}}{ap(s+1)} + C, s \neq -1, a \neq 0$$

$$\int (ax^p + b)^{-1} x^{p-1} dx = \frac{1}{ap} \log|ax^p + b| + C, a \neq 0, p \neq 0$$

$$\int \frac{ax + b}{cx + d} dx = \frac{ax}{c} - \frac{ad - bc}{c^2} \log|cx + d| + C$$

$$\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$$

$$\int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log\left|\frac{x - a}{x + a}\right|$$

$$\int \sqrt{a^2 + x^2} dx = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \log(x + \sqrt{a^2 + x^2}) + C$$

$$\int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin\left(\frac{x}{|a|}\right) + C$$

$$\int \sqrt{x^2 - a^2} dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log|x + \sqrt{x^2 - a^2}| + C$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \log(x + \sqrt{a^2 + x^2}) + C$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \log|x + \sqrt{x^2 - a^2}| + C$$



$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \arcsin\left(\frac{x}{|a|}\right) + C$$

$$\int e^{kx} dx = \frac{1}{k} e^{kx} + C$$

$$\int a^{kx} dx = \frac{1}{k * \log(a)} a^{kx} + C$$

$$\int e^{ax} p(x) dx = e^{ax} (a^{-1} p(x) - a^{-2} p'(x) + a^{-3} p''(x) - \dots + (-1)^n a^{-n-1} p^{(n)}(x)) + C, a \neq 0, p: \text{Polynome of degree } n$$

$$\int e^{kx} \sin(ax+b) dx = \frac{e^{kx}}{a^2 + k^2} \left( k \sin(ax+b) - a \cos(ax+b) \right) + C$$

$$\int e^{kx} \cos(ax+b) dx = \frac{e^{kx}}{a^2 + k^2} \left( k \cos(ax+b) + a \sin(ax+b) \right) + C$$

$$\int \log|x| dx = x(\log|x| - 1) + C$$

$$\int x^k \log(x) dx = \frac{x^{k+1}}{k+1} \left( \log(x) - \frac{1}{k+1} \right) + C, k \neq -1$$

$$\int x^{-1} \log(x) dx = \frac{1}{2} (\log(x))^2 + C$$

$$\int \sin(ax+b) dx = -\frac{1}{a} \cos(ax+b) + C$$

$$\int \cos(ax+b) dx = \frac{1}{a} \sin(ax+b) + C$$

$$\int \sin^2(x) dx = \frac{1}{2} (x - \sin(x) \cos(x)) + C$$

$$\int \cos^2(x) dx = \frac{1}{2} (x + \sin(x) \cos(x)) + C$$

$$\int \tan^2(x) dx = \tan(x) - x + C$$

$$\int \frac{1}{\sin(x)} dx = \log \left| \tan \frac{x}{2} \right| + C$$

$$\int \frac{1}{\cos(x)} dx = \log \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

$$\int \frac{1}{\tan(x)} dx = \log |\sin(x)| + C$$

$$\int_0^{2\pi} \sin(mx) \cos(nx) dx = 0, m, n \in \mathbb{Z}$$

$$\int_0^\infty \frac{\sin(ax)}{x} dx = \frac{\pi}{2}, a > 0$$

$$\int_0^\infty \sin(x^2) dx = \int_0^\infty \cos(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

$$\int_0^\infty e^{-ax} x^n dx = \frac{n!}{a^{n+1}}, a > 0$$

$$\int_0^\infty e^{-ax^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}}, a > 0$$

**Differentiable**  $\implies$  **Continuous**  $\implies$  **Integrable**

$f'(x)$	$f(x)$	$F(x)$
0	$c \ (c \in \mathbb{R})$	$cx$
$c$	$cx$	$\frac{c}{2} x^2$
$r \cdot x^{r-1}$	$x^r \ (r \in \mathbb{R} \setminus \{-1\})$	$\frac{x^{r+1}}{r+1}$
$\frac{-1}{x^2} = -x^{-2}$	$\frac{1}{x} = x^{-1}$	$\log x $
$\frac{1}{2\sqrt{x}} = -x^{-2}$	$\sqrt{x} = x^{\frac{1}{2}}$	$\frac{2}{3} x^{\frac{3}{2}}$
$\cos x$	$\sin x$	$-\cos x$
$-\sin x$	$\cos x$	$\sin x$
$1 + \tan^2 x = \frac{1}{\cos^2 x}$	$\tan x$	$-\log \cos x $
$-\frac{1}{\sin^2(x)}$	$\cot x$	$\log \sin x $
$e^x$	$e^x$	$e^x$
$c \cdot e^{cx}$	$e^{cx}$	$\frac{1}{c} e^{cx}$
$\log a \cdot a^x$	$a^x$	$\frac{a^x}{\log a}$
$\frac{1}{x}$	$\log x $	$x(\log x  - 1)$
$\frac{1}{\log a \cdot x}$	$\log_a x $	$\frac{x}{\log a} (\log x  - 1)$ $= x(\log_a x  - \log_a e)$
$\frac{1}{\sqrt{1-x^2}}$	$\arcsin x$	$x \arcsin x + \sqrt{1-x^2}$
$-\frac{1}{\sqrt{1-x^2}}$	$\arccos x$	$x \arccos x - \sqrt{1-x^2}$
$\frac{1}{1+x^2}$	$\arctan x$	$x \arctan x - \frac{1}{2} \log(1+x^2)$
$\sinh(x)$	$\cosh(x)$	-
$\cosh(x)$	$\sinh(x)$	-
$\frac{1}{\cosh^2(x)}$	$\tanh(x)$	$\log(\cosh(x))$
$2 \sin(x) \cos(x)$	$\sin^2(x)$	$\frac{1}{2} (x - \sin(x) \cos(x))$
$-2 \sin(x) \cos(x)$	$\cos^2(x)$	$\frac{1}{2} (x + \sin(x) \cos(x))$
$\frac{2 \sin(x)}{\cos^3(x)}$	$\tan^2(x)$	$\tan(x) - x$
$\frac{1}{\sqrt{x^2+1}}$	$\operatorname{arsinh} x$	$x \operatorname{arsinh} x - \sqrt{x^2+1}$
$\frac{1}{\sqrt{x^2-1}} \ (x > 1)$	$\operatorname{arcosh} x$	$x \operatorname{arcosh} x - \sqrt{x^2-1}$
$\frac{1}{1-x^2} \ ( x  < 1)$	$\operatorname{artanh} x$	$x \operatorname{artanh} x + \frac{1}{2} \ln(1-x^2)$
$\frac{1}{1-x^2} \ ( x  > 1)$	$\operatorname{arcoth} x$	$x \operatorname{arcoth} x + \frac{1}{2} \ln(x^2-1)$