

# Searches for Heavy Hadronic Resonances with the ATLAS and CMS detectors at the LHC

---

**Caterina Doglioni\***

*University of Geneva*

*E-mail:* [caterina.doglioni@cern.ch](mailto:caterina.doglioni@cern.ch)

**Francesco Santanastasio<sup>†</sup>**

*CERN*

*E-mail:* [francesco.santanastasio@cern.ch](mailto:francesco.santanastasio@cern.ch)

.....

*VI Italian workshop on p-p physics at the LHC,*

*8-10 May 2013*

*Acquario di Genova, Ponte Spinola, Area Porto Antico, Genova, Italy*

---

\*Speaker.

<sup>†</sup>Speaker.

## 1. Introduction

New resonances, that couples to quarks and gluons and decay to final states with hadronic jets, can be produced copiously at the Large Hadron Collider (CERN). This review outlines a selection of searches for heavy resonances in hadronic final states performed at the ATLAS and CMS experiments using the proton-proton collision data collected at a center-of-mass energy of 7 TeV (2011) and 8 TeV (2012). A series of points that are deemed important for current and future searches are also discussed.

## 2. Overview of selected searches in the ATLAS and CMS experiments

The searches for hadronic resonances can be divided into two main categories:

- **Resolved topology** - Searches where the quarks and gluons produced in the final state are each reconstructed at detector level into single, *resolved* hadronic jets. Examples are the single production of a resonance  $X$  decaying to a pair of gluons, or the pair production of two resonances  $X$  at rest each decaying to a quark-antiquark pair;
- **Boosted topology** - Searches where the quarks and gluons produced in the final state are merged into a single reconstructed jet. A typical example is the decay of a resonance  $X$  into a pair of massive particles  $Y$ , with  $M_X \gg M_Y$ , and  $Y$  decays to a pair of light quarks. In this case the resonance  $Y$  will be *boosted* (the resonance has a large momentum compared to its mass) so that a single hadronic jet with a large distance parameter (*fat jet*) encompasses all its decay products. Techniques that exploit the presence of substructure within the *fat jet* are employed to reject background from standard QCD jets and multiple interactions within the same bunch crossing (pile-up).

The quintessential example of hadronic search with resolved jets is the search for heavy resonances in the dijet mass distribution. New particles, or excitations of quarks indicating compositeness, could manifest themselves as narrow 'bumps' in the dijet mass distribution of central leading and subleading jets above the continuum QCD background [5, 6]. The search can be tailored to specific resonances decaying to heavy quark flavors using  $b$ -tagging for one or both jets [7]. No significant excess over the background has been found for the current ATLAS and CMS analyses, and lower limits are set on the masses of new particles including model-independent Gaussian resonances of varying width. Upper limits on cross section times branching fraction to jets times acceptance of the order of 100, 10, and 1 fb at resonance masses of 1.5, 3, and 4.5 TeV, respectively, are set by both experiments.

Searches for resonances in final states with high jet multiplicity, tailored towards R-Parity Violating supersymmetric signatures (such as 3-body decays into quarks of pair produced gluinos, giving a six jet final state), can be performed in both resolved and boosted regimes [8, 9]. Both experiments select events with six or more jets, but the background estimation techniques for ATLAS and CMS differ. In the resolved channel, ATLAS employs the  $p_T$  of the sixth jet as a discriminant variable, while CMS performs a 'bump-search' in the three-jet invariant mass. The combinatorial background (from both the QCD background and the signal) penalizes the CMS search and allows

the ATLAS search to set more stringent limits. A proof-of-principle boosted jet analysis, although not as sensitive as the resolved one, is also carried on by the ATLAS experiment.

In the case of resonances decaying into pairs of top quarks ( $t\bar{t}$ ) or heavy bosons ( $WW$ ,  $ZZ$ ,  $HH$ ,  $HZ$ , ..), the use of jet substructure techniques is crucial to achieve a good background rejection. In these cases, the tops or heavy bosons coming from a TeV-scale resonance are boosted, and therefore their decay products are spatially collimated when reconstructed in the detector.

Specific techniques for top-tagging, based on the presence of three hard energy deposit corresponding to the top decay products, have been employed to distinguish top-jets from jets originated from quarks and gluons that constitute the majority of the QCD background [10, 11, 12, 13]. Both ATLAS and CMS look for heavy resonances decaying in  $t\bar{t}$  at 7 and 8 TeV, in semileptonic and all-hadronic top decays [14, 15, 16, 17, 18]. The ATLAS analysis employs b-tagging to further suppress the QCD background and it sets limits starting from a resonance mass of 500 GeV. The CMS analysis, instead, does not use b-tagging and limits are set starting at a resonance mass of 1 TeV. The CMS expected upper limits on the resonance cross section are about a factor 3-4 lower than the ATLAS ones for resonance masses around 2 TeV. A possible reason might be due to the use of b-tagging: on one side, the b-tag requirement allows the ATLAS analysis to start the search at lower resonance masses compared to CMS, by reducing significantly the QCD background; on the other hand, the low b-tag efficiency at high jet  $p_T$  might penalize the ATLAS search at high resonance masses.

The CMS experiment has also performed a search for RS gravitons decaying to  $WW/ZZ$  and  $W/Z'$  decaying to  $Wq/Zq$  [19]. This analysis employs  $W/Z$ -tagging techniques based on the jet mass and the presence of hard sub-jets to significantly reduce the background and keep a relatively high signal efficiency. Given no excesses above background, limits are set on a number benchmark models.

### 3. Discussion points

#### 3.1 Jet energy scale in ATLAS and CMS

- Some details on high- $p_T$  JES [20][21]
- 2 plots side by side: background rejection after boson tagging and one of the limit plots
- Encourage more work on both sides to make consistent assumptions

#### 3.2 Setting limits on hadronic resonances

- Refer to earlier plots
- Differences: full template/narrow width, inclusive vs dijet-only search
- Recommend to unify definition for 14 TeV (combination)? [22]

#### 3.3 Data for low-mass resonances

- One/two-sentence description of need to look lower in  $m_{jj}$ , hadronic rate problems, delayed triggers and parked data
- one plot only: CMS 2011 low-mass extension

### 3.4 New directions in searches with jet substructure

- Justification on why substructure necessary at 14 TeV (and not for 8 TeV Higgs yet)
- Need for MC generator support and benchmarks among experiments [23][24]
- New ideas:
  - Q/g tagging for lower masses
  - Tools for  $W'$  and  $Z'$ : jet charge and angle between subjects (angular correlations?)
- Open questions:
  - Particle flow and substructure
  - High-lumi and substructure

## 4. Conclusions

### References

- [1] M. Cacciari, G. P. Salam, and G. Soyez, *The anti- $k_t$  jet clustering algorithm*, *JHEP* **04** (2008) 063, [[arXiv:0802.1189](#)].
- [2] ATLAS Collaboration, *Jet energy scale and its systematic uncertainty in proton-proton collisions at  $\sqrt{s}=7$  TeV with atlas 2011 data*, ATLAS-CONF-2013-004, CERN, Jan, 2013.
- [3] CMS Collaboration, *Determination of jet energy calibration and transverse momentum resolution in CMS*, *Journal of Instrumentation* **6** (2011), no. 11 P11002.
- [4] ATLAS Collaboration, *Jet energy measurement with the ATLAS detector in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, *Eur.Phys.J.* **C73** (2013) 2304, [[arXiv:1112.6426](#)].
- [5] CMS Collaboration, *Search for narrow resonances using the dijet mass spectrum with 19.6fb-1 of pp collisions at  $\sqrt{s}=8$  TeV*, CMS-PAS-EXO-12-059, CERN, 2013.
- [6] ATLAS Collaboration, *Search for new phenomena in the dijet mass distribution updated using 13.0 fb<sup>-1</sup> of pp collisions at  $\sqrt{s} = 8$  TeV collected by the ATLAS detector*, ATLAS-CONF-2012-148, CERN, Nov, 2012.
- [7] CMS Collaboration, *Search for heavy resonances decaying into  $bb$  and  $bg$  final states in pp collisions at  $\sqrt{s} = 8$  TeV*, CMS-PAS-EXO-12-023, CERN, 2013.
- [8] CMS Collaboration, *Search for three-jet resonances in pp collisions at*, *Physics Letters B* **718** (2012), no. 2 329 – 347.
- [9] ATLAS Collaboration, *Search for pair production of massive particles decaying into three quarks with the ATLAS detector in  $\sqrt{s} = 7$  TeV pp collisions at the LHC*, *Journal of High Energy Physics* **2012** (2012), no. 12 1–42.
- [10] D. E. Kaplan, K. Rehermann, M. D. Schwartz, and B. Tweedie, *Top Tagging: A Method for Identifying Boosted Hadronically Decaying Top Quarks*, *Phys.Rev.Lett.* **101** (2008) 142001, [[arXiv:0806.0848](#)].
- [11] T. Plehn, M. Spannowsky, M. Takeuchi, and D. Zerwas, *Stop Reconstruction with Tagged Tops*, *JHEP* **1010** (2010) 078, [[arXiv:1006.2833](#)].

- [12] ATLAS Collaboration, *Performance of large- $r$  jets and jet substructure reconstruction with the atlas detector*, ATLAS-CONF-2012-065, CERN, Jul, 2012.
- [13] L. G. Almeida, S. J. Lee, G. Perez, G. Sterman, and I. Sung, *Template Overlap Method for Massive Jets*, *Phys.Rev.* **D82** (2010) 054034, [[arXiv:1006.2035](#)].
- [14] CMS Collaboration, *Search for anomalous  $t\bar{t}$  production in the highly-boosted all-hadronic final state*, *Journal of High Energy Physics* **2012** (2012), no. 9 1–41.
- [15] ATLAS Collaboration, *Search for resonances decaying into top-quark pairs using fully hadronic decays in  $pp$  collisions with ATLAS at  $\sqrt{s}=7$  TeV*, *Journal of High Energy Physics* **2013** (2013), no. 1 1–50.
- [16] CMS Collaboration, *Search for anomalous top quark pair production in the boosted all-hadronic final state using  $pp$  collisions at  $\sqrt{s}=8$  TeV*, CMS-PAS-B2G-12-005, CERN, 2013.
- [17] *A search for  $t\bar{t}$  resonances in the lepton plus jets final state with atlas using  $14\text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s}=8$  tev*, ATLAS-CONF-2013-052, CERN, Geneva, May, 2013.
- [18] CMS Collaboration, *Search for  $t\bar{t}$  resonances in semileptonic final state*, CMS-PAS-B2G-12-006, CERN, 2013.
- [19] CMS Collaboration, *Search for heavy resonances in the W/Z-tagged dijet mass spectrum in  $pp$  collisions at 7 TeV*, [arXiv:1212.1910](#).
- [20] CMS Collaboration, *Single-particle response in the CMS calorimeters*, CMS-PAS-JME-10-008, CERN, 2010.
- [21] ATLAS Collaboration, *Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC*, *Eur.Phys.J.* **C73** (2013) 2305, [[arXiv:1203.1302](#)].
- [22] M. Cacciari, J. Rojo, G. P. Salam, and G. Soyez, *Quantifying the performance of jet definitions for kinematic reconstruction at the LHC*, *JHEP* **0812** (2008) 032, [[arXiv:0810.1304](#)].
- [23] CMS Collaboration, *Studies of jet mass in dijet and W/Z+jet events*, [arXiv:1303.4811](#).
- [24] ATLAS Collaboration, *Jet mass and substructure of inclusive jets in  $\sqrt{s}=7$  TeV  $pp$  collisions with the ATLAS experiment*, *JHEP* **1205** (2012) 128, [[arXiv:1203.4606](#)].