

**TEMPLATE FOR PIC 2012 PROCEEDINGS
PREPARED BY LATEX**

PIC 2012 AUTHOR

*Institute of Experimental Physics SAS
Watsonova 47, 040 01 Košice, Slovak Republic
E-mail: pic@saske.sk*

This is abstract for your PIC 2012 contribution. It should be a few lines briefly describing your contribution to the Proceedings. Please, remember that deadline for your contribution is 1st December 2012.

1 Guidelines

Please, take from the PIC 2012 web-site a Latex class file: 'pic2012.cls' for preparation of your PIC 2012 contribution. This file is also part of archive, which was sent to every author, together with this template.

2 Where to send your contribution

The proceedings will be first reviewed by the session convenors. Please, send a .pdf version to your session convenor before the deadline (for poster presenters with no talk, please send your proceeding to urban@upjs.sk). After having received the comments from the session convenor, the authors are required to submit their papers and source files to the ePrint arXiv server.

All papers for the proceedings of this conference must be submitted to the eprint arXiv (<http://arXiv.org>). The web upload is particularly easy to use, from a PC, Mac or on Unix. You will receive confirmation email upon every step of the process, if you don't, it means you have still more to do. The ultimate goal is to see the full text of your paper on the arXiv.org server with an eprint number in the form "arch-ive/yymmnnn".

Please, include into Comment filed during submission into arXiv (Metadata definition step during submission procedure) the following:

Physics in Collision, Slovakia, 2012

and in the Report field:

PSNUM XX

where XX is the number of your talk on the Conference Web Scientific program page:

<http://www.saske.sk/PIC2012/index.php?id=11>

The online conference proceeding will appear on SLAC eConf server. In addition the regular paper book will be printed and sent to all participants.

Please, send your contribution arXiv number to our conference e-mail:

pic@saske.sk

subject: author name - session

3 File Size

The eprint arXiv (which will be used to submit contribution) will reject files over 650k. This problem generally occurs if the paper contains a lot of large figures. Usually large figures are the result of converting jpegs or scanned images to PS files. To avoid this problem, submit smaller graphics, with lower resolution. There is help on how to do this at the eprint arXiv Web site. In principle, you can provide a link to your Web page in the arXiv Comments section pointing users to better quality pictures.

4 Limits of PIC 2012 contributions

The limit for the size of a PIC 2012 contribution:

4 pages for regular poster,

6 pages for orally presented poster,

10 pages for talks 20-30 min's (as stated on conference Indico page) 14 pages for talks 40-45 min.

5 Deadline for your contribution the the PIC 2012 Proceedings

1st December 2012

6 Figures

Please prepare the figures in high resolution (600-1200 dpi). Half-tone pictures must be sharp enough for reproduction, otherwise they will be rejected.

7 Text sample

QCD is an essential ingredient of the Standard Model, and it is well tested in hard processes when transferred momentum is of the order of the total collision energy (Bjorken limit: $Q^2 \sim s \rightarrow \infty$). The cornerstones of perturbative QCD at this kinematic regime (QCD-improved parton model): the Gribov–Lipatov–Altarelli–Parisi–Dokshitzer (GLAPD) evolution equation and factorization of inclusive hard processes provides a basis for the successful QCD-improved parton model. The factorization theorem for inclusive hard processes ensures that the inclusive cross section factorizes into partonic subprocess(es) and parton distribution function(s). The GLAPD evolution equation governs the $\log Q^2$ -dependence (at $Q^2 \rightarrow \infty$) of the parton distribution functions and the hard subprocess cross-sections at fixed scaling variable $x = Q^2/s$.

Another kinematic domain that is very important at high-energy is given by the (Balitsky–Fadin–Kuraev–Lipatov) BFKL limit [1,2], or QCD Regge limit, whereby at fixed $Q^2 \gg \Lambda_{QCD}^2$, $s \rightarrow \infty$. In the BFKL limit, the BFKL evolution in the leading order (LO) governs $\log(1/x)$ evolution (at $x \rightarrow 0$) of inclusive processes.

Note that the BFKL evolution in the next-to-leading order (NLO) [3,4,5], unlike the LO BFKL [1,2], partly includes GLAPD evolution with the running coupling constant of the LO GLAPD, $\alpha_S(Q^2) = 4\pi/\beta_0 \log(Q^2/\Lambda_{QCD}^2)$.

Photon–photon collisions, particularly $\gamma^*\gamma^*$ processes, play a special role in QCD [6], since their analysis is under much better control than the calculation of lepton–hadron and hadron–hadron processes, which require the input of non-perturbative hadronic structure functions or wave functions. In addition, unitarization (screening) corrections due to multiple Pomeron exchange should be less important for the scattering of γ^* of high virtuality than for hadronic collisions.

Figure 1. The energy dependence of the total cross section for highly virtual photon–photon collisions predicted by the BLM scale-fixed NLO BFKL [10] compared with recently finalized OPAL [11] and L3 [12] data from LEP2 at CERN. The (solid) dashed curves correspond to the (N)LO BFKL predictions for two different choices of the Regge scale: $s_0 = Q^2$ for upper curves and $s_0 = 4Q^2$ for lower curves

The high-energy asymptotic behaviour of the $\gamma\gamma$ total cross section in QED can be calculated [7] by an all-orders resummation of the leading terms: $\sigma \sim \alpha^4 s^\omega$, $\omega = \frac{11}{32}\pi\alpha^2 \simeq 6 \times 10^{-5}$. However, the slowly rising asymptotic behaviour of the QED cross section is not apparent since large contributions come from other sources, such as the cut of the fermion-box contribution: $\sigma \sim \alpha^2(\log s)/s$ [6] (which although subleading in energy dependence, dominates the rising contributions by powers of the QED coupling constant) and QCD-driven processes.

The photon–photon cross sections with LO BFKL resummation was considered in Refs. [2,8,9].

Although the complete NLO impact factor of the virtual photon is not known yet, one can use the LO impact factor of Refs. [7,9], assuming that the main energy-dependent NLO corrections come from the NLO BFKL subprocess rather than from

the photon impact factors [10].

Fig. 1 compares the LO and BLM scale-fixed NLO BFKL predictions $\sigma \sim \alpha^2 \alpha_S^2 s^\omega$ [5,10] with recent CERN LEP2 data from OPAL [11] and L3 [12].

Acknowledgements

We thank all DIS'03 participants for their contributions to the DIS'03 Proceedings. This work was supported in part by the High Energy Foundation and the World Science Agency.

Appendix

This is place for Appendix, if any.

References

1. V.S. Fadin, E.A. Kuraev, and L.N. Lipatov, Phys. Lett. B **60** (1975) 50; E.A. Kuraev, L.N. Lipatov, and V.S. Fadin, Zh. Eksp. Teor. Fiz. **71** (1976) 840 [Sov. Phys. JETP **44** 443 (1976)]; *ibid.* **72** (1977) 377 [**45** (1977) 199].
2. I.I. Balitsky and L.N. Lipatov, Yad. Fiz. **28** (1978) 1597 [Sov. J. Nucl. Phys. **28** (1978) 822].
3. V.S. Fadin and L.N. Lipatov, Phys. Lett. **B429** (1998) 127.
4. G. Camici and M. Ciafaloni, Phys. Lett. **B430** (1998) 349.
5. S.J. Brodsky, V.S. Fadin, V.T. Kim, L.N. Lipatov, and G.B. Pivovarov, Pis'ma ZhETF **70** (1999) 161; [JETP Lett. **70** (1999) 155].
6. V.M. Budnev, I.F. Ginzburg, G.V. Meledin, and V.G. Serbo, Phys. Rep. **C15** (1975) 181.
7. V.N. Gribov, L.N. Lipatov, and G.V. Frolov, Phys. Lett. **B31** (1970) 34; Yad. Fiz. **12** (1970) 994 [Sov. J. Nucl. Phys. **12** (1971) 543]; H. Cheng and T.T. Wu, Phys. Rev. D **1** (1970) 2775.
8. J. Bartels, A. De Roeck, and H. Lotter, Phys. Lett. **B389** (1996) 742; M. Boonekamp, A. De Roeck, C. Royon, and S. Wallon, Nucl. Phys. **B555** (1999) 540.
9. S.J. Brodsky, F. Hautmann, and D.E. Soper, Phys. Rev. **D56** (1997) 6957; Phys. Rev. Lett. **78** (1997) 803; (E) **79** (1997) 3544.
10. S.J. Brodsky, V.S. Fadin, V.T. Kim, L.N. Lipatov and G.B. Pivovarov, Pisma ZhETF **76** (2002) 306 [JETP Lett. **76** (2002) 249].
11. OPAL, G. Abbiendi *et al.*, Eur. Phys. J. **C24** (2002) 17.
12. L3, P. Achard *et al.*, Phys. Lett. **B531** (2002) 39.
13. ZEUS, J. Breitweg *et al.*, Eur. Phys. J. **C6** (1999) 239; H1, C. Adloff *et al.*, Nucl. Phys. **B538** (1999) 3.
14. DØ, B. Abbott *et al.*, Phys. Rev. Lett. **84** (2000) 5722.