Vector problem

$1 \quad 12^{th} \text{ Maths}$ - Chapter 12

This is Problem-6 from Exercise 10.5

- 1. Find a vector of magnitude 5 units, and parallel to the resultant of the vectors $\mathbf{a} = 2\hat{\mathbf{i}} + 3\hat{\mathbf{j}} \hat{\mathbf{k}}$ and $\mathbf{b} = \hat{\mathbf{i}} 2\hat{\mathbf{j}} + \hat{\mathbf{k}}$ Solution:
- 2. sum of the resultant vectors **a** and **b** is

$$\mathbf{a} = \begin{pmatrix} 2\\3\\-1 \end{pmatrix} \tag{1}$$

$$\mathbf{b} = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \tag{2}$$

$$\mathbf{c} = \mathbf{a} + \mathbf{b} \tag{3}$$

$$\mathbf{c} = \begin{pmatrix} 2\\3\\-1 \end{pmatrix} + \begin{pmatrix} 1\\-2\\1 \end{pmatrix} \tag{4}$$

$$\mathbf{c} = \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} \tag{5}$$

(6)

3. The unit direction of the vector \mathbf{c} is

$$\mathbf{\hat{c}} = \frac{\mathbf{c}}{\|\mathbf{c}\|}$$

$$\hat{\mathbf{c}} = \frac{\begin{pmatrix} 3\\1\\0 \end{pmatrix}}{\sqrt{3^2 + 1^2}}$$

$$\hat{\mathbf{c}} = \frac{\begin{pmatrix} 3\\1\\0 \end{pmatrix}}{\sqrt{10}}$$

4. Hence, the vector of magnitude 5 units and parallel to the resultant of vectors ${\bf a}$ and ${\bf b}$ is $\pm 5 {\bf \hat{c}}$

$$\Rightarrow \pm \frac{5}{\sqrt{10}} \begin{pmatrix} 3\\1\\0 \end{pmatrix}$$

$$\Rightarrow \pm \begin{pmatrix} \frac{3\sqrt{10}}{2}\\\frac{\sqrt{10}}{2}\\0 \end{pmatrix}$$
(8)

$$\implies \pm \begin{pmatrix} \frac{3\sqrt{10}}{2} \\ \frac{\sqrt{10}}{2} \\ 0 \end{pmatrix} \tag{8}$$