

UNIMODULAR COVERS OF 3-DIMENSIONAL PARALLELEPIPEDS AND CAYLEY SUMS

ABSTRACT. We show that the following classes of lattice polytopes have unimodular covers, in dimension three: parallelepipeds, smooth centrally symmetric polytopes, and Cayley sums $\text{Cay}(P, Q)$ where the normal fan of Q refines that of P . This improves results of Beck et al. (2018) and Haase et al. (2008) where the last two classes were shown to be IDP.

1. INTRODUCTION

A lattice polytope $P \subset \mathbb{R}^d$ has the *integer decomposition property* if for every positive integer n , every lattice point $p \in nP \cap \mathbb{Z}^d$ can be written as a sum of n lattice points in P . We abbreviate this by saying that “ P is IDP”. Being IDP is interesting in the context of both enumerative combinatorics (Ehrhart theory) and algebraic geometry (projective normality of toric varieties). It falls into a hierarchy of several properties each stronger than the previous one; see, e.g., [3, Section 2.D], [10, Sect. 1.2.5], [13, p. 2097], [14, p. 2313]. Let us here only mention that

P has a unimodular triangulation $\Rightarrow P$ has a unimodular cover $\Rightarrow P$ is IDP.

Remember that a *unimodular triangulation* is a triangulation of P into unimodular simplices, and a *unimodular cover* is a collection of unimodular simplices whose union equals P .

Oda [15] posed several questions regarding smoothness and the IDP property for lattice polytopes. Following [9, 19], we say that a pair (P, Q) of lattice polytopes has the integer decomposition property, or that *the pair (P, Q) is IDP*, if

$$(P + Q) \cap \mathbb{Z}^d = P \cap \mathbb{Z}^d + Q \cap \mathbb{Z}^d,$$

where $A + B := \{a + b : a \in A, b \in B\}$ denotes the *Minkowski sum* of two sets $A, B \subset \mathbb{R}^d$.

A lattice polytope Q is called *smooth* if it is simple and the primitive edge directions at every vertex form a linear basis for the lattice; equivalently, if the projective toric variety defined by the normal fan of Q is smooth. The following versions of Oda’s questions are now considered conjectures [11, 14], and they are open even in dimension three:

- Conjecture 1.1.** (1) (Related to problems 2 and 5 in [15]) Every smooth lattice polytope is IDP.
 (2) (Related to problems 1, 3, 4, 6 in [15]) Every pair (P, Q) of lattice polytopes with Q smooth and the normal fan of Q refining that of P is IDP.

When the normal fan of a polytope Q refines that of another polytope P , as in the second conjecture, we say that P is a *weak Minkowski summand* of Q , since this is easily seen to be equivalent to the existence of a polytope P' such that $P + P' = kQ$ for some dilation constant $k > 0$. This property has the following algebraic implication for the projective toric variety X_Q : P is a weak Minkowski summand of Q if and only if the Cartier divisor defined by P on X_Q is *numerically effective*, or “nef” (see [5, Cor. 6.2.15, Thm. 6.3.12], but observe that what we here call “weak Minkowski summand” is called “N-Minkowski summand” there).

added def of Minkowski sum — P.

Motivated by these and other questions, several authors have studied the IDP property for different classes of lattice polytopes, with special attention to dimension 3 (in dimension 2 it is straightforward that every lattice polygon has unimodular triangulations). For example, very recently Beck et al. [1] proved that all smooth centrally symmetric 3-polytopes are IDP. More precisely, they show that any such polytope can be covered by lattice parallelepipeds (affine images of 3-cubes) and unimodular simplices, both of which are trivially IDP. In Section 2 we show:

Theorem 1.2. *Every 3-dimensional lattice parallelepiped has a unimodular cover.*

This, together with the mentioned result from [1], gives:

Corollary 1.3. *Every smooth centrally symmetric lattice 3-polytope has a unimodular cover.* \square

These results leave open the following important questions regarding parallelo-
topes:

Question 1.4. Do 3-dimensional parallelepipeds have unimodular triangulations?

Question 1.5. Higher dimensional parallelotopes (affine images of cubes) are IDP. Do they have unimodular covers?

The two-dimensional case of Conjecture 1.1(2) is known to hold, with three different proofs by Fakhruddin [7], Ogata [16] and Haase et al. [11]. This last one actually shows that smoothness of Q is not needed. In dimension three, however, the conjecture fails without the smoothness assumption. Indeed, if we let $P = Q$ be any non-unimodular *empty tetrahedron*, then P is obviously a weak Minkowski summand of Q but the pair (P, Q) is not IDP. By an empty tetrahedron we mean a lattice tetrahedron containing no lattice points other than its vertices (see the proof of Lemma 2.2 for a classification of them).

An alternative approach to Conjecture 1.1(2) is via Cayley sums, which we discuss in Section 3. Recall that the *Cayley sum* of two lattice polytopes $P, Q \subset \mathbb{R}^d$ is the lattice polytope

$$\text{Cay}(P, Q) := \text{conv}(P \times \{0\} \cup Q \times \{1\}) \subset \mathbb{R}^{d+1}.$$

We normally require $\text{Cay}(P, Q)$ to be full-dimensional (otherwise we can delete coordinates) but P or Q do not necessarily need to be full-dimensional. We only require the linear subspaces parallel to them to span \mathbb{R}^d .

As we note in Proposition 3.1, if the Cayley sum of P and Q is IDP then the pair (P, Q) is IDP. Hence, the following statement proved in Section 3 is stronger than the afore-mentioned result of [7, 11, 16]:

Theorem 1.6. *Let Q be lattice polygon, and P a weak Minkowski summand of Q . Then the Cayley sum $\text{Cay}(P, Q)$ has a unimodular cover.*

This has the following consequence, also proved in Section 3. Here a *prismatoid* is a polytope whose vertices all lie in two parallel facets.

Corollary 1.7. *Every smooth 3-dimensional lattice prismatoid has a unimodular cover.*

Let us mention that recent work of Gubeladze [8] shows another class of 3-polytopes admitting unimodular covers: the convex hulls of all lattice points inside an ellipsoid; these had previously been shown to be IDP by Bruns, Gubeladze and Michalek [4]. To date there are no known examples of IDP 3-polytopes without a unimodular cover, although such polytopes exist in higher dimension [2].

2. PARALLELEPIPEDS

The main tool for the proof of Theorem 1.2 is what we call the parallelepiped circumscribed to a given tetrahedron, defined as follows:

Definition 2.1. Let T be a tetrahedron with vertices p_1, p_2, p_3 , and p_4 . Consider the points $q_i = \frac{1}{2}(p_1 + p_2 + p_3 + p_4) - p_i$ for each $i \in \{1, 2, 3, 4\}$, and let

$$C(T) = \text{conv}(p_i, q_i : i \in \{1, 2, 3, 4\}).$$

$C(T)$ is a parallelepiped with facets $\text{conv}(p_i, p_j, q_k, q_l)$ for all choices of $\{i, j, k, l\} = \{1, 2, 3, 4\}$. We call it the *parallelepiped circumscribed* to T .

For each $i \in \{1, 2, 3, 4\}$, let $T_i = \text{conv}(q_i, p_j, p_k, p_l)$, with $\{i, j, k, l\} = \{1, 2, 3, 4\}$; we call the T_i *corner tetrahedra* of $C(T)$. Together with T they triangulate $C(T)$.

Modulo an affine transformation, the situation of T and $C(T)$ is exactly that of the regular tetrahedron inscribed in a cube; see Figure 1.




FIGURE 1. In red we have a tetrahedron T , in black its circumscribed parallelepiped $C(T)$, and in blue the corner simplex T_4 .

Observe that the points q_i need not be lattice points. However, the following lemma shows that we can find lattice points in each corner tetrahedron.

Lemma 2.2. Let $T = \text{conv}\{p_1, p_2, p_3, p_4\}$ be an empty lattice tetrahedron that is not unimodular. Let $C(T)$ be the parallelepiped circumscribed to T and let T_1, T_2, T_3 and T_4 be the corresponding corner tetrahedra in $C(T)$. Then, every T_i contains at least one lattice point different from $\{p_1, \dots, p_4\}$.

Proof. By White's classification of empty tetrahedra ([20], see also, e. g. [10, Sect. 4.1]), there is no loss of generality in assuming $T = \text{conv}(p_1, p_2, p_3, p_4)$ with

$$p_1 = (0, 0, 0), \quad p_2 = (1, 0, 0), \quad p_3 = (0, 0, 1), \quad p_4 = (a, b, 1).$$

where $b \geq 2$ is the (normalized) volume of T , and $a \in \{1, \dots, b-1\}$ satisfies $\gcd(a, b) = 1$. This gives

$$\begin{aligned} q_1 &= \left(\frac{1+a}{2}, \frac{b}{2}, 1\right), & q_2 &= \left(\frac{a-1}{2}, \frac{b}{2}, 1\right), \\ q_3 &= \left(\frac{1+a}{2}, \frac{b}{2}, 0\right), & q_4 &= \left(\frac{1-a}{2}, -\frac{b}{2}, 0\right). \end{aligned}$$

Then, the inequalities $b \geq 1 + a \geq 2$ imply:

$$u := (1, 1, 0) \in \text{conv}(p_1, p_2, q_3) \subset T_3, \quad v := (0, -1, 0) \in \text{conv}(p_1, p_2, q_4) \subset T_4.$$

Observe that $u + v = p_1 + p_2 = q_3 + q_4$. Now, this implies that the quadrilateral $\text{conv}(p_1, q_4, p_2, q_3)$ contains a fundamental domain for the lattice $\mathbb{Z}^2 \times \{0\}$. Hence, its translate $\text{conv}(q_2, p_3, q_1, p_4)$ contains a fundamental domain for $\mathbb{Z}^2 \times \{1\}$ and,

in particular, it contains at least one lattice point other than p_3 and p_4 . By central symmetry around its center $(\frac{a}{2}, -\frac{b}{2}, 1)$, $\text{conv}(q_2, p_3, q_1, p_4)$ must contain lattice points in both triangles $\text{conv}(q_2, p_3, p_4) \subset T_1$ and $\text{conv}(q_1, p_3, p_4) \subset T_2$. \square

Lemma 2.3. *Let P be a lattice parallelepiped and let $T \subset P$ be a tetrahedron. Then, at least one of the four corner tetrahedra T_i of the circumscribed parallelepiped $C(T)$ is fully contained in P .*

Proof. Let us denote the vertices of T by p_1, p_2, p_3, p_4 and the vertices of $C(T)$ not in T by q_1, q_2, q_3, q_4 , with the conventions of Definition 2.1.

We call *band* any region of the form $f^{-1}([\alpha, \beta])$ for some linear functional $f \in (\mathbb{R}^3)^* \setminus \{0\}$ and closed interval $[\alpha, \beta] \subset \mathbb{R}$. We claim that any band containing T must contain at least three of the q_i s. This claim implies that the parallelepiped P , which is the intersection of three bands, contains at least one of the q_i s and hence it fully contains the corresponding T_i .

To prove the claim, suppose that $q_1 \notin B := f^{-1}([\alpha, \beta])$ for a certain band $B \supset T$. Without loss of generality, say $f(q_1) < \alpha$. Then the equalities $q_1 + q_i = p_j + p_k$ and $q_1 + p_1 = q_i + p_i$, where $\{i, j, k\} = \{2, 3, 4\}$, respectively give:

- (1) $f(q_i) = f(p_j + p_k - q_1) = f(p_j) + f(p_k) - f(q_1) > 2\alpha - \alpha = \alpha$,
- (2) $f(q_i) = f(q_1 + p_1 - p_2) = f(q_1) + f(p_1) - f(p_i) < \alpha + \beta - \alpha = \beta$,

so that $q_i \in B$ for $i \in \{2, 3, 4\}$. This concludes the proof of the claim, and of the lemma. \square

Corollary 2.4. *Let T be an empty lattice tetrahedron contained in a lattice parallelepiped P . Then, T can be covered by unimodular tetrahedra contained in P .*

Proof. We proceed by induction on the (normalized) volume of T , which is a positive integer. If this volume equals 1 then T is unimodular and there is nothing to prove, so we assume T is not unimodular. Let p_1, p_2, p_3, p_4 denote the vertices of T .

Lemma 2.3 guarantees that one of the corner tetrahedra T_i of the parallelepiped $C(T)$ is contained in P . Without loss of generality, suppose $T_4 = \text{conv}(p_1, p_2, p_3, q_4)$ is in P . By Lemma 2.2, we know that T_4 contains a lattice point other than the p_i s, which we denote by u . Then $S = \text{conv}(T \cup \{u\})$ can be triangulated in two different ways: $S = T \cup T'_4$, where $T'_4 = \text{conv}(p_1, p_2, p_3, u) \subseteq T_4$ and $S = S_1 \cup S_2 \cup S_3$, with

$$S_1 = \text{conv}(p_2, p_3, p_4, u), S_2 = \text{conv}(p_1, p_3, p_4, u), S_3 = \text{conv}(p_1, p_2, p_4, u).$$

Each of the tetrahedra S_i has lattice volume strictly smaller than T because, for each i , p_i is the unique point of $C(T)$ maximizing the distance to the opposite facet $\text{conv}(p_j, p_k, p_l)$ of T . Thus, S_1, S_2 and S_3 cover T and have volume strictly smaller than T . The S_i may not be empty, but we can triangulate them into empty tetrahedra, which by inductive hypothesis can be covered unimodularly. \square

Proof of Theorem 1.2. Arbitrarily triangulate the parallelepiped into empty lattice tetrahedra and apply Corollary 2.4 to these tetrahedra. \square

Let us say that a lattice 3-polytope P has the *circumscribed parallelepiped property* if it satisfies the conclusion of Lemma 2.3: “for every empty tetrahedron T contained in P at least one of the four corner tetrahedra in $C(T)$ is contained in P ”. If this holds then P has a unimodular cover, since then the proofs of Corollary 2.4 and Theorem 1.2 work for P . In turn, our proof that parallelepipeds have the property (Lemma 2.3) is based on the fact that they have only three (pairs of) normal vectors. In the following two examples we show a smooth 3-polytope and two 3-polytopes with four normal vectors that do not have the property. The latter are not IDP:

used i instead of 2, to shorten proof

— P.

Example 2.5 (A smooth 3-polytope without the circumscribed parallelepiped property). Let P be the Cayley embedding of a long horizontal rectangle and a long vertical rectangle. That is, $P = \text{conv}([0, a] \times [0, 1] \times \{0\} \cup [0, 1] \times [0, b] \times \{1\})$, for big a and b . This is smooth and contains a big empty tetrahedron T with vertices $(0, 0, 0)$, $(a, 1, 0)$, $(0, 0, 1)$, $(1, b, 1)$ which occupies most of its volume. In particular, none of the corner tetrahedra of T is contained in P .

More explicitly, the the remaining vertices q_i of the circumscribed parallelepiped are $(\frac{a+1}{2}, \frac{b+1}{2}, 1)$, $(\frac{a+1}{2}, \frac{b+1}{2}, 0)$, $(\frac{1-a}{2}, \frac{b-1}{2}, 1)$, $(\frac{a-1}{2}, \frac{1-b}{2}, 1)$. None of these points are contained in P , and therefore none of the corner tetrahedra are either.

Example 2.6 (Non-IDP polytopes with four facet directions). The following triangular prism P and centrally symmetric octahedron Q are not IDP:

$$(3) \quad P = \text{conv}((0, 1, 1), (1, 0, 1), (1, 1, 0), (-1, 0, 0), (0, -1, 0), (0, 0, -1)),$$

$$(4) \quad Q = \text{conv}((0, 1, 1), (1, 0, 1), (1, 1, 0), (0, -1, -1), (-1, 0, -1), (-1, -1, 0)).$$

Indeed, in both cases the point $(1, 1, 1)$ lies in the second dilation but is not the sum of two lattice points in the polytope.

An affirmative answer to the following question (weaker than the circumscribed parallelepiped property) would still imply that smooth 3-polytopes can be unimodularly covered and, hence, Conjecture 1.1(1) in dimension three:

Question 2.7. If T is an empty tetrahedron contained in a smooth 3-polytope P , can one guarantee that there is a lattice point of P in the circumscribed parallelepiped of T (apart of the vertices of T)?

3. CAYLEY SUMS

Let P and Q be two lattice polytopes in \mathbb{R}^d . We do not require them to be full-dimensional, but we assume their Minkowski sum is. Remember that the *Minkowski sum* $P + Q$ and the *Cayley sum* of P and Q are defined as:

$$P + Q := \{p + q \in \mathbb{R}^d : p \in P, q \in Q\} \subset \mathbb{R}^d,$$

$$\text{Cay}(P, Q) = \text{conv}(P \times \{0\} \cup Q \times \{1\}) \subset \mathbb{R}^{d+1}.$$

The so-called *Cayley Trick* is the isomorphism

$$2 \text{Cay}(P, Q) \cap (\mathbb{R}^d \times \{1\}) = (P + Q) \times \{1\} \cong P + Q,$$

which easily implies:

Proposition 3.1 (see, e.g. [19, Thm. 0.4]). *If $\text{Cay}(P, Q)$ is IDP then the pair (P, Q) is IDP.*

The Cayley Trick also provides the following canonical bijections:

polyhedral subdivisions of $\text{Cay}(P, Q)$	\leftrightarrow	mixed subdivisions of $P + Q$
triangulations of $\text{Cay}(P, Q)$	\leftrightarrow	fine mixed subdivisions of $P + Q$
unimodular simplices in $\text{Cay}(P, Q)$	\leftrightarrow	unimodular prod-simplices in $P + Q$.

See [6] for more details on the Cayley Trick and on triangulations and polyhedral subdivisions of polytopes. In fact these bijections can be taken as definitions of the objects in the right-hand sides. In particular, we call *prod-simplices* in $P + Q$ the Minkowski sums $T_1 + T_2$ where $T_1 \subset P$ and $T_2 \subset Q$ are simplices with complementary affine spans. A prod-simplex is *unimodular* if the edge vectors from any vertex of T_1 and from any vertex of T_2 form a unimodular basis.

We now turn our attention to $d = 2$, in order to prove Theorem 1.6. A triangulation of $\text{Cay}(P, Q) \subset \mathbb{R}^3$ consists of tetrahedra of types $(1, 3)$, $(2, 2)$ and $(3, 1)$, where the type denotes how many vertices they have in P and in Q . Empty tetrahedra of types $(1, 3)$ or $(3, 1)$, which are Cayley sums of an empty (hence unimodular)

Added the last part to answer comment Y.19 — G.

edited it — P.

shortened — P.

We just called the pair IDP in the introduction... Which notation do we prefer? — G.

we stick to just IDP — P.

triangle in P and a point in Q , or viceversa, are automatically unimodular. The case that we need to study are therefore tetrahedra of type $(2, 2)$, which are Cayley sums of a segment $p \subset P$ and a segment $q \subset Q$. These correspond to prod-simplices of two segments in $P + Q$, which are *parallelograms*. The following lemma, whose proof we postpone to Section 4, is crucial to understand how to unimodularly cover these tetrahedra.

We use the following conventions: if a, b are points, we denote by $[a, b]$ and (a, b) respectively the closed and open line segments with endpoints a, b . Given a segment $s = [a, b]$, we denote \vec{s} the vector $b - a$ and denote $\langle \vec{s} \rangle$ the line spanned by \vec{s} . A lattice parallelogram $p + q$ is called *unimodular* if it is a fundamental domain for the lattice. Equivalently, if $\text{Cay}(p, q)$ is a unimodular tetrahedron.

added def of unimodular parallelogram — P.

referee X suggests to give an “example” of the lemma, but I am not sure what could be helpful — P.

Lemma 3.2. *Let $Q \subset \mathbb{R}^2$ be a two-dimensional lattice polytope and $P \subset \mathbb{R}^2$ a weak Minkowski summand of it. Let $p = [p_1, p_2] \subset P$ and $q = [q_1, q_2] \subset Q$ be two primitive and non-parallel lattice segments, and let $\langle \vec{p} \rangle$ and $\langle \vec{q} \rangle$ be the lines spanned by them. If the parallelogram $p + q$ is not unimodular, then at least one of the regions*

$$((p_1, p_2) + \langle \vec{q} \rangle) \cap P, \quad \text{and} \quad ((q_1, q_2) + \langle \vec{p} \rangle) \cap Q$$

contains a lattice point.

See Figure 2 for an illustration of the two regions in the statement, which we call *strips*. In this figure and the forthcoming ones in Section 4 we draw p as a vertical segment and q as a horizontal one for convenience. This is always possible via a linear transformation (which of course changes the lattice; in the proof we do not assume the lattice to be \mathbb{Z}^2).

moved this remark here, since it already affects this proof — P.




FIGURE 2. The strips of Lemma 3.2

Corollary 3.3. *Let T be an empty lattice tetrahedron contained in the Cayley sum $\text{Cay}(P, Q)$, where Q is a lattice polygon and P is a weak Minkowski summand of Q . Then, T can be covered by unimodular tetrahedra contained in $\text{Cay}(P, Q)$.*

Proof. The proof is by induction on the normalized volume of T , which we assume to be at least 2. This implies that T is of type $(2, 2)$, since empty tetrahedra of types $(1, 3)$ and $(3, 1)$ are unimodular. Thus, T is the Cayley sum of primitive segments $p = [p_1, p_2] \subset P$ and $q = [q_1, q_2] \subset Q$. Let u be the lattice point whose existence is guaranteed by Lemma 3.2. Assume (the other case is similar) that

$$u \in ((p_1, p_2) + \langle \vec{q} \rangle) \cap P,$$

and call t the triangle $t = \text{conv}(u, p_1, p_2) \subset P$.

Let us denote $\tilde{u}, \tilde{p}_1, \tilde{p}_2, \tilde{q}_1, \tilde{q}_2$ the points corresponding to u, p_1, p_2, q_1, q_2 in $\text{Cay}(P, Q)$. That is, $\tilde{p}_i = p_i \times \{1\}$, $\tilde{q}_i = q_i \times \{0\}$, and $\tilde{u} = u \times \{1\}$. Observe that

to match the figure, P is now at height 1 instead of 0 — G.

the assumption $u \in ((p_1, p_2) + \langle \vec{q} \rangle)$ implies that one of the segments $[u, q_i]$ crosses the interior of one of the triangles $\text{conv}(p_1, p_2, q_j)$, where $\{i, j\} = \{1, 2\}$. Without loss of generality assume that $[\tilde{u}, \tilde{q}_2]$ crosses $\text{conv}(\tilde{p}_1, \tilde{p}_2, \tilde{q}_1)$, as in Figure 3.




FIGURE 3. $[\tilde{u}, \tilde{q}_2]$ intersects $\text{conv}(\tilde{p}_1, \tilde{p}_2, \tilde{q}_1)$

In turn, this means that the polytope $\text{conv}(\tilde{u}, \tilde{p}_1, \tilde{p}_2, \tilde{q}_1, \tilde{q}_2) = \text{Cay}(t, q)$ has the following two triangulations:

$$\begin{aligned} \mathcal{T}^+ &:= \{\text{Cay}(p, q), \text{Cay}(t, \{q_2\})\}, \\ \mathcal{T}^- &:= \{\text{Cay}([p_1, u], q), \text{Cay}([p_2, u], q), \text{Cay}(t, \{q_1\})\}. \end{aligned}$$

The tetrahedra $\text{Cay}(t, \{q_1\})$ and $\text{Cay}(t, \{q_2\})$ are unimodular, which implies that $T = \text{Cay}(p, q)$ has volume equal to the sum of the volumes of $\text{Cay}([p_1, u], q)$ and $\text{Cay}([p_2, u], q)$. In particular, we have covered T by the three tetrahedra in \mathcal{T}^- , which are of smaller volume and hence have unimodular covers by induction hypothesis. \square

Proof of Theorem 1.6. Arbitrarily triangulate $\text{Cay}(P, Q)$ into empty lattice tetrahedra and apply Corollary 3.3 to these tetrahedra. \square

Let us now show how to derive Corollary 1.7 from this theorem. *Prismatoids* were defined in [17] as polytopes whose vertices all lie in two parallel facets. In particular, a *lattice prismatoid* is any d -polytope $SL(\mathbb{Z}, d)$ -equivalent to one of the form

$$\text{conv}(Q_1 \times \{0\} \cup Q_2 \times \{k\}),$$

where Q_1, Q_2 are lattice $(d-1)$ -polytopes and $k \in \mathbb{Z}_{>0}$. This is almost a generalization of Cayley sums, which would be the case $k = 1$, except the definition of prismatoid requires Q_1 and Q_2 to be full-dimensional, while the Cayley sum only requires this for $Q_1 + Q_2$.

Proposition 3.4. *Let Q_1, Q_2 be two lattice polygons and consider the prismatoid*

$$P := \text{conv}(Q_1 \times \{0\} \cup Q_2 \times \{k\}),$$

with $k \geq 2$. If $P \cap (\mathbb{R}^2 \times \{1\})$ is a lattice polygon then P has a unimodular cover.

Proof. The condition that $P \cap (\mathbb{R}^2 \times \{1\})$ is a lattice polygon implies the same for $P \cap (\mathbb{R}^2 \times \{i\})$, for every i . Indeed, the condition implies that every edge of $\text{Cay}(P, Q)$ of the form $[u \times \{0\}, v \times \{k\}]$ has a lattice point in $\mathbb{R}^2 \times \{i\}$, and hence it has a lattice point in $P \cap (\mathbb{R}^2 \times \{i\})$, for every i .

Observe that for every $i \in \{1, \dots, k-1\}$ the intersection $P \cap (\mathbb{R}^2 \times \{i\})$ has the same normal fan as $Q_1 + Q_2$. Thus, each slice

$$P \cap (\mathbb{R}^2 \times [i-1, i])$$

added tildes to the points
— G.

is a Cayley polytope. For $i \in \{2, \dots, k-1\}$, both bases have the same normal fan (and therefore each is a weak Minkowski summand of the other); for $i \in \{1, k\}$ one base is a weak Minkowski summand of the other. We can therefore apply Theorem 1.6 to each slice and combine the covers thus obtained to get a unimodular cover of P . \square

Proof of Corollary 1.7. The polytope under study satisfies the hypotheses of Proposition 3.4: the smoothness of the prismatoid implies that every edge of the form $[u \times \{0\}, v \times \{k\}]$ has lattice points in all slices. Hence,

$$kP \cap (\mathbb{R}^2 \times \{1\}) = (k-1)Q_1 + Q_2. \quad \square$$

4. PROOF OF LEMMA 3.2

Let f_q be the primitive lattice functional constant on q and f_p the one constant on p . We assume that $f_q(p_1) < f_q(p_2)$ and $f_p(q_1) < f_p(q_2)$.

Since we can perform without loss of generality respective lattice translations to P and to Q , we assume that q_1 is the origin and that $f_q(p_1) = -1$. Observe that then $f_q(p_2)$ must be strictly positive, since $f_q(p_2) = 0$ would imply that $p+q$ is a unimodular parallelogram. Moreover, since p is primitive p_1 and p_2 cannot be in the boundary of $q + \langle \vec{p} \rangle$, which implies that p_1 is the unique lattice point with $f_q(x) = -1$ in $q + \langle \vec{p} \rangle$. Similarly, the unique lattice point in the strip with $f_q(x) = 1$ is $q_1 + q_2 - p_1$.

We let $H_1 = \{f_q(x) \leq 0\}$ and $H_2 = \{f_q(x) \geq 0\}$; similarly let $V_1 = \{f_p(x) \leq f_p(p)\}$ and $V_2 = \{f_p(x) \geq f_p(p)\}$.¹




FIGURE 4. Setup for the proof of Lemma 3.2

Let $w = \text{area}(p+q) \geq 2$, where area denotes the area normalized to a fundamental domain. In what follows, the width of a functional f on a set S , denoted $\text{width}_f(S)$ is defined as the difference $\sup_{x \in S} f(x) - \inf_{x \in S} f(x)$. Then:

$$w = \text{width}_{f_q}(p + \langle \vec{q} \rangle) = \text{width}_{f_q}(p) = \text{width}_{f_p}(q) = \text{width}_{f_p}(q + \langle \vec{p} \rangle).$$

Proof of Lemma 3.2. Suppose by contradiction that there is no lattice point as described in the lemma. In particular, no lattice point on the boundary of Q can be in the interior of the strip $q + \langle \vec{p} \rangle$. Thus the boundary of Q contains two primitive segments which each have one vertex on each side of the strip $q + \langle \vec{p} \rangle$; we will call these $b = [b_1, b_2]$, $t = [t_1, t_2]$, with b and t crossing the strip in H_1 and

¹Since in our figures p and q are vertical and horizontal, we use the letters V and H for the half-planes they defined. Similarly, later in the proof we use the letters b , t , r , and l for certain points and segments meaning “bottom”, “top”, “right” and “left”.

rephrased to address X.10
— P.

added footnote
— P.

H_2 respectively and the convention that $f_p(b_2) > f_p(b_1)$ and $f_p(t_2) > f_p(t_1)$. This readily implies

$$(5) \quad \begin{aligned} f_p(t_1) &\leq f_p(q_1), & f_p(t_2) &\geq f_p(q_2), \\ f_p(b_1) &\leq f_p(q_1), & f_p(b_2) &\geq f_p(q_2). \end{aligned}$$

The same holds for P and the strip $p + \langle \vec{q} \rangle$, and we call the segments $\ell = [l_1, l_2]$ and $r = [r_1, r_2]$, with ℓ and r crossing the strip $p + \langle \vec{q} \rangle$ in V_1 and V_2 respectively. The only difference is that in the case that P is one dimensional we have $\ell = r = p$. Again we have

$$(6) \quad \begin{aligned} f_q(l_1) &\leq f_q(p_1), & f_q(l_2) &\geq f_q(p_2), \\ f_q(r_1) &\leq f_q(p_1), & f_q(r_2) &\geq f_q(p_2). \end{aligned}$$

Observe that a priori one of ℓ and r can coincide with p , if p is on the boundary of P , and similarly one of t, b might be q , if q is on the boundary of Q .

Claim 4.1. *The following inequalities hold,*

$$\text{width}_{f_q}(\ell), \text{width}_{f_q}(r), \text{width}_{f_p}(t), \text{width}_{f_p}(b) \geq w.$$

Each inequality is strict, unless the segment in question coincides with p or q .

Proof. The inequality $\geq w$ follows in each case from (6) and (5).

If one of the inequalities, say the one for ℓ , is not strict, then ℓ has one endpoint on each of the boundary lines of $(p + \langle \vec{q} \rangle)$. Unless $\ell = p$, one of the endpoints of ℓ is not an endpoint of p , say $l_1 \neq p_1$. Thus the triangle $T = \text{conv}(p_2, p_1, l_1)$ is contained in P and its edge $[p_1, l_1]$ is an integer dilation of q .

Since T contains p and a copy of q , its area (normalized to a fundamental domain) is $w/2 \geq 1$, and by Pick's theorem it must contain a lattice point other than its vertices. Since p and q are primitive, this lattice point must lie in the interior of the strip. \square

Claim 4.2. *$f_q(b_2 - b_1)$ and $f_q(t_2 - t_1)$ are non-zero and have the same sign. That is, f_q achieves its maximum over b and over t on the same halfplane V_1 or V_2 .*

Proof. Both t and b must cross the interior of p , or otherwise p_1 or p_2 are the lattice points we are looking for in Q . To seek a contradiction assume, as in Figure 5, that

$$f_q(t_2 - t_1) \leq 0 \leq f_q(b_2 - b_1).$$

That is, f_q decreases (perhaps weakly) along t and increases along b , as f_p increases on both. This implies that $Q \cap V_2$ is contained in the open strip $\{f_q(p_1) < f_q(x) < f_q(p_2)\}$, of width w . This strip cannot contain a translated copy of r , since $\text{width}_{f_q}(r) \geq w$, which gives a contradiction: since P is a weak Minkowski summand of Q , Q must have edges parallel to r both in V_1 and V_2 . \square




FIGURE 5. Illustration of the proof of Claim 4.2

not sure why we use ℓ for the segment and l for the endpoints... — P.

added an explanation here, according to the suggestion Y.32 — G.

simplified proof — P.

We assume without loss of generality that the maximum on t (and hence on b) is achieved in V_2 , that is to say, f_p and f_q increase in the same direction along t (and hence along b). Otherwise the following considerations can be applied to V_1 .

Claim 4.3. *Assume without loss of generality that b and t either are parallel or their affine spans cross in V_2 (if they cross in V_1 , the same claim can be reworded for V_1 and l). Then,*

- (1) *The intersection of Q with any line parallel to p in V_2 has width with respect to f_q strictly smaller than w .*
- (2) *$f_p(r_2) > f_p(r_1)$, that is, f_p achieves its maximum over r in H_2 .*

Proof. Both t and b must intersect p , as said in the proof of Claim 4.2. Their intersections with p are thus endpoints of a segment of width with respect to f_q less than w , the width of p . Since t and b cross in V_2 , the same is true for any segment parallel to p contained in $Q \cap V_2$.




FIGURE 6. Illustration of the proof of Claim 4.3

For part (2), recall that by Claim 4.1, $\text{width}_{f_q}(r) \geq w$. If $f_p(r_2) \leq f_p(r_1)$, it would be impossible to fit a translated copy r' of r in the correct side of Q : since f_q increases along t , r' has width in direction f_q smaller than the segment parallel to p with endpoint r'_1 . This segment by part (1) has width less than w in direction q , a contradiction.

simplified the proof as suggested in Y.38

— G.

□

The last two claims can be summarized as saying that in the pictures b , t and r have positive slope. Observe that this implies that q is not in the boundary of Q and $p \neq r$, so both P and Q are full dimensional.

Let g be the primitive lattice functional constant on $[p_1, q_2]$ (and therefore constant also on $[q_1, q_1 + q_2 - p_1]$). By the assumption on p_1 , the values of g on these segments differ by 1. We choose the sign of g so that

$$g([p_1, q_2]) = g([q_1, q_1 + q_2 - p_1]) - 1.$$

changed the +1 to a -1, so that the inequalities in the following claim are the right way around—we had flipped everything

— G.

Claim 4.4. $g(t_1) > g(t_2)$, $g(b_1) > g(b_2)$, and $g(r_1) < g(r_2)$.

Proof. Since b and t must respectively separate p_1 and $q_1 + q_2 - p_1$ from the other two vertices of the parallelogram $\text{conv}(q_1, p_1, q_2, q_1 + q_2 - p_1)$, they must respectively intersect its (parallel) edges $[p_1, q_2]$ and $[q_1, q_1 + q_2 - p_1]$, which implies the stated inequalities for b and t . The same argument applied to the parallelogram $\text{conv}(p_1, q_2, p_2, p_1 + p_2 - q_2)$, yields the inequalities for r . □

We are now ready to show a contradiction. Since the normal fan of Q refines that of P , Q must have an edge r' which is a translated copy of r . Let r'_1 and r'_2 be its endpoints.

Changed the proof from here

— G.




FIGURE 7. Illustration of the proof of Claim 4.4

Consider the segment s contained in $r'_1 + \langle \vec{p} \rangle$ with endpoints $s_1 = r'_1$ on d and s_2 on the line spanned by t . The width of s with respect to g is $g(s_2) - g(s_1) < 1$, because s is shorter than $Q \cap p$, which is strictly contained between the consecutive parallel lattice lines through p_1 and q_2 and through q_1 and $q_1 + q_2 - p_1$ where g is constant.

We now observe that $g(r_2) < g(s_2)$, because r_2 lies to the right of s_2 (r has positive slope) and below the line spanned by t , and g decreases moving to the right along t (Claim 4.4). Thus $g(r_2) - g(r_1) = g(r'_2) - g(r'_1) < g(s_2) - g(s_1) < 1$, that is, $g(r_2) = g(r_1)$. This contradicts Claim 4.4. \square

REFERENCES

- [1] Matthias Beck, Christian Haase, Akihiro Higashitani, Johannes Hofscheier, Katharina Jochemko, Lukas Katthän, Mateusz Michałek, Smooth Centrally Symmetric Polytopes in Dimension 3 are IDP, *Ann. Combin.*, 23 (2019), 255–262. doi:[10.1007/s00026-019-00418-x](https://doi.org/10.1007/s00026-019-00418-x)
- [2] Winfried Bruns and Joseph Gubeladze, Normality and covering properties of affine semigroups, *J. Reine Angew. Math.* 510 (1999), 151–178. doi:[10.1515/crll.1999.044](https://doi.org/10.1515/crll.1999.044)
- [3] Winfried Bruns and Joseph Gubeladze, *Polytopes, rings, and k-theory*, Monographs in Mathematics, Springer-Verlag, 2009, XIV, 461 p. 52 illus.
- [4] Winfried Bruns, Joseph Gubeladze, and Mateusz Michałek. Quantum Jumps of Normal Polytopes. *Discrete Comput. Geom.*, 56(1) (2016), 181–215. doi:[10.1007/s00454-016-9773-7](https://doi.org/10.1007/s00454-016-9773-7)
- [5] David A. Cox, John B. Little, and Hal K. Schenck. *Toric Varieties*. AMS, Providence, 2011.
- [6] J. A. De Loera, J. Rambau, F. Santos *Triangulations: Structures for Algorithms and Applications*, 539 pp. Algorithms and Computation in Mathematics, Vol. 25, Springer-Verlag. ISBN: 978-3-642-12970-4
- [7] Najmuddin Fakhruddin. Multiplication maps of linear systems on smooth projective toric surfaces. Preprint, [arXiv:math/0208178](https://arxiv.org/abs/math/0208178), August 2002.
- [8] Joseph Gubeladze, Normal polytopes and ellipsoids, *Elect. J. Combin.*, 28:4 (2021), P4.7. doi:[10.37236/10338](https://doi.org/10.37236/10338)
- [9] Christian Haase and Jan Hofmann, Convex-normal (Pairs of) Polytopes, *Canad. Math. Bull.* 60:3 (2017), 510–521. doi:[10.4153/CMB-2016-057-0](https://doi.org/10.4153/CMB-2016-057-0)
- [10] Christian Haase, Andreas Paffenholz, Lindsay C. Piechnik, Francisco Santos. Existence of unimodular triangulations - positive results. *Mem. Amer. Math. Soc.*, Vol. 270 1N. 321, American Math. Society, 2021. DOI: doi:[10.1090/memo/1321](https://doi.org/10.1090/memo/1321)
- [11] Christian Haase, Benjamin Nill, Andreas Paffenholz, and Francisco Santos, Lattice points in Minkowski sums, *Electron. J. Combin.*, 15:1 (2008), Note 11, 5 pp. doi:[10.37236/886](https://doi.org/10.37236/886)
- [12] Jean-Michel Kantor and Karanbir S. Sarkaria. On primitive subdivisions of an elementary tetrahedron, *Pacific J. Math.* **211** (2003), no. 1, 123–155. doi:[10.2140/pjm.2003.211.123](https://doi.org/10.2140/pjm.2003.211.123)
- [13] Mini-workshop: Ehrhart Quasipolynomials: Algebra, Combinatorics, and Geometry, Oberwolfach Rep. 1 (2004), no. 3, 2071–2101, Abstracts from the mini-workshop held August 15–21, 2004, Organized by Jesús A. De Loera and Christian Haase, Oberwolfach Reports. Vol. 1, no. 3. doi:[10.14760/OWR-2004-39](https://doi.org/10.14760/OWR-2004-39)
- [14] Mini-workshop: Projective normality of smooth toric varieties, Oberwolfach Rep. 4 (2007), no. 39/2007, Abstracts from the mini-workshop held August 12–18, 2007. Organized by Christian Haase, Takayuki Hibi and Diane MacLagan. doi:[10.14760/OWR-2007-39](https://doi.org/10.14760/OWR-2007-39)
- [15] Tadao Oda. Problems on Minkowski sums of convex lattice polytopes. Abstract submitted at the Oberwolfach Conference "Combinatorial Convexity and Algebraic Geometry" 26.10–01.11, 1997. Available as arXiv preprint [arXiv:0812.1418](https://arxiv.org/abs/0812.1418), 2008.

- [16] Shoetsu Ogata. Multiplication maps of complete linear systems on projective toric surfaces. *Interdiscip. Inf. Sci.*, 12(2):93–107, 2006. doi:[10.4036/iis.2006.93](https://doi.org/10.4036/iis.2006.93)
- [17] Francisco Santos. A counter-example to the Hirsch Conjecture. *Ann. Math. (2)*, 176 (July 2012), 383–412. doi:[10.4007/annals.2012.176.1.7](https://doi.org/10.4007/annals.2012.176.1.7)
- [18] Francisco Santos and Günter M. Ziegler. Unimodular triangulations of dilated 3-polytopes, *Trans. Moscow Math. Soc.* (2013), 293–311. doi:[10.1007/s00493-014-2959-9](https://doi.org/10.1007/s00493-014-2959-9)
- [19] Akiyoshi Tsuchiya. Cayley sums and Minkowski sums of 2-convex-normal lattice polytopes. Preprint, [arXiv:1804.10538](https://arxiv.org/abs/1804.10538), April 2018.
- [20] G. K. White. Lattice tetrahedra. *Canadian J. Math.* 16 (1964), 389–396. doi:[10.4153/CJM-1964-040-2](https://doi.org/10.4153/CJM-1964-040-2)