

# Two Sum

Try to solve the Two Sum problem.

We'll cover the following ^

- Statement
- Examples
- Understand the problem
- Try it yourself

## Statement

For the given array of integers `arr` and a target `t`, you have to identify the two indices that add up to generate the target `t`. Moreover, you can't use the same index twice, and there will be only one solution.

**Note:** The order of output doesn't matter.

### Constraints:

- $2 \leq arr.length \leq 10^4$
- $-10^9 \leq arr[i] \leq 10^9$
- $-10^9 \leq t \leq 10^9$
- Only one valid answer exists.

## Examples

### Sample example 1

**Input**

arr	1	10	8	4	9
target	17				

-----

**Output**

[2, 4]
--------

1 of 2

### Sample example 2

**Input**

arr	5	12	15	21	6	17
target	33					

-----

**Output**

[1, 3]
--------

?

Tt



## Understand the problem

Let's take a moment to make sure you've correctly understood the problem. The quiz below helps you check if you're solving the correct problem:

### Two Sum

1

What is the output for the following input?

arr = [3, 9, 11, 12, 17, 21]

t = 26

A) [0, 5]

B) [1, 4]

C) [2, 3]

D) [3, 5]

Submit Answer



Question 1 of 3  
0 attempted



Reset Quiz ↻

## Try it yourself

Implement your solution in the following coding playground:

Java



usercode > Solution.java

```
1 import java.util.*;
2 public class Solution{
3     public static int[] twoSum(int[] arr, int t) {
4
5         // Your code will replace this placeholder return statement
6
7         return new int[1];
8     }
9 }
```



Powered by AI

Submit

Test Cases

Results

Case 1

Case 2

Case 3


Input #1

[2,4,6,8,10,19]

Input #2

21

Two Sum

 Hide Hint

You might want to go over the [Knowing What To Track](#) pattern again.

 Back

Maximum Subarray

Next 

Find Minimum in Rotat...

☒ Mark as Completed