

Basic Engineering Mathematics

DRAFT


July 10, 2017



This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/> or send a letter to Creative Commons, PO Box 1866, Mountain View, CA 94042, USA.

Contents

1	Basic Algebra	5
1.1	Law of Negatives	5
1.2	Law of Quotients	5
2	Quadratic Equations	6
3	Logarithmic Functions	7
3.1	Properties of Logarithmic Functions	7
4	Laws of Exponents	8
4.1	Rational Number Exponents	8
5	Complex Numbers	9
5.1	De Moivre's Theorem	9
6	Permutations and Combinations	10
6.1	Binomial Theorem	10
6.2	Summation of Series	10
7	Trigonometry	11
7.1	Trigonometric Identities	11
7.2	Trigonometric ratios of allied angles	11
7.2.1	First Quadrant	12
7.2.2	Second Quadrant	12
7.2.3	Third Quadrant	12
7.2.4	Fourth Quadrant	12
7.3	Compound Angle Formulae	13
7.4	Sum or Difference Formulae	13
7.5	Half Angle Formulae	13
7.6	Sub Multiple Angle Formulae	13
7.7	Triple Angle Formulae	14
7.8	Product Formulae	14
8	Inverse Trigonometric Functions	15
8.1	Properties of Inverse Trigonometric Functions	15
8.2	Expressions and suggested substitutions	16
9	Hyperbolic Functions	17
9.1	Properties of Hyperbolic Functions	17



*“Dedicated to the
mathematicians of
India, who gifted the
world an abundance of
knowledge and a spirit
of enquiry”.*



1 Basic Algebra

$$(a + b)^2 = a^2 + 2ab + b^2 \quad (1)$$

$$(a - b)^2 = a^2 - 2ab + b^2 \quad (2)$$

$$(a + b)(a - b) = a^2 - b^2 \quad (3)$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3 = a^3 + b^3 + 3ab(a + b) \quad (4)$$

$$(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3 = a^3 - b^3 - 3ab(a - b) \quad (5)$$

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2) \quad (6)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2) \quad (7)$$

$$a^4 - b^4 = (a + b)(a - b)(a^2 + b^2) \quad (8)$$

$$a^2 + b^2 = (a + b)^2 - 2ab \quad (9)$$

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac) \quad (10)$$

1.1 Law of Negatives

$$\begin{aligned} -(-a) &= a \\ (-a)(-b) &= ab \\ -ab &= (-a)(b) = a(-b) = -(-a)(-b) \\ a - b &= a + (-b) \end{aligned}$$

1.2 Law of Quotients

$$\begin{aligned} -\frac{a}{b} &= \frac{(-a)}{b} = \frac{a}{(-b)} = -\frac{(-a)}{(-b)} \\ \frac{a}{b} &= \frac{c}{d} \quad \text{if} \quad ad = bc \\ \frac{a}{b} &= \frac{ka}{kb} \end{aligned}$$

2 Quadratic Equations

The quadratic equation

$$ax^2 + bx + c = 0 \tag{11}$$

where a, b and c are constants and $a \neq 0$, has two solutions for the variable x :

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \tag{12}$$

If the *discriminant* Δ with

$$\Delta = b^2 - 4ac = 0 \tag{13}$$

then the equation (11) has real and equal roots given by:

$$x = \frac{-b}{2a} \tag{14}$$

Sum of the roots:

$$x_1 + x_2 = \frac{-b}{a} \tag{15}$$

Product of the roots:

$$x_1.x_2 = \frac{c}{a} \tag{16}$$

3 Logarithmic Functions

For any two variables x and y related as:

$$y = a^x \quad (17)$$

Where a is a constant. We define the logarithmic function as:

$$\log_a y = x \quad (18)$$

3.1 Properties of Logarithmic Functions

$$\log_a mn = \log_a m + \log_a n \quad (19)$$

$$\log_a \frac{m}{n} = \log_a m - \log_a n \quad (20)$$

$$\log_a m^n = n \cdot \log_a m \quad (21)$$

$$\log_a a = 1 \quad (22)$$

$$\log_a 1 = 0 \quad (23)$$

$$\log_a b = \frac{1}{\log_b a} \quad (24)$$

$$a^x = e^{x \cdot \log a} \quad (25)$$

$$\log_a X = \frac{\ln X}{\ln a} \quad (26)$$

4 Laws of Exponents

$$x^a \cdot x^b = x^{a+b} \quad (27)$$

$$\frac{x^a}{x^b} = x^{a-b} \quad (28)$$

$$(x^a)^b = x^{ab} \quad (29)$$

$$x^a = y^b \implies a = b \quad (30)$$

$$x^{-a} = \frac{1}{x^a} \quad (\text{for } a \text{ being any non zero real number}) \quad (31)$$

$$x^0 = 1 \quad (32)$$

$$\sqrt[n]{\frac{x}{y}} = \frac{\sqrt[n]{x}}{\sqrt[n]{y}} \quad (33)$$

$$\sqrt[n]{xy} = \sqrt[n]{x} \cdot \sqrt[n]{y} \quad (34)$$

$$\left(\frac{x}{y}\right)^{-m} = \left(\frac{y}{x}\right)^m \quad (35)$$

$$\frac{x^{-n}}{y^{-m}} = \frac{y^m}{x^n} \quad (36)$$

4.1 Rational Number Exponents

The principle n th root of $x > 0$ is defined as:

$$x^{1/n} = \begin{cases} \text{is a unique real number } y \text{ if } n \text{ is odd or even} \\ 0, \text{ if } x = 0 \\ \text{not a real number if } x < 0 \end{cases}$$
$$(x^{1/n})^n = \begin{cases} x, \text{ if } n \text{ is odd/even and } x \text{ is positive} \\ |x|, \text{ if } n \text{ is even and } x \text{ is negative} \end{cases}$$

5 Complex Numbers

A number of the form $z = x + iy$ consisting of a real part x , denoted by $Re(z)$ and imaginary part y , denoted by $Im(z)$ is known as a complex number.

$$x = r \cos \theta \tag{37}$$

$$y = r \sin \theta \tag{38}$$

$$r = \sqrt{x^2 + y^2} \tag{39}$$

$$\tan \theta = \frac{y}{x} \implies \theta = \tan^{-1} \frac{y}{x} \tag{40}$$

$$e^{i\theta} = \cos \theta + i \sin \theta \tag{41}$$

$$e^{-i\theta} = \cos \theta - i \sin \theta \tag{42}$$

5.1 De Moivre's Theorem

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta) \tag{43}$$

6 Permutations and Combinations

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2)(n-3)\dots(n-(r+1))}{r!} \quad (44)$$

Note that:

$$\binom{n}{0} = 1 \qquad \binom{n}{1} = n \qquad 0! = 1$$

6.1 Binomial Theorem

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \binom{n}{2}x^{n-2}a^2 + \binom{n}{3}x^{n-3}a^3 + \binom{n}{4}x^{n-4}a^4 + \dots\dots\dots a^n \quad (45)$$

6.2 Summation of Series

$$\sum_{n=1}^{\infty} n = 1 + 2 + 3\dots\dots = \frac{(n)(n+1)}{2} \quad (46)$$

$$\sum_{n=1}^{\infty} n^2 = 1^2 + 2^2 + 3^2\dots\dots = \frac{(n)(n+1)(2n+1)}{6} \quad (47)$$

$$\sum_{n=1}^{\infty} n^3 = 1^3 + 2^3 + 3^3\dots\dots = \frac{(n)^2(n+1)^2}{4} \quad (48)$$

7 Trigonometry

$$\begin{aligned} \sin \theta &= \frac{\text{Perpendicular}}{\text{Hypotenuse}} & \cos \theta &= \frac{\text{Base}}{\text{Hypotenuse}} & \tan \theta &= \frac{\text{Perpendicular}}{\text{Base}} \\ \operatorname{cosec} \theta &= \frac{\text{Hypotenuse}}{\text{Perpendicular}} & \sec \theta &= \frac{\text{Hypotenuse}}{\text{Base}} & \cot \theta &= \frac{\text{Base}}{\text{Perpendicular}} \end{aligned}$$

From the above relations it turns out that,

$$\sin \theta \cdot \operatorname{cosec} \theta = 1$$

$$\cos \theta \cdot \sec \theta = 1$$

$$\tan \theta \cdot \cot \theta = 1$$

7.1 Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \tag{49}$$

$$1 + \sec^2 \theta = \tan^2 \theta \tag{50}$$

$$1 + \operatorname{cosec}^2 \theta = \cot^2 \theta \tag{51}$$

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined
$\cot \theta$	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0
$\sec \theta$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
$\operatorname{cosec} \theta$	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1

Figure 1: Most common trigonometric ratios and values

7.2 Trigonometric ratios of allied angles

- *For n being an odd multiple of $\pi/2$ the function changes to its composite function.*
- *For n being an even multiple of $\pi/2$ the function does not change to its composite, but may change sign.*

7.2.1 First Quadrant

$$\begin{array}{ll}\sin(2\pi + \theta) = \sin \theta & \cos(2\pi + \theta) = \cos \theta \\ \operatorname{cosec}(2\pi + \theta) = \operatorname{cosec} \theta & \sec(2\pi + \theta) = \sec \theta \\ \tan(2\pi + \theta) = \tan \theta & \cot(2\pi + \theta) = \cot \theta\end{array}$$

$$\begin{array}{ll}\sin(\pi/2 - \theta) = \cos \theta & \cos(\pi/2 - \theta) = \sin \theta \\ \operatorname{cosec}(\pi/2 - \theta) = \sec \theta & \sec(\pi/2 - \theta) = \operatorname{cosec} \theta \\ \tan(\pi/2 - \theta) = \cot \theta & \cot(\pi/2 - \theta) = \tan \theta\end{array}$$

7.2.2 Second Quadrant

$$\begin{array}{ll}\sin(\pi/2 + \theta) = \cos \theta & \cos(\pi/2 + \theta) = -\sin \theta \\ \operatorname{cosec}(\pi/2 + \theta) = \sec \theta & \sec(\pi/2 + \theta) = -\operatorname{cosec} \theta \\ \tan(\pi/2 + \theta) = -\cot \theta & \cot(\pi/2 + \theta) = -\tan \theta\end{array}$$

$$\begin{array}{ll}\sin(\pi - \theta) = \sin \theta & \cos(\pi - \theta) = -\cos \theta \\ \operatorname{cosec}(\pi - \theta) = \operatorname{cosec} \theta & \sec(\pi - \theta) = -\sec \theta \\ \tan(\pi - \theta) = -\tan \theta & \cot(\pi - \theta) = -\cot \theta\end{array}$$

7.2.3 Third Quadrant

$$\begin{array}{ll}\sin(\pi + \theta) = -\sin \theta & \cos(\pi + \theta) = -\cos \theta \\ \operatorname{cosec}(\pi + \theta) = \operatorname{cosec} \theta & \sec(\pi + \theta) = -\sec \theta \\ \tan(\pi + \theta) = \tan \theta & \cot(\pi + \theta) = \cot \theta\end{array}$$

$$\begin{array}{ll}\sin(3\pi/2 - \theta) = -\cos \theta & \cos(3\pi/2 - \theta) = -\sin \theta \\ \operatorname{cosec}(3\pi/2 - \theta) = -\sec \theta & \sec(3\pi/2 - \theta) = -\operatorname{cosec} \theta \\ \tan(3\pi/2 - \theta) = \cot \theta & \cot(3\pi/2 - \theta) = \tan \theta\end{array}$$

7.2.4 Fourth Quadrant

$$\begin{array}{ll}\sin(3\pi/2 + \theta) = -\cos \theta & \cos(3\pi/2 + \theta) = \sin \theta \\ \operatorname{cosec}(3\pi/2 + \theta) = -\sec \theta & \sec(3\pi/2 + \theta) = \operatorname{cosec} \theta \\ \tan(3\pi/2 + \theta) = -\cot \theta & \cot(3\pi/2 + \theta) = -\tan \theta\end{array}$$

$$\begin{array}{ll}\sin(2\pi - \theta) = -\sin \theta & \cos(2\pi - \theta) = \cos \theta \\ \operatorname{cosec}(2\pi - \theta) = -\operatorname{cosec} \theta & \sec(2\pi - \theta) = \sec \theta \\ \tan(2\pi - \theta) = -\tan \theta & \cot(2\pi - \theta) = -\cot \theta\end{array}$$

- *The values for $(2\pi - \theta)$ and $(-\theta)$ are identical.*

7.3 Compound Angle Formulae

$$\sin(A + B) = \sin(A) \cos(B) + \cos(A) \sin(B)$$

$$\sin(A - B) = \sin(A) \cos(B) - \cos(A) \sin(B)$$

$$\cos(A + B) = \cos(A) \cos(B) - \sin(A) \sin(B)$$

$$\cos(A - B) = \cos(A) \cos(B) + \sin(A) \sin(B)$$

$$\tan(A + B) = \frac{\tan(A) + \tan(B)}{1 - \tan(A) \tan(B)}$$

$$\tan(A - B) = \frac{\tan(A) - \tan(B)}{1 + \tan(A) \tan(B)}$$

7.4 Sum or Difference Formulae

$$\sin(A + B) + \sin(A - B) = 2 \sin(A) \cos(B)$$

$$\sin(A - B) - \sin(A - B) = 2 \cos(A) \sin(B)$$

$$\cos(A + B) + \cos(A - B) = 2 \cos(A) \cos(B)$$

$$\cos(A + B) - \cos(A - B) = -2 \sin(A) \sin(B)$$

7.5 Half Angle Formulae

$$\sin\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 - \cos(A)}{2}}$$

$$\cos\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 + \cos(A)}{2}}$$

$$\tan\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 - \cos(A)}{1 + \cos(A)}}$$

7.6 Sub Multiple Angle Formulae

$$\sin(2A) = 2 \sin(A) \cos(A)$$

$$\cos(2A) = \cos^2(A) - \sin^2(A) = 2 \cos^2(A) - 1 = 1 - 2 \sin^2(A)$$

$$\tan(2A) = \frac{2 \tan(A)}{1 - \tan^2(A)}$$

$$\sin(2A) = \frac{2 \tan(A)}{1 + \tan^2(A)}$$

$$\cos(2A) = \frac{1 - \tan^2(A)}{1 + \tan^2(A)}$$

$$\cos^2(A) = \frac{1 + \cos(2A)}{2}$$

$$\sin^2(A) = \frac{1 - \cos(2A)}{2}$$

$$\sin(A) = 2 \sin(A/2) \cos(A/2)$$

$$\cos(A) = \cos^2(A/2) - \sin^2(A/2) = 2 \cos^2(A/2) - 1 = 1 - 2 \sin^2(A/2)$$

7.7 Triple Angle Formulae

$$\sin(3x) = 3 \sin(x) - 4 \sin^3(x)$$

$$\cos(3x) = 4 \cos^3(x) - 3 \cos(x)$$

$$\tan(3x) = \frac{3 \tan(x) - \tan^3(x)}{1 - 3 \tan^2(x)}$$

$$\sin(4x) = 4 \sin(x) \cos(x) - 8 \sin^3(x) \cos(x)$$

$$\tan(4x) = \frac{4 \tan(x) - 4 \tan^3(x)}{1 - 6 \tan^2(x) + \tan^4(x)}$$

$$\cos(4x) = 8 \cos^4(x) - 8 \cos^2(x) + 1$$

7.8 Product Formulae

If $A + B = C$ and $A - B = D$

Then $A = (C + D)/2$ and $B = (C - D)/2$, the product formulae are defined as

$$\sin(C) + \sin(D) = 2 \sin \frac{(C + D)}{2} \cos \frac{(C - D)}{2}$$

$$\sin(C) - \sin(D) = 2 \cos \frac{(C + D)}{2} \sin \frac{(C - D)}{2}$$

$$\cos(C) + \cos(D) = 2 \cos \frac{(C + D)}{2} \cos \frac{(C - D)}{2}$$

$$\cos(C) - \cos(D) = -2 \sin \frac{(C + D)}{2} \sin \frac{(C - D)}{2}$$

8 Inverse Trigonometric Functions

If $x = \sin \theta$ then we define the inverse trigonometric function as $\sin^{-1}(x) = \theta$, the same holds true for all other trigonometric functions.

8.1 Properties of Inverse Trigonometric Functions

$$\begin{array}{ll} \sin^{-1}(\sin \theta) = \theta & \operatorname{cosec}^{-1}(\operatorname{cosec} \theta) = \theta \\ \cos^{-1}(\cos \theta) = \theta & \sec^{-1}(\sec \theta) = \theta \\ \tan^{-1}(\tan \theta) = \theta & \cot^{-1}(\cot \theta) = \theta \end{array}$$

$$\begin{array}{ll} \sin(\sin^{-1}(x)) = x & \operatorname{cosec}(\operatorname{cosec}^{-1}(x)) = x \\ \cos(\cos^{-1}(x)) = x & \sec(\sec^{-1}(x)) = x \\ \tan(\tan^{-1}(x)) = x & \cot(\cot^{-1}(x)) = x \end{array}$$

Functions	Domain	Range
$\sin^{-1}x$	$[-1, 1]$	$[-\pi/2, \pi/2]$
$\cos^{-1}x$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1}x$	\mathbb{R}	$[-\pi/2, \pi/2]$
$\cot^{-1}x$	\mathbb{R}	$[0, \pi]$
$\sec^{-1}x$	$\mathbb{R} - (-1, 1)$	$[0, \pi] - [\pi/2]$
$\operatorname{cosec}^{-1}x$	$\mathbb{R} - (-1, 1)$	$[-\pi/2, \pi/2] - \{0\}$

Figure 2: Domain and range of inverse trigonometric functions

$$\begin{array}{lll} \sin^{-1}(x) & = & \cos^{-1} \sqrt{1-x^2} = \tan^{-1} \frac{x}{\sqrt{1-x^2}} \\ \cos^{-1}(x) & = & \sin^{-1} \sqrt{1-x^2} = \tan^{-1} \frac{\sqrt{1-x^2}}{x} \\ \tan^{-1}(x) & = & \sin^{-1} \frac{x}{\sqrt{1+x^2}} = \cos^{-1} \frac{1}{\sqrt{1+x^2}} \end{array}$$

$$\begin{aligned}
\sin^{-1}(-x) &= -\sin^{-1}(x) & x \in [-1, 1] \\
\cos^{-1}(-x) &= \pi - \cos^{-1}(x) & x \in [-1, 1] \\
\tan^{-1}(-x) &= -\tan^{-1}(x) & x \in \mathbb{R} \\
\operatorname{cosec}^{-1}(-x) &= -\operatorname{cosec}^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\sec^{-1}(-x) &= \pi - \sec^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\cot^{-1}(-x) &= \pi - \cot^{-1}(x) & x \in \mathbb{R}
\end{aligned}$$

$$\begin{aligned}
\sin^{-1}\left(\frac{1}{x}\right) &= \operatorname{cosec}^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\cos^{-1}\left(\frac{1}{x}\right) &= \sec^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\tan^{-1}(1/x) &= \begin{cases} \cot^{-1}(x), & x > 0 \\ \cot^{-1}(x) - \pi, & x < 0 \end{cases}
\end{aligned}$$

$$\begin{aligned}
\sin^{-1}(x) + \cos^{-1}(x) &= (\pi/2) \\
\tan^{-1}(x) + \cot^{-1}(x) &= (\pi/2) \\
\sec^{-1}(x) + \operatorname{cosec}^{-1}(x) &= (\pi/2)
\end{aligned}$$

$$\begin{aligned}
\tan^{-1}(x) + \tan^{-1}(y) &= \tan^{-1} \frac{x+y}{1-xy} \\
\tan^{-1}(x) - \tan^{-1}(y) &= \tan^{-1} \frac{x-y}{1+xy} \\
2 \tan^{-1}(x) &= \begin{cases} 2 \sin^{-1} \frac{2x}{1+x^2}, & x > 0 \\ \cos^{-1} \frac{1-x^2}{1+x^2}, & x < 0 \end{cases}
\end{aligned}$$

8.2 Expressions and suggested substitutions

Expression	Substitution
$a^2 + x^2$	$x = a \tan \theta \quad \text{or} \quad x = a \cot \theta$
$a^2 - x^2$	$x = a \sin \theta \quad \text{or} \quad x = a \cos \theta$
$x^2 - a^2$	$x = a \operatorname{cosec} \theta \quad \text{or} \quad x = a \sec \theta$
$\sqrt{\frac{a-x}{a+x}} \quad \text{or} \quad \sqrt{\frac{a+x}{a-x}}$	$x = a \cos 2\theta$
$\sqrt{\frac{a^2-x^2}{a^2+x^2}} \quad \text{or} \quad \sqrt{\frac{a^2+x^2}{a^2-x^2}}$	$x = a^2 \cos 2\theta$

9 Hyperbolic Functions

$$\begin{aligned}\sinh(x) &= \frac{e^x - e^{-x}}{2} & \operatorname{cosech}(x) &= \frac{2}{e^x - e^{-x}} \\ \cosh(x) &= \frac{e^x + e^{-x}}{2} & \operatorname{sech}(x) &= \frac{2}{e^x + e^{-x}} \\ \tanh(x) &= \frac{e^x - e^{-x}}{e^x + e^{-x}} & \operatorname{coth}(x) &= \frac{e^x + e^{-x}}{e^x - e^{-x}}\end{aligned}$$

9.1 Properties of Hyperbolic Functions

$$\begin{aligned}\cosh^2(x) - \sinh^2(x) &= 1 \\ \cosh^2(x) + \sinh^2(x) &= \cosh(2x) \\ 1 - \tanh^2(x) &= \operatorname{sech}^2(x) \\ \operatorname{coth}^2(x) - 1 &= \operatorname{cosech}^2(x) \\ \sinh(2x) &= 2 \sinh(x) \cosh(x) \\ \sin(ix) &= \frac{e^{ix} - e^{-ix}}{2} \\ \cos(ix) &= \frac{e^{ix} + e^{-ix}}{2} \\ \sin(ix) &= i \sinh(x) \\ \cos(ix) &= \cosh(x) \\ \tan(ix) &= i \tanh(x)\end{aligned}$$