

Basics of Engineering Mathematics

Santosh S Malagi

Technische Universiteit Delft

Email: s.s.malagi@student.tudelft.nl

August 2, 2017




This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/> or send a letter to Creative Commons, PO Box 1866, Mountain View, CA 94042, USA.

Contents

1	Basic Algebra	4
1.1	Law of Negatives	4
1.2	Law of Quotients	4
2	Polynomial Functions	5
2.1	Corollary to the Fundamental Theorem	5
2.2	Synthetic Division	6
3	Quadratic Equations	7
3.1	Factorization method	7
3.2	Completion of squares method	7
4	Logarithmic Functions	8
4.1	Properties of Logarithmic Functions	8
5	Laws of Exponents	9
5.1	Rational Number Exponents	10
5.2	Simplest Radical Form	10
6	Complex Numbers	11
6.1	De Moivre's Theorem	11
7	Permutations and Combinations	12
7.1	Binomial Theorem	12
7.2	Summation of Series	12
8	Trigonometry	13
8.1	Trigonometric Identities	13
8.2	Trigonometric ratios of allied angles	13
8.2.1	First Quadrant	14
8.2.2	Second Quadrant	14
8.2.3	Third Quadrant	14
8.2.4	Fourth Quadrant	14
8.3	Compound Angle Formulae	15
8.4	Sum or Difference Formulae	15
8.5	Half Angle Formulae	15
8.6	Sub Multiple Angle Formulae	15
8.7	Triple Angle Formulae	16
8.8	Product Formulae	16
9	Inverse Trigonometric Functions	17
9.1	Properties of Inverse Trigonometric Functions	17
9.2	Expressions and suggested substitutions	19
10	Hyperbolic Functions	20
10.1	Properties of Hyperbolic Functions	20
11	Linear Inequalities	21

12 Limits and Differentiation	22
12.1 Fundamental principles of limits	22
12.2 Properties	22
12.3 Standard Limits	22
12.4 Dealing with indeterminate forms	23
12.5 L'Hospital's Rule	23
12.6 Differentiation Formulae	23
12.7 Differentiation of Trigonometric Functions	24
12.8 Differentiation of Inverse Trigonometric Functions	24
12.9 Differentiation of Hyperbolic Functions	24
12.10 Product Rule	25
12.11 Quotient Rule	25
12.12 Derivative of Inverse of a function	25
12.13 Chain Rule of Differentiation (Derivative of composite function)	25



*“Dedicated to the
mathematicians of
India, who gifted the
world an abundance of
knowledge and a spirit
of enquiry”.*



1 Basic Algebra

$$(a + b)^2 = a^2 + 2ab + b^2 \quad (1)$$

$$(a - b)^2 = a^2 - 2ab + b^2 \quad (2)$$

$$(a + b)(a - b) = a^2 - b^2 \quad (3)$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3 = a^3 + b^3 + 3ab(a + b) \quad (4)$$

$$(a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3 = a^3 - b^3 - 3ab(a - b) \quad (5)$$

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2) \quad (6)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2) \quad (7)$$

$$a^4 - b^4 = (a + b)(a - b)(a^2 + b^2) \quad (8)$$

$$a^2 + b^2 = (a + b)^2 - 2ab \quad (9)$$

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac) \quad (10)$$

1.1 Law of Negatives

$$\begin{aligned} -(-a) &= a \\ (-a)(-b) &= ab \\ -ab &= (-a)(b) = a(-b) = -(-a)(-b) \\ a - b &= a + (-b) \end{aligned}$$

1.2 Law of Quotients

$$\begin{aligned} -\frac{a}{b} &= \frac{(-a)}{b} = \frac{a}{(-b)} = -\frac{(-a)}{(-b)} \\ \frac{a}{b} &= \frac{c}{d} \quad \text{if} \quad ad = bc \\ \frac{a}{b} &= \frac{ka}{kb} \end{aligned}$$

2 Polynomial Functions

Zeros of a polynomial: If $f(c) = 0$, then c is called a zero of the polynomial $f(x)$.

Division algorithm for a polynomial: If $f(x)$ and $g(x)$ are two polynomials such that $g(x) \neq 0$ then there exist two unique polynomials $q(x)$ and $r(x)$ such that:

$$\frac{f(x)}{g(x)} = q(x) + \frac{r(x)}{g(x)} \quad (11)$$

Fundamental Theorem of Algebra: Every polynomial with positive degree has at least one complex zero.

Remainder Theorem: If a polynomial $f(x)$ is divided by $x - c$ then the remainder is $f(c)$.

Factor Theorem: Polynomial $f(x)$ has a factor $x - c$ if and only if $f(c) = 0$.

2.1 Corollary to the Fundamental Theorem

1. Every polynomial of positive degree n has a factorization of the form:

$$P(x) = a_n(x - r_1)(x - r_2)\dots\dots(x - r_n) \quad (12)$$

where r_i are not necessarily distinct. If $(x - r_i)$ occurs n times it is said to have a multiplicity of n .

2. It is not always possible to find the factors using exact methods.
3. A polynomial of degree n has at most n complex zeros.
4. Complex zeros of real polynomials with real coefficients occur in complex conjugate pairs.
5. Any polynomial of degree $n > 0$ with real coefficients has complete factorization using linear and quadratic factors, multiplied by the leading coefficient of the polynomial.
6. **Intermediate Value Theorem:** Given a polynomial $f(x)$ with $a < b$ and $f(a) \neq f(b)$ then $f(c)$ takes on every value c in the interval (a, b) .
7. For a polynomial $f(x)$ if $f(a)$ and $f(b)$ have opposite signs then $f(x)$ has at least one zero between a and b .
8. For a polynomial $f(x)$ if $f(a)$ and $f(b)$ have opposite signs then $f(x)$ has at least one zero between a and b .

9. **Descartes rule of signs:** If $f(x)$ is a polynomial with terms arranged in descending order, then the number of positive real roots of $f(x)$ is either equal to the number of sign changes between the successive terms of $f(x)$ or less than this number by an even number. Number of negative real zeros of $f(x)$ is obtained by applying this rule to $f(-x)$.

2.2 Synthetic Division

c is said to be a zero of the polynomial $f(x)$ if $f(c) = 0 \implies x - c$ is a factor of the polynomial $f(x)$. The graph of $f(x)$ has an intercept at c .

1. Arrange the coefficients in descending order in the first row.
2. Third row is formed by bringing down the first coefficient of $f(x)$ then successively multiplying each coefficient in the third row by c , placing the results in second row adding this to the corresponding coefficients in the first row, and placing result in the next position of the third row.

3 Quadratic Equations

The quadratic equation

$$ax^2 + bx + c = 0 \quad (13)$$

where a, b and c are constants and $a \neq 0$, has two solutions for the variable x :

$$x_{\alpha, \beta} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (14)$$

If the *discriminant* Δ with

$$\Delta = b^2 - 4ac = 0 \quad (15)$$

then the equation (13) has real and equal roots given by:

$$x = \frac{-b}{2a} \quad (16)$$

Sum of the roots:

$$\alpha + \beta = \frac{-b}{a} \quad (17)$$

Product of the roots:

$$\alpha \cdot \beta = \frac{c}{a} \quad (18)$$

3.1 Factorization method

Find the two factors $x - \alpha$ and $x - \beta$ which when multiplied produce the quadratic equation $ax^2 + bx + c$.

3.2 Completion of squares method

Consider any quadratic equation of the form $ax^2 + bx + c = 0$. Dividing throughout by a .

$$\implies x^2 + \left(\frac{b}{a}\right)x + \frac{c}{a} = 0$$

$$\text{Adding and subtracting } \left(\frac{b}{2a}\right)^2,$$

$$x^2 + \left(\frac{b}{a}\right)x + \left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{2a}\right)^2 + \frac{c}{a} - \left(\frac{b}{2a}\right)^2 = 0$$

$$\therefore ax^2 + bx + c = 0 \implies a(x + d)^2 + e = 0 \quad \text{where,} \quad d = \frac{b}{2a} \quad \text{and} \quad e = \frac{c}{a} - \left(\frac{b}{2a}\right)^2.$$

4 Logarithmic Functions

For any two variables x and y related as:

$$y = a^x \tag{19}$$

Where a is a constant. We define the logarithmic function as:

$$\log_a y = x \tag{20}$$

4.1 Properties of Logarithmic Functions

$$\log_a mn = \log_a m + \log_a n \tag{21}$$

$$\log_a \frac{m}{n} = \log_a m - \log_a n \tag{22}$$

$$\log_a m^n = n \cdot \log_a m \tag{23}$$

$$\log_a a = 1 \tag{24}$$

$$\log_a 1 = 0 \tag{25}$$

$$\log_a b = \frac{1}{\log_b a} \tag{26}$$

$$a^x = e^{x \cdot \log a} \tag{27}$$

$$\log_a X = \frac{\ln X}{\ln a} \tag{28}$$

5 Laws of Exponents

$$x^a . x^b = x^{a+b} \tag{29}$$

$$\frac{x^a}{x^b} = x^{a-b} \tag{30}$$

$$(x^a)^b = x^{ab} \tag{31}$$

$$x^a = y^b \implies a = b \tag{32}$$

$$x^{-a} = \frac{1}{x^a} \quad (\text{for } a \text{ being any non zero real number}) \tag{33}$$

$$x^0 = 1 \tag{34}$$

$$\sqrt[n]{\frac{x}{y}} = \frac{\sqrt[n]{x}}{\sqrt[n]{y}} \tag{35}$$

$$\sqrt[n]{xy} = \sqrt[n]{x} . \sqrt[n]{y} \tag{36}$$

$$\left(\frac{x}{y}\right)^{-m} = \left(\frac{y}{x}\right)^m \tag{37}$$

$$\frac{x^{-n}}{y^{-m}} = \frac{y^m}{x^n} \tag{38}$$

5.1 Rational Number Exponents

The principle n th root of $x > 0$ is defined as:

$$x^{1/n} = \begin{cases} \text{is a unique real number } y \text{ if } n \text{ is odd or even} \\ 0, \text{ if } x = 0 \\ \text{not a real number if } x < 0 \end{cases}$$

$$(x^{1/n})^n = \begin{cases} x, \text{ if } n \text{ is odd/even and } x \text{ is positive} \\ |x|, \text{ if } n \text{ is even and } x \text{ is negative} \end{cases}$$

5.2 Simplest Radical Form

No radicand can contain a factor with an exponent greater than or equal to the index of the radical e.g. $\sqrt[3]{16x^3y^5} \implies$ is not in it's standard form. No power of a radicand and index of the radical can have common factor other than 1. e.g. $2xy\sqrt[3]{2y^2} \implies$ not in standard form. No radical appears in denominator. No fractional part appears in radical. e.g. $\sqrt[6]{t^3} = \sqrt{t}$ is in it's standard form.

6 Complex Numbers

A number of the form $z = x + iy$ consisting of a real part x , denoted by $Re(z)$ and imaginary part y , denoted by $Im(z)$ is known as a complex number.

$$x = r \cos \theta \tag{39}$$

$$y = r \sin \theta \tag{40}$$

$$r = \sqrt{x^2 + y^2} \tag{41}$$

$$\tan \theta = \frac{y}{x} \implies \theta = \tan^{-1} \frac{y}{x} \tag{42}$$

$$e^{i\theta} = \cos \theta + i \sin \theta \tag{43}$$

$$e^{-i\theta} = \cos \theta - i \sin \theta \tag{44}$$

6.1 De Moivre's Theorem

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta) \tag{45}$$

7 Permutations and Combinations

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2)(n-3)\dots(n-(r+1))}{r!} \quad (46)$$

Note that:

$$\binom{n}{0} = 1 \qquad \binom{n}{1} = n \qquad 0! = 1$$

7.1 Binomial Theorem

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \binom{n}{2}x^{n-2}a^2 + \binom{n}{3}x^{n-3}a^3 + \binom{n}{4}x^{n-4}a^4 + \dots\dots\dots a^n \quad (47)$$

7.2 Summation of Series

$$\sum_{n=1}^{\infty} n = 1 + 2 + 3\dots\dots = \frac{(n)(n+1)}{2} \quad (48)$$

$$\sum_{n=1}^{\infty} n^2 = 1^2 + 2^2 + 3^2\dots\dots = \frac{(n)(n+1)(2n+1)}{6} \quad (49)$$

$$\sum_{n=1}^{\infty} n^3 = 1^3 + 2^3 + 3^3\dots\dots = \frac{(n)^2(n+1)^2}{4} \quad (50)$$

8 Trigonometry

$$\begin{aligned}\sin \theta &= \frac{\text{Perpendicular}}{\text{Hypotenuse}} & \cos \theta &= \frac{\text{Base}}{\text{Hypotenuse}} & \tan \theta &= \frac{\text{Perpendicular}}{\text{Base}} \\ \operatorname{cosec} \theta &= \frac{\text{Hypotenuse}}{\text{Perpendicular}} & \sec \theta &= \frac{\text{Hypotenuse}}{\text{Base}} & \cot \theta &= \frac{\text{Base}}{\text{Perpendicular}}\end{aligned}$$

From the above relations it turns out that,

$$\sin \theta \cdot \operatorname{cosec} \theta = 1$$

$$\cos \theta \cdot \sec \theta = 1$$

$$\tan \theta \cdot \cot \theta = 1$$

8.1 Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \tag{51}$$

$$1 + \sec^2 \theta = \tan^2 \theta \tag{52}$$

$$1 + \operatorname{cosec}^2 \theta = \cot^2 \theta \tag{53}$$

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined
$\cot \theta$	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0
$\sec \theta$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
$\operatorname{cosec} \theta$	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1

Figure 1: Most common trigonometric ratios and values

8.2 Trigonometric ratios of allied angles

- *For n being an odd multiple of $\pi/2$ the function changes to its composite function.*
- *For n being an even multiple of $\pi/2$ the function does not change to its composite, but may change sign.*

8.2.1 First Quadrant

$$\begin{array}{ll}\sin(2\pi + \theta) = \sin \theta & \cos(2\pi + \theta) = \cos \theta \\ \operatorname{cosec}(2\pi + \theta) = \operatorname{cosec} \theta & \sec(2\pi + \theta) = \sec \theta \\ \tan(2\pi + \theta) = \tan \theta & \cot(2\pi + \theta) = \cot \theta\end{array}$$

$$\begin{array}{ll}\sin(\pi/2 - \theta) = \cos \theta & \cos(\pi/2 - \theta) = \sin \theta \\ \operatorname{cosec}(\pi/2 - \theta) = \sec \theta & \sec(\pi/2 - \theta) = \operatorname{cosec} \theta \\ \tan(\pi/2 - \theta) = \cot \theta & \cot(\pi/2 - \theta) = \tan \theta\end{array}$$

8.2.2 Second Quadrant

$$\begin{array}{ll}\sin(\pi/2 + \theta) = \cos \theta & \cos(\pi/2 + \theta) = -\sin \theta \\ \operatorname{cosec}(\pi/2 + \theta) = \sec \theta & \sec(\pi/2 + \theta) = -\operatorname{cosec} \theta \\ \tan(\pi/2 + \theta) = -\cot \theta & \cot(\pi/2 + \theta) = -\tan \theta\end{array}$$

$$\begin{array}{ll}\sin(\pi - \theta) = \sin \theta & \cos(\pi - \theta) = -\cos \theta \\ \operatorname{cosec}(\pi - \theta) = \operatorname{cosec} \theta & \sec(\pi - \theta) = -\sec \theta \\ \tan(\pi - \theta) = -\tan \theta & \cot(\pi - \theta) = -\cot \theta\end{array}$$

8.2.3 Third Quadrant

$$\begin{array}{ll}\sin(\pi + \theta) = -\sin \theta & \cos(\pi + \theta) = -\cos \theta \\ \operatorname{cosec}(\pi + \theta) = \operatorname{cosec} \theta & \sec(\pi + \theta) = -\sec \theta \\ \tan(\pi + \theta) = \tan \theta & \cot(\pi + \theta) = \cot \theta\end{array}$$

$$\begin{array}{ll}\sin(3\pi/2 - \theta) = -\cos \theta & \cos(3\pi/2 - \theta) = -\sin \theta \\ \operatorname{cosec}(3\pi/2 - \theta) = -\sec \theta & \sec(3\pi/2 - \theta) = -\operatorname{cosec} \theta \\ \tan(3\pi/2 - \theta) = \cot \theta & \cot(3\pi/2 - \theta) = \tan \theta\end{array}$$

8.2.4 Fourth Quadrant

$$\begin{array}{ll}\sin(3\pi/2 + \theta) = -\cos \theta & \cos(3\pi/2 + \theta) = \sin \theta \\ \operatorname{cosec}(3\pi/2 + \theta) = -\sec \theta & \sec(3\pi/2 + \theta) = \operatorname{cosec} \theta \\ \tan(3\pi/2 + \theta) = -\cot \theta & \cot(3\pi/2 + \theta) = -\tan \theta\end{array}$$

$$\begin{array}{ll}\sin(2\pi - \theta) = -\sin \theta & \cos(2\pi - \theta) = \cos \theta \\ \operatorname{cosec}(2\pi - \theta) = -\operatorname{cosec} \theta & \sec(2\pi - \theta) = \sec \theta \\ \tan(2\pi - \theta) = -\tan \theta & \cot(2\pi - \theta) = -\cot \theta\end{array}$$

- *The values for $(2\pi - \theta)$ and $(-\theta)$ are identical.*

8.3 Compound Angle Formulae

$$\sin(A + B) = \sin(A) \cos(B) + \cos(A) \sin(B)$$

$$\sin(A - B) = \sin(A) \cos(B) - \cos(A) \sin(B)$$

$$\cos(A + B) = \cos(A) \cos(B) - \sin(A) \sin(B)$$

$$\cos(A - B) = \cos(A) \cos(B) + \sin(A) \sin(B)$$

$$\tan(A + B) = \frac{\tan(A) + \tan(B)}{1 - \tan(A) \tan(B)}$$

$$\tan(A - B) = \frac{\tan(A) - \tan(B)}{1 + \tan(A) \tan(B)}$$

8.4 Sum or Difference Formulae

$$\sin(A + B) + \sin(A - B) = 2 \sin(A) \cos(B)$$

$$\sin(A - B) - \sin(A - B) = 2 \cos(A) \sin(B)$$

$$\cos(A + B) + \cos(A - B) = 2 \cos(A) \cos(B)$$

$$\cos(A + B) - \cos(A - B) = -2 \sin(A) \sin(B)$$

8.5 Half Angle Formulae

$$\sin\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 - \cos(A)}{2}}$$

$$\cos\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 + \cos(A)}{2}}$$

$$\tan\left(\frac{A}{2}\right) = \pm \sqrt{\frac{1 - \cos(A)}{1 + \cos(A)}}$$

8.6 Sub Multiple Angle Formulae

$$\sin(2A) = 2 \sin(A) \cos(A)$$

$$\cos(2A) = \cos^2(A) - \sin^2(A) = 2 \cos^2(A) - 1 = 1 - 2 \sin^2(A)$$

$$\tan(2A) = \frac{2 \tan(A)}{1 - \tan^2(A)}$$

$$\sin(2A) = \frac{2 \tan(A)}{1 + \tan^2(A)}$$

$$\cos(2A) = \frac{1 - \tan^2(A)}{1 + \tan^2(A)}$$

$$\cos^2(A) = \frac{1 + \cos(2A)}{2}$$

$$\sin^2(A) = \frac{1 - \cos(2A)}{2}$$

$$\sin(A) = 2 \sin(A/2) \cos(A/2)$$

$$\cos(A) = \cos^2(A/2) - \sin^2(A/2) = 2 \cos^2(A/2) - 1 = 1 - 2 \sin^2(A/2)$$

8.7 Triple Angle Formulae

$$\sin(3x) = 3\sin(x) - 4\sin^3(x)$$

$$\cos(3x) = 4\cos^3(x) - 3\cos(x)$$

$$\tan(3x) = \frac{3\tan(x) - \tan^3(x)}{1 - 3\tan^2(x)}$$

$$\sin(4x) = 4\sin(x)\cos(x) - 8\sin^3(x)\cos(x)$$

$$\tan(4x) = \frac{4\tan(x) - 4\tan^3(x)}{1 - 6\tan^2(x) + \tan^4(x)}$$

$$\cos(4x) = 8\cos^4(x) - 8\cos^2(x) + 1$$

8.8 Product Formulae

If $A + B = C$ and $A - B = D$

Then $A = (C + D)/2$ and $B = (C - D)/2$, the product formulae are defined as

$$\sin(C) + \sin(D) = 2\sin\frac{(C + D)}{2}\cos\frac{(C - D)}{2}$$

$$\sin(C) - \sin(D) = 2\cos\frac{(C + D)}{2}\sin\frac{(C - D)}{2}$$

$$\cos(C) + \cos(D) = 2\cos\frac{(C + D)}{2}\cos\frac{(C - D)}{2}$$

$$\cos(C) - \cos(D) = -2\sin\frac{(C + D)}{2}\sin\frac{(C - D)}{2}$$

9 Inverse Trigonometric Functions

If $x = \sin \theta$ then we define the inverse trigonometric function as $\sin^{-1}(x) = \theta$, the same holds true for all other trigonometric functions.

9.1 Properties of Inverse Trigonometric Functions

$$\begin{array}{ll} \sin^{-1}(\sin \theta) = \theta & \operatorname{cosec}^{-1}(\operatorname{cosec} \theta) = \theta \\ \cos^{-1}(\cos \theta) = \theta & \sec^{-1}(\sec \theta) = \theta \\ \tan^{-1}(\tan \theta) = \theta & \cot^{-1}(\cot \theta) = \theta \end{array}$$

$$\begin{array}{ll} \sin(\sin^{-1}(x)) = x & \operatorname{cosec}(\operatorname{cosec}^{-1}(x)) = x \\ \cos(\cos^{-1}(x)) = x & \sec(\sec^{-1}(x)) = x \\ \tan(\tan^{-1}(x)) = x & \cot(\cot^{-1}(x)) = x \end{array}$$

Functions	Domain	Range
$\sin^{-1}x$	$[-1, 1]$	$[-\pi/2, \pi/2]$
$\cos^{-1}x$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1}x$	\mathbb{R}	$[-\pi/2, \pi/2]$
$\cot^{-1}x$	\mathbb{R}	$[0, \pi]$
$\sec^{-1}x$	$\mathbb{R} - (-1, 1)$	$[0, \pi] - [\pi/2]$
$\operatorname{cosec}^{-1}x$	$\mathbb{R} - (-1, 1)$	$[-\pi/2, \pi/2] - \{0\}$

Figure 2: Domain and range of inverse trigonometric functions

$$\begin{array}{llll} \sin^{-1}(x) & = & \cos^{-1} \sqrt{1-x^2} & = & \tan^{-1} \frac{x}{\sqrt{1-x^2}} \\ \cos^{-1}(x) & = & \sin^{-1} \sqrt{1-x^2} & = & \tan^{-1} \frac{\sqrt{1-x^2}}{x} \\ \tan^{-1}(x) & = & \sin^{-1} \frac{x}{\sqrt{1+x^2}} & = & \cos^{-1} \frac{1}{\sqrt{1+x^2}} \end{array}$$

$$\begin{aligned}
\sin^{-1}(-x) &= -\sin^{-1}(x) & x \in [-1, 1] \\
\cos^{-1}(-x) &= \pi - \cos^{-1}(x) & x \in [-1, 1] \\
\tan^{-1}(-x) &= -\tan^{-1}(x) & x \in \mathbb{R} \\
\operatorname{cosec}^{-1}(-x) &= -\operatorname{cosec}^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\sec^{-1}(-x) &= \pi - \sec^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\cot^{-1}(-x) &= \pi - \cot^{-1}(x) & x \in \mathbb{R}
\end{aligned}$$

$$\begin{aligned}
\sin^{-1}\left(\frac{1}{x}\right) &= \operatorname{cosec}^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty) \\
\cos^{-1}\left(\frac{1}{x}\right) &= \sec^{-1}(x) & x \in (-\infty, -1] \cup [1, \infty)
\end{aligned}$$

$$\tan^{-1}(1/x) = \begin{cases} \cot^{-1}(x), & x > 0 \\ \cot^{-1}(x) - \pi, & x < 0 \end{cases}$$

$$\begin{aligned}
\sin^{-1}(x) + \cos^{-1}(x) &= (\pi/2) \\
\tan^{-1}(x) + \cot^{-1}(x) &= (\pi/2) \\
\sec^{-1}(x) + \operatorname{cosec}^{-1}(x) &= (\pi/2)
\end{aligned}$$

$$\begin{aligned}
\tan^{-1}(x) + \tan^{-1}(y) &= \tan^{-1} \frac{x+y}{1-xy} \\
\tan^{-1}(x) - \tan^{-1}(y) &= \tan^{-1} \frac{x-y}{1+xy}
\end{aligned}$$

$$2 \tan^{-1}(x) = \begin{cases} 2 \sin^{-1} \frac{2x}{1+x^2}, & x > 0 \\ \cos^{-1} \frac{1-x^2}{1+x^2}, & x < 0 \end{cases}$$

9.2 Expressions and suggested substitutions

Expression	Substitution
$a^2 + x^2$	$x = a \tan \theta \quad \text{or} \quad x = a \cot \theta$
$a^2 - x^2$	$x = a \sin \theta \quad \text{or} \quad x = a \cos \theta$
$x^2 - a^2$	$x = a \operatorname{cosec} \theta \quad \text{or} \quad x = a \sec \theta$
$\sqrt{\frac{a-x}{a+x}} \quad \text{or} \quad \sqrt{\frac{a+x}{a-x}}$	$x = a \cos 2\theta$
$\sqrt{\frac{a^2-x^2}{a^2+x^2}} \quad \text{or} \quad \sqrt{\frac{a^2+x^2}{a^2-x^2}}$	$x = a^2 \cos 2\theta$

10 Hyperbolic Functions

$$\begin{aligned}\sinh(x) &= \frac{e^x - e^{-x}}{2} & \operatorname{cosech}(x) &= \frac{2}{e^x - e^{-x}} \\ \cosh(x) &= \frac{e^x + e^{-x}}{2} & \operatorname{sech}(x) &= \frac{2}{e^x + e^{-x}} \\ \tanh(x) &= \frac{e^x - e^{-x}}{e^x + e^{-x}} & \operatorname{coth}(x) &= \frac{e^x + e^{-x}}{e^x - e^{-x}}\end{aligned}$$

10.1 Properties of Hyperbolic Functions

$$\begin{aligned}\cosh^2(x) - \sinh^2(x) &= 1 \\ \cosh^2(x) + \sinh^2(x) &= \cosh(2x) \\ 1 - \tanh^2(x) &= \operatorname{sech}^2(x) \\ \operatorname{coth}^2(x) - 1 &= \operatorname{cosech}^2(x) \\ \sinh(2x) &= 2 \sinh(x) \cosh(x) \\ \sin(ix) &= \frac{e^{ix} - e^{-ix}}{2} \\ \cos(ix) &= \frac{e^{ix} + e^{-ix}}{2} \\ \sin(ix) &= i \sinh(x) \\ \cos(ix) &= \cosh(x) \\ \tan(ix) &= i \tanh(x)\end{aligned}$$

11 Linear Inequalities

Inequalities of the form:

$$ax + b < 0$$

$$ax + b > 0$$

$$ax + b \leq 0$$

$$ax + b \geq 0$$

are known as linear inequalities.

1. Add and subtract any number k without change in the inequality on both sides.
2. When multiplying or dividing by constant k , reverse the sign of the inequality only when k is negative.
3. Linear inequalities have infinite solution sets, these can be obtained by isolating the variable and solving them in a manner similar to solving equations.

Inequality	Representation	Graph
$a < x < b$	(a, b)	
$a \leq x \leq b$	$[a, b]$	
$a < x \leq b$	$(a, b]$	
$a \leq x < b$	$[a, b)$	
$x > a$	(a, ∞)	
$x \geq a$	$[a, \infty)$	
$x < b$	$(-\infty, b)$	
$x \leq b$	$(-\infty, b]$	

12 Limits and Differentiation

12.1 Fundamental principles of limits

1. The limit may exist at a point even if the function is undefined.
2. If a function $f(x)$ is defined at a point a i.e. $f(a)$ exists, it is not necessary that the limit at a must exist. Moreover, even if the limit exists it need not be equal to $f(a)$.
3. **Indeterminate forms:** Any function assuming either of these forms is said to be indeterminate - $\frac{0}{0}, 0.\infty, \frac{\infty}{\infty}, \infty - \infty, 0^0, 1^\infty, \infty^0$.

12.2 Properties

$$\begin{aligned}\lim_{x \rightarrow a} [f(x) \pm g(x)] &= \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x). \\ \lim_{x \rightarrow a} [f(x).g(x)] &= \lim_{x \rightarrow a} f(x). \lim_{x \rightarrow a} g(x). \\ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} &= \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} \quad \text{provided } \lim_{x \rightarrow a} g(x) \neq 0.\end{aligned}$$

12.3 Standard Limits

$$\lim_{x \rightarrow 0} \sin x = 0 \quad (54)$$

$$\lim_{x \rightarrow 0} \tan x = 0 \quad (55)$$

$$\lim_{x \rightarrow 0} \cos x = 1 \quad (56)$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad (57)$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \quad (58)$$

$$\lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} = 1 \quad (59)$$

$$\lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} = 1 \quad (60)$$

$$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a \quad \text{where } a > 0 \quad (61)$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \quad (62)$$

$$\lim_{x \rightarrow 0} \frac{\log_a(1+x)}{x} = \log_a e \quad \text{where } a > 0 \text{ and } a \neq 1 \quad (63)$$

$$\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1 \quad (64)$$

$$\lim_{x \rightarrow 0} \frac{\log x}{x^m} = 0 \quad \text{where } m > 0 \quad (65)$$

$$\lim_{x \rightarrow 0} \frac{(1+x)^m - 1}{x} = m \quad (66)$$

$$\lim_{x \rightarrow 0} \frac{x^n - a^n}{x - a} = n.a^{n-1} \quad (67)$$

12.4 Dealing with indeterminate forms

If $f(x)$ and $g(x)$ are two functions such that $f(x) \rightarrow 0$ as $x \rightarrow a$ and $g(x) \rightarrow \infty$ as $x \rightarrow a$ such that $[1 + f(x)]^{g(x)}$ assumes the form 1^∞ then:

$$\lim_{x \rightarrow a} [1 + f(x)]^{g(x)} = e^{\lim_{x \rightarrow a} f(x) \cdot g(x)}$$

12.5 L'Hospital's Rule

L'Hospital's rule states that for functions $f(x)$ and $g(x)$ which are differentiable on an open interval I except possibly at a point c contained in I , if:

$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} g(x) = 0$ or $\pm\infty$ and $g'(x) \neq 0$ for all x in I with $x \neq c$ and $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ exists then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

12.6 Differentiation Formulae

$$\frac{d}{dx}[f(x) \pm g(x)] = \frac{df(x)}{dx} \pm \frac{dg(x)}{dx}$$

$$\frac{d}{dx} f(x) \cdot g(x) = \frac{d}{dx} f(x) \cdot \frac{d}{dx} g(x)$$

$$\frac{dc}{dx} = 0 \quad \text{where } c \text{ is a real constant}$$

$$\frac{dx^n}{dx} = nx^{n-1} \quad \text{where } n \text{ is any real number}$$

$$\frac{de^x}{dx} = e^x$$

$$\frac{da^x}{dx} = a^x \log_e a$$

$$\frac{d}{dx} \log_a x = \frac{1}{x \cdot \log a}$$

$$\frac{d}{dx} \ln x = \frac{1}{x}$$

12.7 Differentiation of Trigonometric Functions

$$\frac{d}{dx} \sin x = \cos x$$

$$\frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tan x = \sec^2 x$$

$$\frac{d}{dx} \operatorname{cosec} x = -\operatorname{cosec} x \cdot \cot x$$

$$\frac{d}{dx} \sec x = \sec x \cdot \tan x$$

$$\frac{d}{dx} \cot x = -\operatorname{cosec}^2 x$$

12.8 Differentiation of Inverse Trigonometric Functions

$$\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$$

$$\frac{d}{dx} \operatorname{cosec}^{-1} x = -\frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} \sec^{-1} x = \frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} \cot^{-1} x = -\frac{1}{1+x^2}$$

12.9 Differentiation of Hyperbolic Functions

$$\frac{d}{dx} \sinh x = \cosh x$$

$$\frac{d}{dx} \cosh x = \sinh x$$

$$\frac{d}{dx} \tanh x = \operatorname{sech}^2 x$$

$$\frac{d}{dx} \operatorname{cosech} x = -\operatorname{cosech} x \cdot \coth x$$

$$\frac{d}{dx} \operatorname{sech} x = -\operatorname{sech} x \cdot \tanh x$$

$$\frac{d}{dx} \coth x = -\operatorname{cosech}^2 x$$

12.10 Product Rule

$$\frac{d}{dx}.uv = u.\frac{dv}{dx} + v.\frac{du}{dx}$$

12.11 Quotient Rule

$$\frac{d}{dx} \cdot \frac{u}{v} = \frac{v \cdot \frac{du}{dx} - u \cdot \frac{dv}{dx}}{v^2}$$

12.12 Derivative of Inverse of a function

$$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$$

12.13 Chain Rule of Differentiation (Derivative of composite function)

If $u = \phi(x)$
 $y = \phi(u)$, then

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$