

## EE5605: Kernel Methods, Fall 2022 (56)

Indian Institute of Technology Hyderabad  
HW 1, 50 points. Assigned: Thursday 10.11.2022.  
**Due: Friday 18.11.2022 at 11:59 pm.**

1. Show that product of kernels is a kernel. (5)
2. Use one or more of the following results about kernels to answer the question that follows:
  - The sums of kernels are kernels.
  - The products of kernels are kernels.
  - If  $T : \mathcal{X} \rightarrow \mathcal{X}'$  is a mapping from the set  $\mathcal{X}$  to the set  $\mathcal{X}'$  and  $k$  is a kernel defined on  $\mathcal{X}'$ , then  $k(T(x), T(x'))$  is a kernel on  $\mathcal{X}$ .
  - Assume we can define a Taylor series  $f(x) = \sum_{n=0}^{\infty} a_n x^n, |x| < r, x \in \mathbb{R}$  for  $r \in (0, \infty]$ , with  $a_n \geq 0$  for all  $n \geq 0$ .

Define  $\mathcal{X}$  to be a  $\sqrt{r}$ -ball in  $\mathbb{R}^d$ . Then for  $x, x' \in \mathbb{R}^d$  such that  $\|x\| < \sqrt{r}, k(x, x') = f(\langle x, x' \rangle) = \sum_{n=0}^{\infty} a_n (\langle x, x' \rangle)^n$  is a kernel.

Show that  $k(x, y) := \exp(\langle x, y \rangle), x, y \in \mathbb{R}^d, d \geq 1$  is a valid kernel. (5)

3. Implement the vanilla (linear) and kernel versions of a support vector machine (SVM). Use python. Do not use built-in functions for SVM. You can use open-source solvers for the convex optimization. (20)
4. Use *Xsvm.csv*, *ysvm.csv* to train the linear classifier. Test on the following points  $x_t^1 = [1.9, 0.4], x_t^2 = [0.9, 0.9], x_t^3 = [1, 4, 1.5], x_t^4 = [0.01, 0.005]$ . Report your predictions. (10)
5. Demonstrate the ability of the kernel SVM to learn the non-linear XOR function. Experiment with different kernels. Report the result of the kernel SVM and compare it with the vanilla version. (10)