

Pushing the precision frontier at the LHC with V+jets

**Ulla Blumenschein,^d Shane Breeze,^b Chiara Debenedetti,^e Engin Eren,^g
Simona Gargiulo,^k Nigel Glover,^b Jonas Lindert,^b Daniel Maitre,^b Sarah A. Malik,^a
Bjoern Penning,^c Svenja K. Pflitsch,^g Darren Price,^f Stefan Richter,^l
Marek Schoenherr,^h Peter Schichter,^b Paolo Torrielli,^j Maria Ubiali,ⁱ
Nicholas Wardle,^a Angelo G. Zecchinelli^a**

^a*High Energy Physics Group, Blackett Laboratory, Imperial College, Prince Consort Road, London, SW7 2AZ, UK*

^b*Institute for Particle Physics Phenomenology, Durham University, Durham, DH1 3LE, UK*

^c*Bristol University, HH Wills Physics Laboratory, Tyndall Avenue, Bristol, BS8 1TL, UK*

^d*Queen Mary University of London, Mile End Road, London, E1 4NS, UK*

^e*University of California Santa Cruz, 1156 High Street, Santa Cruz, California, US*

^f*The University of Manchester, Oxford Rd, Manchester, M13 9PL, UK*

^g*Deutsches Elektronen-Synchrotron DESY, Notkestrae 85, 22607 Hamburg, Germany*

^h*University of Zurich, Rmistrasse 71, CH-8006, Zurich, Switzerland*

ⁱ*University of Cambridge, The Old Schools, Trinity Ln, Cambridge CB2 1TN, UK*

^j*Universit di Torino, Via Giuseppe Verdi, 8, 10124 Torino, Italy*

^k*Albert Ludwigs University of Freiburg, Fahnbergplatz, 79085 Freiburg im Breisgau, Germany*

^l*University College London, Gower St, Bloomsbury, London WC1E 6BT, UK*

ABSTRACT: This documents the proceedings from a workshop titled ‘Illuminating Standard candles at the LHC: V+jets’ held at Imperial College London on 25th-26th April 2017. It summarises the numerous contributions to the workshop, from the experimental overview of V+jets measurements at CMS and ATLAS and their role in searching for physics beyond the Standard Model to the status of higher order perturbative calculations to these processes and their inclusion in state of the art Monte Carlos. An executive summary of the ensuing discussions including a list of outcomes and wishlist for future consideration is also presented.

Contents

1	Introduction	1
2	Experimental overview of V+jet measurements	2
3	Summary of theoretical developments	5
3.1	Higher order QCD and EWK corrections	5
3.2	Status of generators	11
3.3	PDF constraints from V+jets	13
4	Backgrounds to BSM searches	13
5	Outcome/wishlist	14

1 Introduction

Processes in which a vector boson is produced in association with jet(s) in proton-proton collisions (V+jets) at the Large Hadron Collider (LHC) provide valuable benchmarks for precision tests of the Standard Model (SM), probing perturbative QCD and constraining Parton Distribution Functions (PDFs). They also contribute as dominant backgrounds to searches for a wide range of theoretical scenarios beyond the SM, such as searches for Supersymmetry, Dark Matter, and exotic decays of the Higgs boson to invisible particles.

The LHC is now in its second phase of operation (Run 2), colliding protons at the higher center of mass energy of 13 TeV and expecting to accumulate 100 fb^{-1} of data by the end of Run 2 in 2018, 5 times higher than previously studied in Run 1. As the LHC collects an unprecedented dataset and enters an era of precision, it presents a tremendous opportunity to perform ever more precise measurements of V+jets processes and do so in regions of phase space that were previously limited by statistics, such as the high transverse momentum region that is also characteristic of the phase space probed by our searches. In parallel, developments in theoretical calculations have led to improved predictions and state of the art Monte Carlos becoming available, with the near term prospect of having automated next-to-leading-order (NLO) QCD and EWK corrections. The motivation to study V+jets processes is hence two fold; (1) as a precision test of the SM in a new era of LHC, thus validating fundamental ingredients of state of the art Monte Carlo generators, and (2) increasing the sensitivity of searches for new physics that rely critically on reducing the systematic uncertainties on the V+jets background processes.

This paper is organised as follows. In Section 2 we provide an experimental overview of the V+jets measurements from ATLAS and CMS, highlighting a small subset of these measurements. The progress in theoretical developments is discussed in Section 3, including

the calculations of higher order perturbative QCD and EWK corrections, the status of various Monte Carlo generators and the constraints from V+jets processes on fits to PDFs. In Section 4 we select a few analyses to highlight the impact of V+jets processes on searches for new physics. We conclude in Section 5 by listing the key outcomes/wishlist resulting from the discussions at the workshop.

2 Experimental overview of V+jet measurements

Processes with a vector boson and jets are produced with large cross sections at a hadron collider and the high statistics allow for a wide range of measurements to be performed. In this section we use select measurements from ATLAS and CMS to highlight the critical role they play in testing perturbative QCD, constraining PDFs and estimating backgrounds to new physics searches.

The measurement of the production of a Z boson with jets, Z+jets, is a powerful test of perturbative QCD. The large production cross section and the fully reconstructable decay products in the Z boson decay to charged leptons gives a clean experimental signature that can be precisely measured. The process also constitutes a non-negligible background to searches for new physics and studies of the Higgs boson. The Z boson production cross section in association with up to seven jets was measured with the ATLAS detector using 3.16 fb^{-1} of data collected at a center of mass energy of 13 TeV [?]. The measurement is performed separately in the electron and muon decay channels and subsequently combined taking into account the correlations between systematic uncertainties. The cross section is measured as a function of the following observables; inclusive and exclusive jet multiplicity, ratio of jet multiplicities $N_{\text{jets}+1}/N_{\text{jets}}$, p_T of the leading jet, jet rapidity, angular separation between the two leading jets and their invariant mass, and the H_T where H_T represents the scalar sum of the p_T of all selected jets and leptons in the event.

The measured fiducial cross section after unfolding for detector effects is compared to the following theoretical predictions from both multi-leg matrix element matched and merged calculations and also fixed order calculations; SHERPA 2.2 with a matrix element calculation for up to 2 additional partons at NLO and up to 4 partons at LO using the Comix and OpenLoops matrix element generators merged with the SHERPA parton shower, MADGRAPH_aMC@NLO using matrix elements including up to 4 partons at LO interfaced to PYTHIA 8 and using the CKKWL merging scheme, MADGRAPH_aMC@NLO with matrix elements for up to 2 jets at NLO and matched to PYTHIA 8 using the FxFx merging scheme, ALPGEN 2.14 interfaced with PYTHIA 6, fixed order parton level calculation at NLO using BLACKHAT+SHERPA with up to 4 partons, and calculations of $Z+\geq 1\text{jet}$ at NNLO from N_{jetti} .

The measured cross section as a function of the inclusive jet multiplicity and H_T for $Z+\geq 1$ jet events is shown in Figure 1. The error bars denote the statistical uncertainty while the hatched bands represent the total uncertainty taken by adding the statistical and systematic uncertainties in quadrature. The typical uncertainty is of the order of 10% in the 1-2 jet bin, where the largest contribution is from the jet energy scale and resolution. The jet multiplicity distribution is well described by all theoretical calculations at low




Figure 1. Measured cross section as a function of the leading jet p_T for inclusive $Z + \geq 1, 2, 3, 4$ events.

multiplicity but the data starts to diverge from the predictions at higher multiplicity where the parton shower takes over. In general, distributions dominated by a single jet multiplicity are modelled well by fixed order NLO calculations. The LO MADGRAPH_aMC@NLO matrix element calculation produces a harder H_T spectrum compared to the data. This modeling of the H_T and related observables is significantly improved by the NLO matrix element and parton shower matched generators, SHERPA and MADGRAPH_aMC@NLO with FxFx. The recent $Z + \geq 1$ jet N_{jetti} NNLO prediction also describes well the H_T distribution and other key observables such as the leading jet p_T distribution not shown here. BLACKHAT+SHERPA underpredicts the high H_T tail, as can be expected from a fixed order NLO calculation missing the higher parton multiplicities added by a parton shower. This agreement is recovered by adding higher orders in pQCD, the recent $Z + \geq 1$ jet N_{jetti} NNLO prediction describes well the H_T distribution and other key observables such as the leading jet p_T distribution not shown here.

In addition to the cross section measurements of individual V+jets processes, the ratio of their cross sections are also interesting quantities, such as W/Z , Z/γ and W^+/W^- . A differential measurement of the ratio of cross sections of Z+jets and γ +jets was performed using the CMS detector at a center of mass energy of 8 TeV and using a dataset corresponding to an integrated luminosity of 19.7 fb^{-1} [1]. In the limit of high boson transverse momentum the LO QCD effects from the mass of the Z boson on the Z/γ ratio is small and

hence the ratio is expected to become constant at a boson p_T where the effects of the finite Z mass becomes negligible, around 300 GeV. At higher boson p_T , corrections from higher order perturbative QCD and EWK processes (as discussed in next section) can lead to a non-negligible dependence of the cross section on logarithmic terms that can become large, thus altering the flat behaviour of this ratio. In addition, this ratio is a crucial theoretical input in the determination of one of the key backgrounds to searches for new physics in the jets plus missing transverse momentum channel, $Z(\rightarrow \nu\nu)+\text{jets}$. This constitutes a dominant and irreducible background, and can account for up to 70% of the events in searches for Supersymmetry, dark matter and the invisible decay of the Higgs. These searches typically employ data driven techniques to determine the number of $Z(\rightarrow \nu\nu)+\text{jets}$ events in the signal region by defining orthogonal control regions in data and using simulation to extrapolate from the control region to the signal region. The Z/γ ratio is one of the key inputs in estimating $Z(\rightarrow \nu\nu)+\text{jets}$ from the statistically powerful control sample of $\gamma+\text{jet}$ events and the largest uncertainty is the theoretical uncertainty assigned to this ratio from missing higher order corrections, as discussed in detail in Section 4.

The CMS measurement of this ratio is performed in four regions; $N_{\text{jets}} \geq 1, 2, 3$, and $H_T > 300$ GeV and requires the vector bosons to have transverse momentum larger than 100 GeV. The unfolded data distributions are compared to predictions from several theoretical calculations; a QCD calculation at NLO for $Z+\text{jets}$ and $\gamma+\text{jets}$ from BLACKHAT+SHERPA for up to 3 jets, a LO multiparton matrix element calculation from MADGRAPH with up to 4 additional partons and interfaced with PYTHIA using the MLM matching scheme, and a simulation of $Z+\text{jets}$ using SHERPA. The $Z+\text{jet}$ events from MADGRAPH+PYTHIA generation and SHERPA are rescaled using a global NNLO K-factor calculated from FEWZ 3.1.

The differential cross section for $Z+\text{jets}$ production as a function of p_T^Z and $\gamma+\text{jets}$ production as a function of $p_{T\gamma}$ is shown in Figure 2 together with the ratio of the various theoretical predictions to the data. For $Z+\text{jets}$, MADGRAPH+PYTHIA describes the data well up to approx 150 GeV in p_T^Z , and then predicts a harder spectrum than the data, overpredicting the data by up to 40-50% above 600 GeV. SHERPA undershoots the data below p_T^Z of 50 GeV and then overpredicts by up to 30% at high p_T . BLACKHAT underpredicts the data by almost a consistent 10% for p_T^Z above 100 GeV. For $p_{T\gamma}$, BLACKHAT roughly reproduces the shape of the data distribution, but underestimates the rate by approximately 10-15%. MADGRAPH undershoots the data by up to 30% at low p_T but models well the region above 500 GeV.

The differential ratio of the Z and γ cross sections as a function of the boson p_T is shown in Figure 2 for the inclusive selection and $H_T > 300$ GeV. Systematics from jet energy scale, resolution, luminosity are considered as correlated between Z and γ and cancel in the ratio. The prediction from MADGRAPH is consistently 20% higher than data. BLACKHAT also overestimates the data at high p_T by around 20%. More discussion on this data-MC discrepancy follows in the next section and the inclusion of higher order QCD and EWK corrections.




Figure 2. Differential cross section of Z+jets production as a function of $p_{T,Z}$ and γ +jets production as a function of $p_{T,\gamma}$ for the detector corrected CMS data compared to several theoretical predictions.

3 Summary of theoretical developments

This section discussed the progress in theoretical developments relevant to V+jets processes, including the calculation of higher orders in perturbation theory and the current status and upcoming developments in state of the art Monte Carlo generators.

3.1 Higher order QCD and EWK corrections

Theoretical uncertainties on V+jets processes principally arise from three main sources; (1) missing higher order corrections, (2) uncertainties in input parameters such as parton distributions, masses and couplings, and (3) uncertainties in the parton/hadron transition including the fragmentation which is modeled by the parton shower, the hadronisation and the underlying event. The uncertainties from missing higher order corrections can be improved by the inclusion of higher orders and the resummation of large logarithms, those




Figure 3. Differential cross section of the ratio of Z+jets and γ +jets cross sections as a function of the boson p_T as measured by CMS and its comparison to theoretical predictions from BLACKHAT and MADGRAPH.

on the input parameters are be improved by a better description of the benchmark processes and on the parton-hadron transition by improving the matching/merging at higher orders and a better estimation of non-perturbative effects. While NLO QCD is the current state of the art, there have been rapid developments in the calculation of NNLO QCD with many results becoming available. The inclusion of higher order corrections from NLO EW effects have also become important as they are roughly similar in size to the NNLO QCD and become significantly large at high energies when probing higher transverse momenta and also near resonances. NNLO QCD calculations are emerging as the standard for high statistics benchmark processes like V+jets and are now available for all such processes. The calculations are available at the parton level and can compute the fiducial cross-sections but the codes are complicated to use and require significant CPU resources. However, they demonstrate all the features expected of such a calculation; a reduced dependence on the renormalisation scale and hence a reduction in the scale uncertainty, stabilisation of the perturbative series, more partons in the final state so perturbation theory can begin to reconstruct some of the shower, and will eventually lead to improved PDFs, hence further reducing the theory uncertainty. A comparison between the p_T^Z distribution from




Figure 4. Comparison of the NNLO p_T^Z distribution normalised by the NNLO Drell-Yan cross section with data from ATLAS.

ATLAS data and the NNLO calculation normalised to the NNLO Drell-Yan cross-section in Figure 8 shows that it significantly improves the agreement with data resulting in a substantially reduced scale uncertainty. The photon p_T has also recently been calculated to NNLO and a comparison with the p_T spectrum from ATLAS is shown in Figure 8. While the shape of the distribution does not show a significant improvement with data compared to the NLO description, the theoretical uncertainty is significantly reduced. The NNLO/NLO k-factor is around 10% and reasonably flat, with a slight increase at higher p_T , and both predictions are consistently below the data. The uncertainty from varying the scale is 2-3% for the NNLO prediction compared to 8-10% for the NLO. The effects of the electroweak corrections are included by rescaling the complete NNLO result by the change in the LO prediction when including one-loop electroweak effects. While the overall agreement between data and prediction gets worse when including the electroweak corrections, the normalised distribution shows that the shape is much better described with the corrections. The total uncertainty on the photon p_T distribution is approx 4% at low p_T and 10% at a p_T of 1 TeV, dominated by the PDF uncertainty.

The inclusive W+1jet production is important for calibrating the missing transverse momentum attributed to the neutrino. It has large NLO corrections of the order of 40% owing to new partonic configurations from the soft/collinear W radiation from dijet events. The NNLO corrections are relatively small leading to a significant reduction in the scale uncertainty and display good convergence of the QCD perturbative expansion.

The fiducial cross sections at 13 TeV for the inclusive and exclusive 1-jet bin are compared in Table 3.1. The NLO correction increases the LO result by 42% for the inclusive case and by 16% for the exclusive bin, while including the NNLO corrections increases the inclusive cross section by 3% while reducing the exclusive 1-jet cross section by 4%. These different corrections for the inclusive and exclusive case are due to jet veto logarithms




Figure 5. Comparison of the NNLO and NLO $p_{T\gamma}$ distribution with data from ATLAS and the ratio to the NNLO calculation (left). Shaded bands represent the uncertainty on the theoretical calculations. Also shown is the ratio to the NNLO calculation when including the effects of NLO EWK corrections (right) for the unnormalised (top) and normalised (bottom) distributions.

which can have a large impact on fixed-order cross sections in exclusive jet bins. The p_T^W distribution is shown in Figure 9. The NLO corrections above $p_T^W = 200$ GeV are at a maximum of 60% and then slowly decrease to 40% at a p_T^W of 1 TeV with an uncertainty from scale variation of 20%, while the NNLO corrections are 10% at p_T^W of 200 GeV and remain roughly constant out to high p_T , with an uncertainty from scale variation of a few percent. The corrections have a very different impact on the exclusive jet distribution owing to the jet veto logarithms which increase with the transverse momentum. The NLO correction is 10% at p_T^W of 200 GeV and increases to 70% for p_T^W of 800 GeV. The NNLO correction is roughly constant from p_T^W of 50 GeV at 10%. For the H_T distribution, there is a large k-factor for the NLO and significantly reduced but still sizeable NNLO corrections. The NLO corrections grow to 75% when the H_T is > 1 TeV, with a residual scale dependence of $\pm 15\%$. At NLO there are configurations containing 2 hard jets and a soft/collinear W boson that are logarithmically enhanced. These cannot occur at LO since the W boson must balance in the transverse plane against the single jet, thus leading to NLO corrections that are large but the QCD perturbative expansion shows convergence and stabilises when the NNLO corrections are included.

	σ_{LO} (pb)	σ_{LO} (pb)	σ_{LO} (pb)	K_{NLO}	K_{NNLO}
inclusive	$773.7^{+33.7}_{-36.8}$	$1099.3^{+57.8}_{-44.6}$	$1130.2^{+5.2}_{-8.7}$	1.42	1.03
exclusive	$773.7^{+33.7}_{-36.8}$	$895.7^{+16.0}_{-11.6}$	$863.2^{+10.5}_{-13.0}$	1.16	0.96




Figure 6. The sources of systematic uncertainty on the $p_{T\gamma}$ distribution from variations in the scale, PDF and the photon isolation (top) and the total uncertainty together with the ratio of the CMS data to the NNLO distribution including EWK corrections (bottom).




Figure 7. Comparison of the NNLO p_T^Z distribution normalised by the NNLO Drell-Yan cross section with data from ATLAS.

Also studied is the case of the boosted W where the the leading jet is required to have a $p_T > 500$ GeV. There are 2 distinct category of events that pass these selection cuts;




Figure 8. Comparison of the NNLO p_T^Z distribution normalised by the NNLO Drell-Yan cross section with data from ATLAS.




Figure 9. Comparison of the LO, NLO and NNLO p_T^W distribution for inclusive and exclusive $W+1$ jet production and the behaviour of the NLO and NNLO k-factors (below).

those where the leading jet is back to back with the high p_T boson and dijet events with the emission of a soft or collinear W boson from one of the jets. The first class of events occur at LO in the perturbative expansion of the W +jet process, while the second type of event first occurs at NLO. The correction in the fiducial cross section in going from the LO to NLO is large, with a k-factor of 2.84 due to this new event category that appears at NLO. The NNLO correction is smaller at 16% and the scale variation also decreases from 20% at NLO to less than 7% at NNLO. The NNLO correction is contained within the NLO scale variation band, indicating convergence of the perturbative expansion. The separation between the closest jet and W boson is shown in Figure 11. Events where the jet and W




Figure 10. Comparison of the LO, NLO and NNLO H_T distribution for inclusive and exclusive $W+1$ jet production and the behaviour of the NLO and NNLO k-factors (below).




Figure 11. The $\Delta R_{j,W}$ distribution for the case of a boosted W with a leading jet $p_T > 500$ GeV for LO, NLO and NNLO. Also shown are the k-factors for the NLO and NNLO are (below).

are back to back is shown in the region where $\Delta R_{j,W} > \pi$, while the region below this is quite broad and populated by the NLO configuration where a soft W boson is emitted from one of the jets. The NNLO effects are very similar to NLO below π . Since the lepton is emitted preferentially along the direction of the W , the $\Delta R_{j,l}$ distribution is similar.

3.2 Status of generators

In this section we review the status of several Monte Carlo event generators and their forthcoming releases. A new version of the SHERPA Monte Carlo event generator, SHERPA-2.2.3, was released in April 2017. In addition to bug fixes for all known issues in SHERPA-2.2.2, it contains extended support for UFO BSM physics, a new parton shower and the functionality to do on-the-fly variations of renormalisation scale, factorisation scale, α_s and PDFs. The multijet merging for loop induced processes has also been further tested. Further, the generator now includes approximate NLO EWK corrections in the existing NLO QCD multijet merging. This represents a first and useful step towards a complete NLO




Figure 12. Comparison of the Z/γ ratio vs. boson p_T in data from CMS with NLO QCD and NLO QCD+EWK predictions from SHERPA+OPENLOOPS.

QCD+EW matching and multijet merging. Figure 12 shows a comparison of the theoretical predictions from SHERPA+OPENLOOPS to CMS data for the Z/γ ratio versus boson p_T for events with $n_{jets} \geq 1$. The data is compared to NLO QCD and NLO QCD+EW and demonstrates an improvement in the data-MC comparison with the inclusion of EWK corrections.

The NLO QCD+EWK prediction from SHERPA+OPENLOOPS for the angular separation between the closest jet and the muon in the W+jets inclusive and exclusive process is shown in Figure 13 including the comparatively large subleading Born contributions owing to the phase space opened up by the W+2jet process. The NLO corrections are negative in the peak of the distribution at $\Delta R(\mu, j) \approx \pi$ and the subleading Born contribution becomes important at large ΔR . Also shown is the ATLAS data and its comparison with predictions from ALPGEN+PYTHIA W+jets MLM merged process, PYTHIA 8 with a W+jets QCD shower and dijet with a QCD and EWK shower, and SHERPA+OPENLOOPS with NLO QCD+EWK+subLO. The ATLAS data shows excellent agreement with the SHERPA+OPENLOOPS prediction.

The forthcoming release of SHERPA-2.3.0 will include parton shower reweighting and the full NLO EWK corrections.

V+jets processes are key for the phenomenological validation of NLO multi-jet merging as implemented in Monte Carlo event generators like MADGRAPH_aMC@NLO owing to their high statistics at the LHC and their utility in probing regions of phase space that are affected by both fixed order and matrix element calculations. The FxFx NLO multi-jet merging method is the default tool in MADGRAPH_aMC@NLO and has worked well, giving a very good overall description of the data with only 0, 1, 2 jets. It can be interfaced




Figure 13. Comparison of $\Delta R(\mu, j)$ NNLO p_T^Z distribution normalised by the NNLO Drell-Yan cross section with data from ATLAS.

with both PYTHIA 8 and HERWIG++ and has shown substantial improvement with respect to inclusive processes. One of the possible foreseen improvements will be inclusion of processes with massless particles at Born, for instance γ +jets and VBF. Also incorporated is an automated interface to UNLOPS to enable an independent evaluation of NLO multi-jet merging systematics on top of scale variations and underlying event shower modelling. The automated event generation for loop-induced processes have also been added to include gg effects for $Z+0, 1, 2$ jets.

3.3 PDF constraints from V+jets

Inclusion of transverse momentum distribution of the Z boson into PDF fits has significant constraining power. The experimental precision on the Z p_T distribution for $p_{T,\text{up}}$ to 200 GeV is $< 1\%$. Inclusion of Z p_T data constrains the intermediate- x gluon and light quark distributions.

4 Backgrounds to BSM searches

Backgrounds from V+jets processes contribute to many searches for BSM physics, in particular those involving missing transverse energy. The relatively large cross-sections for processes like $Z(\rightarrow \nu\nu)$ +jets and $W(\rightarrow l\nu)$ +jets means backgrounds from them are sizeable compared to the signal process. This section briefly highlights a few searches where

the our improved understanding of these processes in certain regions of phase space will play a critical role in driving the future sensitivity of these searches.

In the ‘monojet’ search looking for the presence of at least 1 jet and substantial \cancel{E}_T , the dominant backgrounds from $Z(\rightarrow \nu\nu)$ +jets and $W(\rightarrow l\nu)$ +jets are determined using a set of independent control regions in data. The control regions are defined such that they share similar kinematic characteristics with the signal region but are orthogonal to it. The control regions most commonly used are; W +jets, $Z(\rightarrow ll)$ and γ +jets. Transfer factors are determined that account for the lepton acceptance and efficiency, the difference in branching fractions between the control region process and the background process and the ratio of the production cross-sections. One of the key systematic uncertainties in the analysis is from the uncertainty on these transfer factors, in particular the theoretical component associated with the ratio of production cross sections in the extreme regions of phase space where the search is conducted. As seen in the previous section, the effects from higher order QCD and EWK corrections for V +jets processes and in the ratio of cross sections for Z/γ and W/Z at high transverse momentum become substantial. Hence, understanding these processes to better accuracy is critical.

The search for invisible decays of the Higgs boson also sees a large background contribution from V +jets processes. This background is particularly enhanced where the invariant mass of the dijet pair is large, with VBF production of Z +jets contributing around 30% to the signal region and carrying large uncertainties of 20-30%. The main source of uncertainty for the VBF production of the Higgs is from the W/Z ratio in this VBF phase space and has the largest impact on the result, while for a $V(jj)$ tagged analysis the dominant source of systematic uncertainty is from the theoretical uncertainty on the γ/Z ratio, followed by the W/Z ratio, as shown in Figure ???. It is therefore highly desirable to have Monte Carlo generators with NLO QCD and EWK corrections also for VBF topologies and the phase space probed by multi-jet searches typical of Supersymmetry searches.

5 Outcome/wishlist

This section gives a concise list of the key outcomes of the workshop and the ‘wishlist’ discussed by the theorists and experimentalists.

- V +jets processes play a very important role as fundamental tests of the Standard Model, from probing perturbative QCD to constraining PDFs and are also a crucial background in a multitude of BSM searches.
- Theoretical modeling of the inclusive V +jets process is under very good control, evidenced by the very good agreement between data and simulation for a range of different Monte Carlos.
- The inclusion of NLO EW corrections is crucial in the tails of high-energy distributions. Approximate fixed order NLO EWK corrections are available in SHERPA+OPENLOOPS 2.2. The exact fixed order EWK corrections will be available in SHERPA 2.3. Including these corrections in MADGRAPH_aMC@NLO is being worked on.

Systematic uncertainty	Impact		
Common			
γ +jets/ $Z(\nu\nu)$ +jets ratio theory	32%		
$W(l\nu)$ +jets/ $Z(\nu\nu)$ +jets ratio theory	21%		
Jet energy scale+resolution	12%		
V-tagging efficiency	12%		
Lepton veto efficiency	13%		
Electron efficiency	13%		
Muon efficiency	8.6%		
b jet tag efficiency	5.7%		
Photon efficiency	3.1%		
E_T^{miss} scale	4.6%		
Top quark background normalisation	6.0%		
Diboson background normalisation	< 1%		
Luminosity	< 1%		
Signal specific			
ggH p_T -spectrum	12%		
QCD scale + PDF (ggH)	3.0%		
QCD scale + PDF (VH)	1.4%		
Total statistical only	-46 / + 50%		
Total uncertainty	-69 / + 74%		
		Systematic uncertainty	Impact
		Common	
		W to Z ratio in QCD produced V+jets	13%
		W to Z ratio in EW produced V+jets	6.3%
		Jet energy scale+resolution	6.0%
		QCD multijet normalisation	4.3%
		PU mis-modelling	4.2%
		Lepton efficiencies	2.5%
		Luminosity	2.2%
		Signal specific	
		ggH acceptance	3.8%
		QCD scale + PDF (qqH)	1.8%
		QCD scale + PDF (ggH)	< 0.2%
		Total statistical only	-27 / + 28%
		Total uncertainty	-33 / + 32%

Figure 14. The key sources of systematic uncertainties and their impact on the fitted value of $B(H \rightarrow \text{inv.})$ in the (a) V(jj)- tagged analysis and (b) VBF-tagged analysis.

- Recommendations for applying corrections to account for NLO EWK effects and evaluate uncertainties associated with them for inclusive searches looking for jets and \cancel{E}_T have recently been finalised and are available in[].
- NNLO QCD reduces scale uncertainties to the O(5%) level for individual distributions. It is desirable to quantify the correlations between kinematic distributions and validate the different calculations using different methods, for instance antenna subtraction vs Njettiness slicing.
- The version number used for Monte Carlo generators should be specified.
- The agreement between data and simulation deteriorates in more exclusive phase-space regions, for instance the high invariant mass of dijet pair in VBF production. It is desirable to understand the reasons for these differences between the various Monte Carlos.
- Important to publish more exclusive distributions of kinematic quantities e.g H_T distribution in bins of jet multiplicity, 2D distributions to show correlations between variables e.g H_T vs p_T^Z , in $p_T(V)$ vs $\Delta\phi(j1, j2)$. For the case of collinear boson emission and the observable of interest, the angular separation between the boson and the closest jet, the region of this distribution in between the two extremes (dijet collinear with boson and back to back dijets) is interesting.
- Publish more jet-observables & leptonic observables.
- Where possible, make available the unnormalized distributions, or the provision of k-factor used to normalize the overall cross-section. NNLO k-factors obtained for

inclusive sample ($N_{\text{jet}} \geq 0$) are not always applicable to less inclusive distributions ($N_{\text{jet}} \geq 1, 2$). One possibility is to have normalized distributions in the public note and unnormalized ones in HEPDATA.

- Need higher-order EW corrections for QCD & EW V+jets in VBF topologies. SHERPA includes QCD corrections for VBF topologies, also EWK corrections to QCD production but not QCD corrections to EWK production. At higher order there are also interferences between QCD and EWK production, need to calculate V+jets at all sub-leading one loop orders[?].

References

- [1] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1510** (2015) 128 Erratum: [JHEP **1604** (2016) 010] doi:10.1007/JHEP04(2016)010, 10.1007/JHEP10(2015)128 [arXiv:1505.06520 [hep-ex]].