

Preliminaries

- There is a pre-class assignment due on Monday (see Canvas).

As part of this pre-class assignment, if you have not already, record the pronunciation of your name via my.harvard


- Skill Check 02 will be on Monday. An example problem is below.
- The first problem set is posted on Gradescope and is due at noon on Friday. Find the office hours schedule for next week on Canvas.

All students can extend problem set submissions until 8pm on Friday, and have access to up to three 32 hour late days for problem sets over the course of the semester.

Skill Check 02 practice

The skill check will be during class on Monday. It will be similar, but not identical, to the question below.

Consider the differential equation $\dot{x} = f(x)$ for $f(x)$ given by the graph of $f(x)$ vs x below. Draw a phase portrait for $\dot{x} = f(x)$ on the blank x -axis provided below the graph. Use the conventional shading scheme to indicate stability type for fixed points (including properly half-shaded for half-stable fixed points).



Skill check solution



Answer:

Extra notes:

1. Avoid arrows at the ends of the x axis (because those can be confused with the direction arrows for the vector field flow).
2. Label the axis (x).
3. Add fixed points as open circles. Fixed points occur when $f(x) = 0$.
4. Add a single arrow showing the direction of $\frac{dx}{dt}$ in each region where $f(x) < 0$ or $f(x) > 0$.
5. Use the arrows to shade the fixed points. Fill in the fixed point (or the appropriate half of the fixed point) when flow is towards it.

Activity

Teams

- | | |
|--------------------------------|----------------------------|
| 1. George, Alexander, Margaret | 5. Emily, Shefali, Michail |
| 2. Camilo, Sophie, David H | 6. Noah, Iona, Isaiah |
| 3. Thea, Joseph, Mariana | 7. Katheryn, Mallory |
| 4. Van, David A, Hiro | 8. Ada, Alice |

All Teams: Write your names in the corner of the whiteboard.

Teams 1 and 2: Post screenshots of your work to the course Google Drive today (make or use a 'C03' folder). Include words, labels, and other short notes on your whiteboard that might make those solutions useful to you or your classmates. Find the link in Canvas.

Team activity

- (practice classifying stability: Strogatz 2.4.7)

Let $\dot{x} = rx - x^3$ where the parameter r satisfies either $r < 0$, $r = 0$, or $r > 0$.

Find the fixed points and classify their stability using linear stability analysis (an algebraic method). If linear stability analysis does not allow you to classify the point because $f'(x^*) = 0$ then note that. Such fixed points are called *non-hyperbolic*.

Discuss all three cases.

- (more parameter dependence) Let $\dot{x} = r + x^2$.

(a) Find the fixed points algebraically as a function of r

(b) Make phase portraits for $r = -2, -\frac{1}{4}, 0, 1$.

(c) Using r as the vertical axis, place these phase portraits in an xr -plane.

The $r = -2$ portrait will be at the bottom with the others above it, sketched at the appropriate values of r .

(d) Draw the location of stable fixed points in the xr -plane using a solid curve. Draw the location of unstable fixed points using a dashed curve.

(e) Rotate your axes: in the rx -plane (r as the horizontal axis), sketch the solid and dashed lines that summarize the locations and stability of the fixed points.

How does this diagram encode the information in the phase portraits?

This diagram is referred to as a 'bifurcation diagram'

- (oscillation: Strogatz 2.6.1) A simple harmonic oscillator, defined by $\ddot{x} = -\frac{k}{m}x$, has a solution $x(t) = A \sin \omega t + B \cos \omega t$ that oscillates with time.

(a) Plug this expression for $x(t)$ into the differential equation to show that it is a solution for some ω . Find that ω .

(b) What happens to A and B ?

(c) We learned that oscillations are not possible in a one-dimensional system. This system is showing oscillations. Reconcile those two facts.

4. Compare the populations models

$$\dot{N} = N(1 - N/K)$$

(logistic) and

$$\dot{N} = N(1 - N/K)(N/A - 1)$$

(strong Allee effect) where $0 < A < K$.

- (a) Based on the differential equation, what is the Allee effect?
 - (b) Try to imagine a scenario where it is relevant (it was initially described in experiments on small fish).
 - (c) Consider solutions, $N(t)$, to both equations. How, if at all, do solutions between the two equations differ qualitatively?
 - (d) The term *basin of attraction* refers to the set of initial conditions that approach a particular fixed point. What is the basin of attraction of the extinction fixed point, $N^* = 0$, for each equation?
5. (even more parameter practice. Strogatz 3.2.3) Let $\dot{x} = x - rx(1 - x)$. Sketch each of the qualitatively different phase portraits that occurs as r is varied. Sketch the bifurcation diagram of fixed points vs r in the rx -plane. Use solid and dashed lines to indicate the stability of the fixed points in your diagram.