

- Which of the following anterior teeth is widest mesiodistally:
- Mandibular central incisor.
 - Mandibular canine.
 - c Maxillary central incisor.
 - Maxillary canine.

The number of major fossae in occlusal surface of permanent maxilla

- S
- a Three fossae.
 - Two fossae.
 - Five fossae.
 - Four fossae.

3. The mamelons is the external manifestation of :

- S
- Cusps.
 - Lobes.
 - c Labial ridge.
 - Tubercle.

4. The mandibular first premolar develops from the following number

- a
- a 4 lobes.
 - b 5 lobes.
 - 3 lobes.
 - 6 lobes.

5. Small, irregular grooves do not denote the union of primary

- S
- a Supplemental grooves.
 - b Developmental grooves.
 - Buccal grooves.
 - Lingual grooves.

6. The neck of the tooth is called

- Crypt.
- Socket.
- c Cervical line.
- Root trunk.

7. The point angles:

- S
- a Four in number in anterior and posterior teeth.
 - x b Do not include the incisal surface.
 - Are found in posterior teeth only.
 - Derive their name from the 2 surfaces that

The dental formula of 4 is:

- Left mandibular first premolar.
- b Right maxillary first premolar.
- Primary right mandibular first premolar.
- Permanent right mandibular first molar.

9 . The second deciduous molars similar to:

- a. Permanent first molars. ●
- b. Premolars.
- c. Permanent second molars.
- d. Permanent third molars.

10 . The oblique ridge in the maxillary first molar connects:

- 3.
- a. Distolingual and Mesio Buccal triangular ridges.
 - b. Distal and mesial triangular ridges.
 - c. Mesiolingual and Distobuccal triangular ridges. ●
 - d. None of the above.

11 . There are two types of maxillary second molars when viewing

- a. 5 cusps type and 4 cusps type.
- b. 4 cusps type and 3 cusps type. ●
- c. 3 cusps type and 2 cusps type.
- d. None of the above.

12 . In occlusal aspect, the outline form of permanent maxillary first

- a. Rhomboidal. ●
- b. Trapezoidal.
- c. Rectangular.
- d. Oval.

13 . From the buccal aspect, the mesiobuccal cusp of the permanent

- 3.
- a. Narrower than distobuccal cusp and its slopes meets e
 - b. Broader than distobuccal cusp and its slopes meets eac
 - c. Sharper than the distobuccal cusp.
 - d. Broader than mesio-lingual cusp.

14 . In maxillary third molar:

- 3.
- a. The distobuccal cusp is wider and longer than mesiob
 - b. The distal contact area is flat and larger than mesial co
 - c. The occlusal surface has numerous supplemental groo
 - d. Both a and b.

15 . The linear elevation on the lingual aspect of incisors are :

- a. Marginal ridges. ●
- b. Lingual ridges.
- c. Cingulum.
- d. Mamellons.

16 . From the lingual aspect of the permanent mandibular first mola

- a. The mesiobuccal and distobuccal cusps.
- b. The distobuccal cusp and the distal cusp.
- c. Two lingual grooves.
- d. Two lingual cusps and the lingual portion of the dista

Mandibular primary second molar has:

- ☒ a. Five cusps.
- b. Four cusps.
- c. Three cusps.
- d. Six cusps.

Regarding the permanent mandibular first molar:

- ☒ a. Mesiodistal measurement is 1mm greater than the buccolingual.
- b. The buccolingual measurement is greater distally than mesially.
- ☒ c. The distal cusp is the biggest cusp.
- d. There are two major fossae and one minor fossa.

19. The occlusal aspect of maxillary first premolar is:

- a. Rhomboid.
- b. Square.
- ☒ c. Hexagonal.
- d. Pentagonal.

20. From the occlusal aspect, the maxillary second premolar differs from the first premolar:

- a. It has more rounded outline form.
- b. It has shorter central groove.
- c. It has no mesio-marginal developmental groove.
- ☒ d. All of the above.

The mesial aspect of maxillary first premolar characterized by:

- ☒ a. Presence of mesial developmental depression.
- b. Longer lingual cusp compared to the buccal cusp.
- c. The buccal crest of contour located in the occlusal third of the crown.
- d. None of the above.

22. The oblique ridge is found on the occlusal surfaces of:

- a. The deciduous maxillary first and permanent upper first molar.
- b. The deciduous mandibular second and permanent upper first molar.
- ☒ c. The deciduous maxillary second and permanent upper first molar.
- d. None of the above.

23. Which of the following permanent posterior teeth has "Y" shaped occlusal surface:

- a. Permanent maxillary first molar.
- b. Permanent mandibular first molar.
- ☒ c. Permanent mandibular second premolar.
- d. Permanent maxillary second premolar.

24. Which statement is true about maxillary first premolar cusps:

- ☒ a. The buccal cusp is about 1 mm longer than lingual cusp.
- b. The lingual cusp is about 1 mm longer than buccal cusp.
- c. Both the buccal and lingual cusps are of same height.
- d. None of the above.

25 . Which characteristic(s) of mandibular first premolar resemble(s) second premolar:

- a. The contact areas mesially and distally are nearly at the same level.
- b. The curvatures of cervical line mesially and distally is similar.
- c. The tooth has more than one cusp.
- d. All of the above.

26 . When maxillary first premolars have two roots, they are placed

- a. One mesially and other distally.
- b. One buccally and other lingually.
- c. Both roots are placed lingually.
- d. None of the above.

27 . Concerning the pulp cavities:

- a. The outline of the pulp cavity corresponds to the external contour of the tooth.
- b. The dental pulp is the soft tissue component of the tooth.
- c. The space in which the pulp tissue is located.
- d. All of the above.

28 . Lateral (Accessory) canal:

- a. It is a canal that extends in a horizontal direction from the root into the periodontal ligament.
- b. It is most often found in the cervical half of the root.
- c. It is located in the anatomical root of the tooth.
- d. None of the above.

29 . In deciduous pulp cavities differ from the permanent teeth in:

- a. The pulp chamber of deciduous teeth is irregular.
- b. The pulp horns in deciduous molars are high and more numerous.
- c. The pulp cavities are proportionally much smaller than in permanent teeth.
- d. All of the above.

30 . The canine fossa is important character to distinguish:

- a. Permanent maxillary canine.
- b. Permanent mandibular canine.
- c. Maxillary first premolar.
- d. Mandibular first molar.

31 . The cross section of the root of Maxillary canine at cervix:

- a. Oval.
- b. Rounded.
- c. Rectangular.
- d. Triangular.

32 . The distinguish feature in mandibular first premolar is:

- a. The mesio-marginal developmental groove.
- b. Round occlusal form.
- c. No lingual inclination.
- d. Transverse ridge between buccal and lingual cusp.

constrict openings at the root apex through which the blood and nerve supply enter to the pulp called:

- a. Apical foramen. ●
- b. Lateral canal.
- c. Supplemental canals.
- d. Anastomosis

34 . The tooth that have three root canals is:

- a. Permanent mandibular first molar.
- b. Permanent maxillary first molar.
- c. Both a and b. ●
- d. Maxillary first premolar.

35 . The significant point of the proximal contact area:

- a. Stabilize the dental arches. ●
- b. It allows escapements of food.
- c. Prevent soft tissue biting during mastication.
- d. Increase the force of mastication.

36 . Embrasure:

- a. Spillways to bring the food toward the teeth.
- b. A triangular spaces filled by gingival tissue.
- c. A (V) shaped spaces away from contact areas. ●
- d. A lingual height of contours.

To prevent cusps clash between upper and lower posterior teeth, the proximal form should :

- a. Rhomboidal in maxillary and trapezoidal in mandibular.
- b. Rhomboidal in mandibular and trapezoidal in maxillary. ●
- c. Trapezoidal in all maxillary and mandibular posterior teeth.
- d. Triangular in maxillary and mandibular posterior teeth.

38 . Triangular crown form of anterior teeth from proximal aspects:

- a. Guide the mandible through various lateral movement
- b. Increase anchor of root and prevent rotation.
- c. Wide base for crown strength. ●
- d. Increase the force of mastication.

The tooth supporting structure:

- a. Alveoli.
- b. Crypt.
- c. Periodontium. ●
- d. Dentin.

The mixed dentition:

- a. Started at age 6 to 12 years. ●
- b. Started by eruption of premolars at 12 years.
- c. Started at age 6 month-6 years.
- d. A and C.

The difference between deciduous and permanent dentition:

- a. The roots of deciduous molars are more divergent. ●
- b. The pulp chamber is much smaller in deciduous teeth.
- c. The cervical ridge buccally on deciduous molars is less pronounced than the permanent.
- d. The crowns of permanent teeth appear constricted at the cervix.

The following is character of deciduous dentition:

- a. Longer crown than root.
- b. Crown length equal to root length.
- c. Crown width more than crown length.
- d. None of the above. ●

3 . The successors of deciduous molars are:

- a. Permanent incisors.
- b. Permanent molars.
- c. Permanent canines.
- d. Premolars. ●

44 . Over developed facial and lingual contours of the teeth results in:

- a. Proper gingival massage.
- b. Food will push gingival tissue apically.
- c. Provide too much protection of the gingiva. ●
- d. Gingival recession.

45 . Concerning the permanent dentition :

- a. The maxillary teeth tend to erupt before mandibular teeth.
- b. The eruption may be affected by sex. ●
- c. The canine erupts before premolar.
- d. The eruption not affected by racial factor.

The trapezoidal form of proximal surfaces of upper posterior teeth provides :

- a. Wide interproximal spaces.
- b. Proper embrasures.
- c. Narrow occlusal surfaces to be easily forced through the food. ●
- d. Positive contact between the teeth.

The interproximal spaces are triangular in shape, the base, side walls and apex of the triangle respectively is formed of:

- a. Alveolar process, proximal surfaces of adjacent teeth and contact areas.
- b. Contact areas, proximal surfaces of adjacent teeth and alveolar process.
- c. Cervical lines of adjacent teeth, contact areas and alveolar process.
- d. Alveolar process, facial surfaces of adjacent teeth and contact areas.

axial proximal contact areas are:

- a. Narrow in the posterior teeth.
- b. Centered buccolingually in posterior teeth.
- c. Situated more to lingual in anterior teeth.
- d. Situated slight to the buccal in posterior teeth. ●

The size of embrasure is determined by :

- a. Location of contact area. ●
- b. Location of interproximal space.
- c. Location of cervical line.
- d. None of the above.

Under developed facial and lingual contours of the teeth lead to :

- a. Gingival recession.
- b. Food will push gingival tissue apically.
- c. Excessive frictional irritation of the gingiva.
- d. All of the above. ●

Oral exam 10 mark Put (T) or (F) for the following.

-) Permanent maxillary first molar is same to the deciduous maxillary second
-) Permanent maxillary first molar is smaller than permanent maxillary second
-) Permanent maxillary second premolar has long central groove.
-) five well formed cusps are located at $\overline{6}$.
-) The segments of dental arch are: anterior, middle and posterior.
-) The embrasure is " V " shape space started away of trifurcation of root.
-) cross-shape central groove is seen at permanent mandibular third molar.
-) Curve of Spee is mediolateral relation of maxillary and mandibular teeth
-) Leeway space is the difference in the size of permanent premolars.
-) sphere of Monsoon is both of curve of Spee and Bonwill equilateral triangle