

CANINES

DR. M. 3EED

CANINES

General characteristics:

- *There are one canine in each quadrant. Canine may be named as **Cuspid**.
- *Canines have longer and stronger roots than those of other teeth.
- *over the labial aspect of their roots, there is well developed bone ridge called the **canine eminence**, giving the canine a very important cosmetic value because they ensure facial expression.
- *Canines are formed of **4 lobes (3 labial and 1 lingual)**, the middle lobe is highly developed **labially** forming the **labial ridge** and **lingually** forming the **lingual ridge** and **incisally** forming a **cusp**, therefore canines are called the cuspids.
- *the canines are located between incisors and premolars, so they support the two groups in function (**tearing and partial cutting**).

	Begainig of calcification	Crown completion	Eruption	Root completion
$\frac{1}{1}$	3-4 month	3 years before eruption date	7 years	3 years after eruption date
1			6 years	
$\frac{2}{2}$	10-12 months		8 years	
2	3-4 months		7 years	
$\frac{3}{3}$	4-5 months		11 years	
3			9 years	



1) Labial aspect	Upper Canine	Lower Canine
GEOL	Trapezoid with the smallest uneven side cervically and the largest side incisally.	
		
All outlines	☒ -Mesial: convex ☒ -Distal: concave ☒ -Incisal: ✓ Mesial cusp slope : concave ✓ Distal cusp slope : convex DCS > MCS ☒ Cervical line : convex root wise	☒ The same but with small differences :- ☒ -its is narrower than the upper canine. ☒ -it's crown appear longer than upper canine because of the narrow neck and the more incisally located incisors. ☒ -the mesial outline is straight ☒ -the cusp tip is less pointed
Contact areas	MCA: at the J. of M and I 1/3 DCA: more cervically , located in middle 1/3	MCA: in the incisal 1/3 DCA: more cervically. at the junction of middle and incisal 1/3
Surface anatomy		
1)Elevation	☒ (cervical ridge) : at the cervical 1/3 ☒ Labial ridge: prominent from cusp tip to cervical ridge.	The same but less developed.
2)Depressions	-2 shallow developmental grooves which divide the labial surface into 3 lobes .	
ROOT	Single root The apex is pointed and <u>curved distally</u> . The root is the longest in all teeth.	Single root curved distally , but slightly shorter than the lower canine.

2) Lingual aspect	Upper Canine	Lower Canine
N.B	Both crown and root are narrower lingually than labially due to (lingual convergence)	
		
Surface anatomy	<u>well</u> developed :	Less developed :
1) Elevation	✓ Cingulum : ✓ MMR ✓ DMR ✓ lingual ridge is prominent and extends from cusp tip to cingulum	✓ Cingulum: ✓ MMR ✓ DMR ✓ The lingual ridge is restricted and found <u>only in incisal 1/3.</u>
2) Depressions	- 2 lingual fossae	- 1 lingual fossa , due to the less developed lingual ridge.

3) Mesial aspect	Upper Canine	Lower Canine
Geol	Triangular ☒ Base : Cervically ☒ Apex : Incisally	
Outines		
A) Labial outline:	❖ In Cervical 1/3 : convex > Cervical Ridge ❖ In middle and incisal 1/3 : Sl. Convex	
B) Lingual outline:	❖ In Cervical 1/3: Convex > Cingulum ❖ In Middle 1/3: concave > Fossa ❖ In Incisal 1/3 : Convex > Incisal ridge	
N.B	The cusp tip of the crown is labial to the center of the root	-the cusp tip is located lingual to the center of the root due to lingual inclination.

4) distal aspect	Upper canine	Lower Canine
N.B	Same as mesial aspect except :- -The cervical line has less curvature toward the cusp tip -The contact areas is still centralized but is located more cervically. -The depression on the distal aspect of the root is more pronounced than on mesial aspect . - this depression on the root helps in anchorage and stanilization of the tooth in the arch , to preventrotation and displacement.	
اوعى تنسى دي		The developmental depression on the root in lower canine may be deep that it can divide the root into 2 roots : 1 labial + 1lingual roots 

DR.

5) Incisal aspect	Upper Canine	Lower Canine
Geol	Diamond , and has (lingual convergence)	
Dimensions	Thickness > width	
N.B	-the cusp tip is labially and mesially situated <u>Labially due to : this teeth is upper tooth</u> <u>Mesially due to : the DCS is > MCS)</u> -The labial ridge is very well developed -The lingual ridge is well developed so there are 2 lingual fossae. -The cingulum and marginal ridges are also seen from this aspect. 	The cusp tip is lingually and mesially situated <u>lingually due to lingual inclination.</u> <u>Mesially due to : the DCS is > MCS)</u> -The lingual ridge is less developed restricted to the incisal third so there is only one fossa 

6)Pulp cavity	Upper Canine	Lower Canine
Pulp cavity	One pulp cavity	The possibility of presence 2 seprate root canals (1 labial and 1 lingual) if the lower canine has 2 roots .
Pulp horn	One pulp horn	One pulp horn

DR. M. 3EED