

Dental eruption dates

Emergence of the primary dentition takes place between the sixth and thirtieth months of postnatal life.

primary dentition is completely formed by about age 3.

primary teeth lost completely at about age 11-12.

The predominant sequence of eruption of the primary teeth in the individual jaw is:

- 1- central incisor.
- 2- lateral incisor.
- 3- first molar.
- 4- Canine.
- 5- and second molar.

At about (6 to 10) months of age, **the mandibular central incisors** emerge through the alveolar gingiva, followed by the other anterior teeth.

so that by about 13 to 16 months, **all eight primary incisors have erupted.**

The first primary molars emerge by about 16 months of age.

NOTE: The maxillary first molar tending most often to erupt earlier than the mandibular first molar.

The primary maxillary canines erupt at about 19 months, and **the primary mandibular canines** erupt at 20 months.

The primary second mandibular molar erupts at a mean age of 27 months, and **the primary maxillary second molar** follows at a mean age of 29 months.

Permanent teeth Begins to emerge at 6 years.

Eruption process of Permanent teeth is completed by 12 -13 years except for the 3rd molars.

Stages of dentition in human

(A) Deciduous dentition period (8 months to 6 years):

From the eruption of mandibular primary central incisors to the eruption of mandibular Permanent 1st molar.

(mandibular primary central incisors -to- mandibular Permanent 1st molar).

(B) Transitional (Mixed) dentition period (6 to 12 years):

The first transition dentition begins with the emergence and eruption of **the mandibular first permanent molars** (6 years) and ends with the loss of the last primary tooth, which usually occurs at about age 11 to 12 (usually Maxillary primary canine).

(the mandibular first permanent molars -to- loss of the last primary tooth (usually Maxillary primary canine) (11 to 12 years)).

Primary teeth exfoliated followed by eruption of their permanent successors.

(C) Permanent dentition period (12 years and beyond):

Is well established by about 13 years of age with the eruption of all Permanent teeth except the 3rd molars.

NOTE: The mandibular permanent teeth tend to erupt before maxillary teeth.

The mandibular central incisor usually erupts around (6 to 7 years).

(Erupts before the maxillary central incisor and may erupt simultaneously with or even before the mandibular first molar).

Mandibular lateral incisors: (7-8 years). (erupt very soon after the central incisors, often simultaneously).

Maxillary central incisors: (7-8) years.

Maxillary lateral incisors: (8-9) years.

Mandibular canines (cuspids): (9-10) years.

Maxillary first premolars: (10-11) years.

Mandibular 1st premolar & Mx 2nd premolars: (10 -12) years.

Mandibular 2nd premolar & maxillary canines: (11-12) years.

Mandibular second molars: (11-13) years.

Mx second molar: (12-13) years.