


A dark blue vertical bar runs down the left side of the page. A light blue arrow points to the right from the bar, containing the date.

5/27/2021

Dental Anatomy

MCQs

Several thin, curved lines in shades of blue and grey sweep upwards from the bottom left corner of the page.

Helpers 48

FACULTY OF DENTISTRY, TANTA

Section 1

Introduction.

Choose the correct answer(s):

1. If you read an article in a British dental journal that refers to tooth No. 48, you would suspect that the authors were using the International Numbering System. What universal number (or letter) would they be talking about?

- a. 25 b. J c. 30 d. T **e. 32**

2. Using the Universal Numbering System, what numbers are used to identify maxillary canines?

- a. 6** b. 8 c. 10 **d. 11** e. 27

3. Which tooth junctions are NOT normally visible on a handheld intact tooth?

- a. Cementoenamel junction **b. Dentinoenamel junction**
c. Dentinocemental junction **d. Dentinopulpal junction**

4. Which statement(s) is (are) likely to be true on a person with a barely erupted tooth No. 9?


- a. The clinical crown is larger than the anatomic crown
b. The clinical crown is smaller than the anatomic crown
c. The clinical root is larger than the anatomic root
d. The clinical root is smaller than the anatomic root

5. Which tooth surface(s) face(s) the lips or cheeks?

- a. Facial** b. Distal **c. Buccal** d. Occlusal **e. Labial**

6. Which pairs of teeth have a mesial surface touching a mesial surface?

- a. 25 and 26 b. 16 and 17 c. 7 and 8
d. 1 and 32 **e. 8 and 9**



7. When viewing tooth No. 8 from the distal view, it can be divided into thirds from the incisal to the cervical and from the facial to the lingual. Which third is NOT possible to see from the distal view?

- a. Facial b. Cervical c. Middle **d. Mesial** e. Incisal

8. If you were observing the faciolingual dimension of a tooth, what surface(s) could you be viewing?

- a. Mesial** **b. Occlusal** **c. Proximal** d. Labial **e. Distal**

9. If the root-to-crown ratio of a maxillary molar (No. 14) is 1.72 and that of another molar, No. 16, is 1.49, which tooth has the longest root relative to its shorter crown?

- a. No. 14** b. No. 16 c. More information is required in order to answer this question

10. Which of the following bumps or ridges is NOT likely to be found on a maxillary premolar?

- a. Oblique ridge** **b. Cingulum** c. Mesial marginal ridge
d. Transverse ridge e. Triangular ridge

11. Which ridges surround the perimeter of the occlusal surface (occlusal table) of a two-cusped premolar?

- a. Mesial marginal ridge** **b. Distal marginal ridge**
c. Mesial cusp ridge of the buccal cusp **d. Distal cusp ridge of the lingual cusp**
e. Transverse ridge

12. What is the correct order of anatomic landmarks of a tooth with two roots from the cementoenamel junction to the root tip?

- a. Cervix, trunk, furcation, apex** b. Trunk, cervix, furcation, apex
c. Trunk, furcation, cervix, apex d. Cervix, trunk, apex, furcation
e. Furcation, trunk, cervix, apex



Helpers 48

13. When viewed from the proximal views, what is the location of the greatest bulge (crest of curvature or height of contour) on the facial surface of all teeth?

- a. Occlusal third b. Lingual third
- c. Buccal third d. Middle third **e. Cervical third**

14. Which space(s) contain(s) the part of the gingiva known as the interdental papilla?

- a. The buccal embrasure b. Occlusal embrasure
- c. Lingual embrasure **d. Cervical embrasure**
- e. Interproximal space**

15. Ideal class I occlusion involves an important first permanent molar relationship where the mesiobuccal cusp of the maxillary first molar is located within the

- a. Mesiobuccal groove of the mandibular first molar**
- b. Distobuccal groove of the mandibular first molar
- c. Buccal groove of the mandibular second molar
- d. Mesiobuccal groove of the mandibular second molar
- e. Distobuccal groove of the mandibular second molar

16. Where do lingual cusps of maxillary teeth occlude in ideal class I occlusion?

- a. In the buccal embrasure space between mandibular teeth
- b. In the lingual embrasure space between mandibular teeth
- c. In occlusal fossae of mandibular teeth**

17. How many developmental lobes form a premolar with two cusps (one buccal cusp and one lingual cusp)?

- a. 1 b. 2 c. 3 **d. 4** e. 5



Helpers 48

Section 2

Permanent Incisors.

A) Choose the correct answer(s):

- a. Maxillary central incisor
- b. Maxillary lateral incisor
- c. Mandibular central incisor
- d. Mandibular lateral incisor

1. Mesiodistal dimension of the crown is larger than the labiolingual dimension.
A **b** c d
2. The incisal ridge exhibits a distolingual twist.
A b c **d**
3. The root is very narrow mesiodistally with mesial and distal root depressions
A b **c** **d**
4. The incisal edge is positioned more to the lingual of the root axis line.
A b **c** **d**
5. The distal proximal height of contour is more cervical than the mesial height of contour.
A **b** c **d**
6. This tooth has the widest (mesiodistally) incisor crown.
A b c d
7. This tooth has the shortest root relative to its crown.
A b c d
8. This tooth is the most symmetrical incisor.
A b **c** d
9. This tooth has the largest curvature of the mesial cervical line.
A b c d
10. This tooth has the narrowest incisor crown (mesiodistally).
A b **c** d



Helpers 48

B) Choose the correct answer:

1. The following are the first permanent teeth to erupt along with the first permanent molars:
 - a. Maxillary central incisors
 - b. Mandibular central incisors**
 - c. Maxillary lateral incisors
 - d. Mandibular lateral incisors
2. Smallest tooth in permanent dentition is:
 - a. Maxillary central incisor
 - b. Maxillary lateral incisor
 - c. Mandibular central incisor**
 - d. Mandibular lateral incisor
3. Among incisor teeth, the crowns are more symmetrical in:
 - a. Maxillary incisors
 - b. Mandibular incisors**
 - c. Both of the above
 - d. None of the above
4. The mesial and distal contact areas are nearly at the same level in:
 - a. Maxillary incisors
 - b. Mandibular incisors**
 - c. Both of the above
 - d. None of the above
5. Which of the following statement is true?
 - a. In the maxillary arch, central incisor is larger than the lateral
 - b. In the mandibular arch, lateral incisor is larger than the central
 - c. Both of the above**
 - d. None of the above
6. In the incisor group, which tooth exhibits greatest variation in its morphology?
 - a. Maxillary central incisor
 - b. Maxillary lateral incisor**
 - c. Mandibular central incisor
 - d. Mandibular lateral incisor



7. Both the contact areas are very much nearer to incisal edge in:
- a. Maxillary incisors
 - b. Mandibular incisors**
 - c. Both of the above
 - d. None of the above
8. When compared to maxillary incisors, the lingual fossa of mandibular incisors is:
- a. Deeper
 - b. Exhibit more developmental grooves
 - c. Shallower without developmental grooves**
 - d. Both a and b
9. Lingual pits are often present in:
- a. Maxillary incisors**
 - b. Mandibular incisors
 - c. Both of the above
 - d. None of the above
10. When viewed from proximal aspect, the incisal ridge/edge of maxillary incisors is placed:
- a. In line with the vertical root axis**
 - b. Labial to the line with the vertical root axis
 - c. Lingual to the line with the vertical root axis
 - d. None of the above



Helpers 48

Section 3

Permanent Canines.

A) For each trait described below, indicate the letter of the best response from the five selections provided. Each trait has only one best answer.

- a. Maxillary central incisor
- b. Maxillary canine
- c. Mandibular canine
- d. All of the above
- e. None of the above



Helpers 48

1. This tooth exhibits less cervical line curvature on the distal aspect than on the mesial aspect.

A B C **D** E

2. The cingulum is centered mesiodistally.

A **B** C D E

3. There is an almost continuous crown–root outline on the mesial surface of this tooth.

A B **C** D E

4. The mesial contact area is located more incisally than the distal contact area on the same tooth.

A B C D E

5. The cusp tip is positioned lingual to the mid-root axis line from the proximal view.

A B C **D** E

6. Mamelons could be observed on this tooth.

A B **C** D E

7. On which tooth is the cusp angle most acute?

A B C D E

8 The mesiodistal width of this tooth is greater than its labiolingual width.

A B C D E

9. The mesial and distal marginal ridges are aligned more vertically than horizontally on the lingual surface.

A B C D E

10. The teeth (tooth) develop(s) from four lobes.

A B C D E

11. The teeth (tooth) develop(s) from three lobes.

A B C D E



Helpers 48

B) Choose the correct answer:

1. The permanent maxillary canine erupts:

- a. After the eruption of mandibular canine
- b. Before eruption of both the maxillary premolars
- c. After the eruption of one or both maxillary premolars
- d. Both a and c

2. The labial ridge is prominent in:

- a. Maxillary canine
- b. Mandibular canine
- c. Both
- d. None



3. Among the permanent canines:

- a. The maxillary canine crown is bulkier and broader
- b. Mandibular canine crown is longer and narrower
- c. Crowns of both the canines are of same size
- d. Both a and b

4. Which of the following statements is true about permanent canines?

- a. The cusp is sharp and well developed in maxillary canine
- b. Angle formed by the mesial and distal cusp ridges is wider in mandibular canine
- c. The occlusal portion occupies 1/3rd of the crown length in maxillary canine ; while it occupies 1/5th of the crown length in mandibular canine
- d. All of the above

5. Which of the following statements is true regarding contact areas of permanent canines?

- a. Mesial and distal contact areas are markedly at dissimilar levels in maxillary canine
- b. Contact areas are placed more incisally in mandibular canine
- c. Contact areas are placed at same level in maxillary and mandibular canines
- d. Both a and b

6. When permanent mandibular canine is viewed from labial aspect, the crown is:

- a. Upright on the root base
- b. Tilted mesially on the root base
- c. Tilted distally on the root base
- d. None of the above

7. The cingulum is:

- a. More prominent in maxillary canine
- b. More prominent in mandibular canine
- c. Small in both the canines
- d. Prominent in both the canines

8. Lingual ridge is often seen dividing the lingual fossa into two small fossae in:

- a. Maxillary canine
- b. Mandibular canine
- c. Maxillary central incisor
- d. Maxillary lateral incisor

9. When maxillary canine is viewed proximally, the cusp tip is:

- a. On line with the vertical root axis
- b. Labial to the vertical root axis
- c. Lingual to the vertical root axis
- d. Distal to the vertical root axis

10. When mandibular canine is viewed proximally, the cusp tip is:

- a. On line with the vertical root axis
- b. Labial to the vertical root axis
- c. Lingual to the vertical root axis
- d. Distal to the vertical root axis



Helpers 48

Section 4

Permanent Premolars.

A) For each of the following traits or statements, circle the letter (or letters) of the premolars (if any) that apply.

More than one answer may be correct.

- a. Maxillary first premolar
- b. Maxillary second premolar
- c. Mandibular first premolar
- d. Mandibular second premolar (two-cusp type)
- e. Mandibular second premolar (three-cusp type)



Helpers 48

1. Mesial cusp ridge of the buccal cusp is longer than the distal cusp ridge.

A B C D E

2. Has a nonfunctioning lingual cusp.

A B **C** D E

3. Two premolars that most frequently have a groove crossing the mesial marginal ridge or one groove just lingual to it.

A B **C** D E

4. Has a depression in the cervical one third of the mesial side of the crown and root.

A B C D E

5. Maxillary premolar that has the longer sharper buccal cusp.

A B C D E

6. Largest maxillary premolar.

A B C D E

7. Mandibular premolar with the longest and sharpest buccal cusp.

A B **C** D E

8. Maxillary premolar that is most symmetrical (occlusal view).

A **B** C D E

9. Two premolars without a central groove.

A B **C** D **E**

10. Crowns tipped lingually with respect to the root axis line (proximal view).

A B **C D E**

11. From buccal view, crown is tipped distally from the root axis.

A B **C D E**

12. Mesial marginal ridge is more cervically located than its distal marginal ridge

A B **C** D E

13. Has no transverse ridge.

A B C D **E**

14. Has the longer central groove.

A B C D E

15. Has two major cusps almost the same size and length.

A **B** C D E

16. Has a central fossa.

A B C D **E**



Helpers 48

17. Premolars with only two fossae: both are triangular fossae.

A**B**

C

D

E

18. Has a central fossa and two triangular fossae.

A

B

C

D

E

19. Has a lingual groove.

A

B

C

D

E

Helpers 48

B) Choose the correct answer:

1. Which of the following statements is true regarding the premolar teeth?

- a. Maxillary 1st and 2nd premolars are similar in form
- b. Mandibular 1st and 2nd premolars are similar in form
- c. Mandibular 1st and 2nd premolars are widely different in form
- d. Both a and c**

2. The distal cusp ridge is longer than the mesial cusp ridge in all the premolars except:

- a. Maxillary 1st premolar**
- b. Maxillary 2nd premolar
- c. Mandibular 1st premolar
- d. Mandibular 2nd premolar

3. From proximal aspect, the maxillary premolar crowns appear:

- a. Triangular
- b. Trapezoidal**
- c. Rhomboidal
- d. Ovoid

4. From proximal aspect, the mandibular premolar crowns appear:

- a. Triangular
- b. Trapezoidal
- c. Rhomboidal
- d. Ovoid

5. When viewed proximally, the maxillary premolar crowns are:

- a. Upright on the root base
- b. Tilted buccally on the root base
- c. Tilted lingually on the root base
- d. Tilted mesially on the root base



Helpers 48

6. When viewed proximally, mandibular premolar crowns are:

- a. Upright on the root base
- b. Tilted buccally on the root base
- c. Tilted lingually on the root base
- d. Tilted distally on the root base

7. Compared to maxillary premolars, the buccal and lingual cusp tips of mandibular premolars are:

- a. Wide apart
- b. More nearer
- c. Blunt/flattened
- d. None of above

8. When viewed proximally, buccal cusp tip of maxillary premolars is located:

- a. Buccal to the vertical root axis
- b. Lingual to the vertical root axis
- c. On line with vertical root axis
- d. Both b and c

9. When viewed proximally, the buccal cusp tip of mandibular premolars is:

- a. Buccal to the vertical root axis
- b. Lingual to the vertical root axis
- c. On line with the vertical root axis
- d. Distally to the vertical root axis

10. When viewed proximally, compared to maxillary premolars the mandibular premolars have:

- a. Buccal and lingual crown outlines well within the confines of root base
- b. Buccal crown outlines out of the confines of root base
- c. Lingual crown outline out of the confines of root base
- d. Buccal and lingual crown outlines out of the confines of root base



Helpers 48

Section 5

Permanent Upper Molars.

1. Which three grooves radiate out from the central fossa in a maxillary first molar?

a. Central

b. Distolingual

c. Transverse groove of oblique ridge (when present)

d. Buccal

e. Lingual

2. Which cusp on a maxillary first molar has two ridges: one that forms part of a transverse ridge and the other that forms part of an oblique ridge?

a. Mesiobuccal

b. Mesiolingual

c. Distobuccal

d. Distolingual

e. Cusp of Carabelli



Helpers 48

3. Which cusp is the largest and longest on a maxillary second molar?

a. Mesiobuccal

b. Mesiolingual

c. Distobuccal

d. Distolingual

e. Cusp of Carabelli

4. Which cusp is most likely to be absent on a maxillary second molar?

- a. Mesiobuccal
- b. Mesiolingual
- c. Distobuccal
- d. Distolingual**
- e. Distal

5. When the cusp is absent in question 4 above, which groove(s) would not be present?

- a. Central
- b. Buccal
- c. Distal oblique**
- d. Lingual**

6. Of the four fossae on a maxillary first molar, which is the largest?

- a. Mesial triangular
- b. Distal triangular
- c. Central**
- d. Distal

7. From which view are only two roots visible on a maxillary first molar?

- a. Mesial**
- b. Distal
- c. Buccal
- d. Lingual



Helpers 48

8. Which grooves are likely to radiate out of the mesial triangular fossa on the maxillary first molar?

- a. Mesio Buccal fossa groove
- b. Mesio lingual fossa groove
- c. Mesial marginal ridge groove (when present)
- d. Central
- e. Buccal

9. Which two cusps have the ridges that make up or join to form the oblique ridge on a maxillary molar?

- a. Mesio Buccal
- b. Distobuccal
- c. Mesio lingual
- d. Distolingual
- e. Cusp of Carabelli

10. Which two cusps have a triangular ridge that make up or join to form a transverse (not oblique) ridge on most maxillary molars?

- a. Mesio Buccal
- b. Distobuccal
- c. Mesio lingual
- d. Distolingual
- e. Cusp of Carabelli



Helpers 48

Section 6

Permanent Lower Molars.

1. Which of the following grooves radiate out from the central fossa in a mandibular second molar?

a. Central

b. Mesiobuccal

c. Distobuccal

d. Lingual

e. Buccal

2. Which cusp is the largest and longest on a mandibular second molar?

a. Mesiobuccal

b. Distobuccal

c. Mesiolingual

d. Distolingual

e. Distal

3. Which cusp may be absent on a mandibular first or third molar?

a. Mesiobuccal

b. Distobuccal

c. Mesiolingual

d. Distolingual

e. Distal



Helpers 48

4. When this cusp is absent in question No. 3 above, which groove(s) would not be present?

- a. Buccal
- b. Lingual
- c. Mesibuccal
- d. Distobuccal
- e. Lingual

5. Which fossae are found on a mandibular first molar?

- a. Mesial triangular
- b. Distal triangular
- c. Buccal
- d. Lingual
- e. Central



Helpers 48

6. Which developmental groove connects with the lingual groove running in the same direction on a mandibular second molar?

- a. Mesibuccal
- b. Distobuccal
- c. Buccal
- d. Mesiolingual
- e. Distolingual

7. From which view is only one root visible on a mandibular first molar?

- a. Mesial
- b. Distal
- c. Buccal
- d. Lingual
- e. Apical

8. Which root may occasionally be divided or bifurcated on a mandibular first molar?

- a. Buccal
- b. Lingual
- c. Mesial
- d. Distal
- e. Mesiobuccal

9. Which cusp triangular ridge does not meet to form a transverse ridge on a five-cusp first molar?

- a. Mesiobuccal
- b. Distobuccal
- c. Mesiolingual
- d. Distolingual
- e. Distal

10. Which ridges form the boundaries of the mesial triangular fossa of a mandibular molar?

- a. Triangular ridge of mesiobuccal cusp
- b. Triangular ridge of mesiolingual cusp
- c. Mesial marginal ridge
- d. Buccal cusp ridge of mesiobuccal cusp
- e. Lingual cusp ridge of mesiolingual cusp



Section 7

Permanent Molars.



1. The cusp of Carabelli is a feature of:
 - a. Permanent maxillary 1st molar
 - b. Primary maxillary 1st molar
 - c. Primary maxillary 2nd molar
 - d. Both a and c

2. Which of the following statements is true regarding the molars?
 - a. Maxillary molars have 2 roots; mandibular molars have 3 roots
 - b. Maxillary molars have 3 roots; mandibular molars have 2 roots
 - c. Both maxillary and mandibular molars have 3 roots
 - d. Both maxillary and mandibular molars have 2 roots

3. Maxillary molars have 3 roots. They are:
 - a. Buccal, lingual and distal
 - b. 1 buccal, 2 lingual
 - c. Mesiobuccal, distobuccal and palatal
 - d. 2 mesial and 1 distal

4. The mandibular molars have 2 roots. They are:
 - a. Mesial and distal
 - b. Buccal and lingual
 - c. Buccal and palatal
 - d. Both b and c

5. Oblique ridge is a feature of:

- a. Maxillary premolars
- b. Maxillary molars**
- c. Mandibular premolars
- d. Mandibular molars



6. The following are the features of maxillary molars except:

- a. Cusp of Carabelli on 1st molar
- b. Oblique ridge on occlusal surface
- c. Trifurcated roots
- d. Rectangular occlusal crown outline**

7. Maxillary molar crowns have:

- a. Mesiodistal dimension greater than the buccolingual dimension
- b. Buccolingual dimension greater than the mesiodistal dimension**
- c. Mesiodistal and buccolingual dimensions nearly equal
- d. None of the above

8. Mandibular molar crowns have:

- a. Mesiodistal dimension greater than the buccolingual dimension**
- b. Buccolingual dimension greater than the mesiodistal dimension
- c. Mesiodistal and buccolingual dimensions nearly equal
- d. None of the above

9. Compared to maxillary molars, the buccal surface of mandibular molar crowns:

- a. Is more vertical
- b. Bends buccally from middle third
- c. Bends lingually from cervical third
- d. Bends mesially

10. Compare to mandibular molars, the maxillary molar crowns exhibit:

- a. Less lingual convergence
- b. Greater lingual convergence
- c. No lingual convergence in 1st molar, which may show a buccal taper
- d. Both a and c



Helpers 48

Section 7

Primary Dentition.

1. Which primary teeth have crowns that are wider mesiodistally than they are long inciso- or occluso-cervically?

- a. Maxillary central incisor
- b. Maxillary first molar
- c. Mandibular lateral incisor
- d. Mandibular first molar
- e. Mandibular canine



Helpers 48

2. Which one tooth is adjacent and distal to the primary maxillary second molar in a 7-year-old?

- a. Maxillary first premolar
- b. Maxillary second premolar
- c. Secondary maxillary first molar
- d. Secondary maxillary second molar
- e. Primary maxillary first molar

3. How many teeth should be visible in the mouth of a 3-year-old?

- a. None
- b. 10
- c. 20
- d. 24
- e. 28

4. How many teeth should be present in the mouth of a 13-year-old?

- a. 10
- b. 20
- c. 24
- d. 28
- e. 32

5. Which primary molar most resembles a secondary maxillary right first molar?

- a. Tooth A
- b. Tooth E
- c. Tooth F
- d. Tooth T
- e. Tooth B

6. What would you estimate to be the dental age of a child with the following teeth: all primary maxillary incisors, canines, and molars; secondary mandibular incisors and first molars.

- a. 2 to 4 years
- b. 5 to 7 years
- c. 8 to 9 years
- d. 10 to 11 years
- e. Over 12 years

7. Which teeth (primary or secondary) have the mesial proximal contact positioned more cervically than the distal proximal contact?

- a. Mandibular first premolar
- b. Maxillary first premolar
- c. Primary maxillary canine
- d. Primary mandibular canine
- e. Mandibular second premolar

8. Which teeth (secondary or primary) have the mesial cusp ridge of the facial cusp longer than the distal cusp ridge of the facial cusp?

- a. Mandibular first premolar
- b. Maxillary first premolar
- c. Primary maxillary canine
- d. Primary mandibular canine
- e. Mandibular second premolar



Helpers 48

9. Which succedaneous tooth erupts beneath tooth J?

- a. No. 1
- b. No. 5
- c. No. 10
- d. No. 13
- e. No. 16

10. Which of the following traits can be used to differentiate primary teeth from secondary teeth?

- a. Primary teeth have greater facial cervical bulges.
- b. Primary teeth have relatively thinner and longer roots.
- c. Primary teeth are whiter.
- d. Primary anterior teeth are larger than their successors.
- e. Primary teeth have relatively larger pulps.

11. Which of the following secondary teeth would you expect to be erupted in the average 9- to 10-year-old?

- a. Maxillary lateral incisor
- b. Maxillary central incisor
- c. Mandibular canine
- d. Maxillary canine
- e. Mandibular second molar



Helpers 48

12. Which traits apply to a primary mandibular first molar?

- a. Its roots are resorbed by the eruption of the 6-year mandibular first molar.
- b. It resembles a mandibular 6-year first molar.
- c. It has a prominent buccal cervical bulge.
- d. It has a prominent mesial marginal ridge.
- e. It has a prominent transverse ridge.

Section 8

Occlusion

Answer the following questions with the one best answer based on this sketch of teeth in class I occlusion.

1. In the maximum intercuspal position (MIP), which two teeth occlude with the maxillary first premolar?

- a. Mandibular canine and first premolar
- b. Mandibular first premolar and second premolar**
- c. Mandibular second premolar and first molar

2. In MIP, which two teeth occlude with the mandibular second molar?

- a. Maxillary first premolar and second premolar
- b. Maxillary second premolar and first molar
- c. Maxillary first molar and second molar**

3. Which two teeth would occlude with the incisal edge of the right mandibular lateral incisor during protrusion of the mandible?

- a. Maxillary right central incisor and left central incisor
- b. Maxillary right central incisor and right lateral incisor**
- c. Maxillary right lateral incisor and canine
- d. Maxillary right canine and first premolar



Helpers 48

4. In MIP, with what landmark would the lingual cusp of the maxillary second premolar occlude?

- a. The mesial marginal ridge of the mandibular second premolar
- b. The mesial fossa of the mandibular second premolar
- c. The distal marginal ridge of the mandibular second premolar
- d. The mesial fossa of the mandibular first molar

5. If this person did not have class I occlusion but had class II occlusion (where the mandible was positioned one full tooth distal to its class I position), which tooth or teeth would contact the maxillary second premolar?

- a. Mandibular canine and first premolar
- b. Mandibular first premolar and second premolar
- c. Mandibular second premolar only
- d. Mandibular second premolar and first molar
- e. Mandibular first molar only

6. If this person did not have class I occlusion but had class III occlusion (where the mandible was positioned one full tooth mesial to its class I position), which tooth or teeth would contact the maxillary second premolar?

- a. Mandibular canine and first premolar
- b. Mandibular first premolar and second premolar
- c. Mandibular second premolar only
- d. Mandibular second premolar and first molar
- e. Mandibular first molar only



Helpers 48

Best Wishes

