

mcq

In Dental Anatomy

&

Physiology

CHOOSE THE CORRECT ANSWER:

1. At the age of 9 years, the mandibular permanent first molar has its distal contact with which of the following teeth?
 - a. First premolar.
 - b. No distal contact.
 - c. Second deciduous molar
 - d. Second permanent molar.

2. Which of the following is the last succedaneous tooth to erupt?
 - a. maxillary canine.
 - b. mandibular canine.
 - c. maxillary first premolar
 - d. mandibular third molar.

3. The identifying number of the permanent mandibular left second molar in the two digit numbering system
 - a. 52
 - b. 18
 - c. 17
 - d. 37

4. Which of the following is generally the longest root of maxillary permanent 1st molar?
 - a. distal.
 - b. mesiobuccal.
 - c. palatal.
 - d. distobuccal.

- 5. Which of the following is the most poorly developed cusp in maxillary second permanent molar?**
- Mesiobuccal.
 - Mesia-lingual.
 - Distobuccal.
 - Disto-lingual
- 6. A Y shaped occlusal groove is most likely to be present in:**
- mandibular first premolar.
 - mandibular second premolar.
 - maxillary first premolar.
 - maxillary second premolar
- 7. The roots of which of the following permanent teeth are normally completely formed by the age of 10 years?**
- upper canine.
 - lower second molar.
 - lower second premolar
 - upper central.
- 8. Obtuse occlusal angle in permanent molars usually include:**
- the mesiobuccal in upper 1s molar.
 - the mesiolingual in lower 1s molar.
 - the mesiobuccal in upper 2n molar
 - the distobuccal in upper 1st molar.
- 9. Teeth which usually have transverse ridge on the occlusal surface include:**
- Deciduous lower 1st molars.
 - Permanent 2nd molars.
 - lower 1rt premolars
 - Upper 2nd Premolars.
- 10. The mixed dentition period begins with the emergence of which of the following permanent teeth?**
- First premolar
 - Lower canine
 - Lateral incisors
 - First molar

MCQ IN DENTAL ANATOMY & HISTOLOGY

11. Which of the following teeth develop from five lobes?

- a. upper6&upperE.
c. upper 7 &lower 7.

12. According to Palmer notation system, tooth C| represents?

- Deciduous lower right incisor.
- Deciduous upper right incisor.
- Deciduous upper right canine.
- Deciduous upper left first molar.

13. According to the universal numbering system, tooth #7d refers to:

- a. Deciduous right maxillary first molar.
- b. Deciduous left mandibular second molar.
- c. Permanent right maxillary third molar
- d. Deciduous left maxillary lateral incisor.

14. According to the two digit system of teeth numbering the 21 is:

- a. upper permanent left central incisor.
- b. upper deciduous right lateral incisor.
- c. upper deciduous left central incisor
- d. upper permanent right central incisor.

15. Which of the following is not a representative of a lobe?

- a. cusp.
b. mamelon.
c. tubercle.
d. cingulum.

16. Tooth development starts primarily by formation of:

- a. Tooth germ.
c. dental organ.

17. The portion of tooth that is exposed in the mouth is known as the:

- a Operative crown.
c. Anatomical crown.

18. The term trifurcation refers to the:

- a. Point where three line angles meet
- b. Point at which three occlusal grooves meet
- c. Point at which three roots divide
- d. Point at which three ridges meet

MCQ IN DENTAL ANATOMY & HISTOLOGY

19. The apex of the mesiobuccal root of the permanent upper first molar is:
- In line with the mesiobuccal developmental groove.
 - In line with the distobuccal developmental groove.
 - directly in one line with the mesiobuccal cusp
 - directly below the mesiolingual cusp.
20. The roots of the deciduous molars are more divergent than those of the permanent ones to:
- Provides maximum support & strength to the teeth.
 - Facilitate their resorption.
 - Allow more blood supply to their investing tissues
 - Allow enough room for the permanent successors
21. Which one of the molars is most likely to exhibit a cross occlusal groove pattern?
- #2
 - #14
 - #19
 - #31
22. How many roots does the maxillary second molar have?
- One lingual and two mesial.
 - Two facial and two lingual.
 - One palatal and two facial.
 - Two lingual and one facial.
23. The crown root ratio of deciduous teeth is:
- 1:1
 - 2:3
 - 1:1.5
 - 1:2
24. The transverse groove of the oblique ridge for the upper permanent first molar extends between:
- The mesial and distal triangular fossae.
 - The central fossa and the mesial triangular fossa
 - The distal fossa and the distal triangular fossa
 - The central fossa and the distal fossa
25. The mesial cusp slope in the deciduous canine is:
- Eqixil to the distal cusp slope.
 - Longer than the distal cusp slope.
 - Shoiter than the distal cusp slope
 - Variable.

MCQ IN DENTAL ANATOMY & HISTOLOGY

- 26. Linear incomplete fusion of two developmental lobes of a tooth is known as**
- a. Pit
 - b. Supplemental groove.
 - c. Developmental groove
 - d. Fissure.
- 27. There are two lingual fossae on the lingual surface of:**
- a. Upper lateral incisor.
 - b. Maxillary canine.
 - c. Mandibular canine,
 - d. Lower lateral incisor.
- 28. The location of the mental foramen at three years is:**
- a. Midway between the upper & lower borders.
 - b. Closer to the upper border.
 - c. Closer to the lower border
 - d. Below the crypt of lower D.
- 29. In old age the condyle becomes inclined backward due to:**
- a. Increase in the mandibular angle measurement
 - b. Posterior muscle traction
 - c. Elongation of the coronoid process.
 - d. Retraction of the mandible during mastication.
- 30. One of the direct factors protecting the Periodontium is**
- a. Crown form
 - b. Self-cleansing ability of the tooth
 - c. Proximal contact relation
 - d. Angulation of tooth surfaces.
- 31. All the following are considered as types of embrasures except:**
- a. Occlusal.
 - b. Cervical.
 - c. Facial
 - d. Lingual.

MCQ IN DENTAL ANATOMY & HISTOLOGY

32. The mesio-lingual cusp of upper first permanent molar strikes its antagonist at:
- Central fossa
 - Mesial marginal ridge.
 - Distal marginal ridge
 - Distal cusp.
33. Which of the following is the deciduous dental formula of man?
- $12/2 \text{ CI} 1 \text{ M} 2/2 = 10 \times 2 = 20$.
 - $12/2 \text{ CI} 1 \text{ M} 3/3 = 12 \times 2 = 24$.
 - $12/2 \text{ CI} 1 \text{ PM} 2/2 \text{ M} 3/3 = 16 \times 2 = 32$
 - $12/2 \text{ CI} 1 \text{ PM} 1/1 \text{ M} 2/2 = 12 \times 2 = 24$.
34. All the following teeth have no predecessors except:
- The maxillary and mandibular premolars.
 - The permanent maxillary and mandibular first molars.
 - The permanent maxillary and mandibular second molars
 - The permanent maxillary and mandibular third molars.
35. All the following are nonsuccedaneous, except:
- the maxillary and mandibular premolars.
 - the permanent maxillary and mandibular first molars.
 - the permanent maxillary and mandibular second molars
 - the permanent maxillary and mandibular third molar.
36. The successor of the upper first deciduous molar is:
- Upper first permanent molar.
 - Upper second premolar,
 - Upper first premolar.
 - Upper second permanent molar.
37. The cusp of the canine develops from:
- Mesial lobe.
 - Mid die lobe.
 - Distil lobe.
 - Lingual lobe
38. Which of the following is not true regarding surfaces of the teeth?
- Mesial surfaces are those towards the median line of the face,
 - Distal surfaces are those most distant from the median line of the face.
 - The surface that is towards the lip is the lingual surface.
 - The surface that is towards the cheek on posterior teeth and the lips on anterior teeth is the fecial surface.

MCQ IN DENTAL ANATOMY & HISTOLOGY

39. Which of the following teeth may have two roots?

- a. Maxillary central incisor.
- b. Maxillary lateral incisor.
- c. Mandibular canine
- d. Maxillary canine.

40. The proximal surfaces of mandibular premolars are:

- a. Trapezoidal.
- b. Triangular.
- c. Rhomboidal
- d. Diamond

41. All of the following are point angles except:

- a. Dislo-bucco-occlusal.
- b. Mesio-linguo-occlusal.
- c. Mesio-bucco-occlusal
- d. Disto-mesio-occlusal.

42. The curve of Wilson in the maxillary segment is:

- a. Concave
- b. Convex,
- c. Concave - convex.
- d. Sphere.

43. Normal intact healthy interproximal spaces are filled with:

- a. Free gingiva
- b. Periodontal fibers.
- c. Inter-dental papillae.
- d. Attached gingiva.

44. The occlusal surface of the upper first premolar is bounded by how many lines:

- a. two.
- b. four.
- c. five.
- d. six.

45. According to the two digit system, 81 & 53 refer to:

- a. Upper right central incisor and lower left permanent 3rd molar.
- b. Upper right 2nd premolar and lower right deciduous canine.
- c. Lower right deciduous central incisor and upper right deciduous canine.
- d. Upper left permanent central incisor and upper right deciduous canine.

MCQ IN DENTAL ANATOMY & HISTOLOGY

53. Which of the following teeth is **MOST LIKELY** to have a lingual pit according to the universal numbering system?
a. #23 b. #11 c. #9 d. #7
54. On the crown of the maxillary canine, **WHAT LOBE** includes the cusp tip?
a. mesial facial lobe, b. middle facial lobe,
c. distal facial lobe, d. lingual lobe.
55. How many cingula are there in the permanent dentition?
a. 12 b. 6 c. 8 d. 4
56. The linear elevations that form the lateral boundaries of the occlusal surfaces of posterior teeth and the lingual surfaces of anterior teeth are called:
a. Triangular ridges. b. Cervical ridges,
c. Lingual ridges. d. Marginal ridges.
57. What is the name of the feature on the lingual surface of maxillary canine that extends from the cusp tip on down to the cingulum of the tooth.
a. mesial marginal ridge,
b. cervical ridge,
c. lingual ridge
d. incisal ridge.
58. The shape of the central & distal fossae of the occlusal surface of the 6 respectively is:
a. Circular & linear. b. Triangular & linear,
c. Rounded & triangle. d. Triangular & circular.
59. The occlusal surface of 6 is rhomboidal in shape with:
a. Acute mesiobuccal & distolingual angles.
b. Obtuse mesiobuccal & distolingual angles.
c. Acute distobuccal & distolingual angles.
d. Obtuse distobuccal & acute mesiolingual angles.
60. The height of contour of the facial and lingual surfaces of the anterior teeth occurs in the:
a. mesial third. b. incisal third,
c. middle third d. cervical third

MCQ IN DENTAL ANATOMY & HISTOLOGY

- 61. When viewed from the occlusal aspect, which of the following is a difference in the crown of 5 as compared to 4 :**
- A more pointed outline.
 - A shorter central groove.
 - A longer central groove
 - Thinner marginal ridges.
- 62. The maxillary canine can be distinguished from the mandibular canine by which of the following characteristics:**
- The mandibular canine has a more prominent cingulum.
 - In the mandibular canine, the cusp slopes are long.
 - In the mandibular canine, the cusp tip is located labially to the center of the tooth,
 - The mandibular canine has less prominent cingulum.
- 63. A mesio-lingual developmental groove is found on the surface of :**
- Lower second premolar (2 cusp-type).
 - Lower second premolar (3 cusp type).
 - Lower first premolar
 - Upper first premolar.
- 64. The mesiobuccal cervical ridge is most marked & considered a characteristic feature of:**
- Deciduous & permanent upper 1st molars.
 - Deciduous & permanent lower 1st molars.
 - Deciduous upper & lower 1st molars
 - Permanent upper & lower 1st molars.
- 65. The tooth follicle gives:**
- The dental lamina, periodontal ligament and cementum.
 - The dental lamina, alveolar bone and cementum
 - Enamel, dentin and cementum.
 - Cementum, periodontal ligament and alveolar bone.
- 66. The sequence of eruption of deciduous teeth is as follows:**
- A,B,C,D,E.
 - A,B,D,C,E.
 - B,E,C,D.
 - A,C,B,D.

MCQ IN DENTAL ANATOMY & HISTOLOGY

- 67. The mesial marginal ridge is so well developed that it resembles a cusp in:**
- a. The deciduous lower first molar.
 - b. The deciduous lower second molar.
 - c. The permanent upper first molar.
 - d. The deciduous upper second molar.
- 68. The enamel of the deciduous teeth is more easily worn down because:**
- a. It is white- in colour than that of the permanent teeth.
 - b. It is thinner than that of the permanent teeth.
 - c. It is more opaque than that of permanent teeth.
 - d. It is less mineralized than that of permanent teeth.
- 69. Eruption of the lower 5 occurs at:**
- a. 9-10 years.
 - b. 11-12 years,
 - c. 12-13 years.
 - d. 8-9 years.
- 70. Appearance of the enamel organ of the deciduous upper first molar occurs at:**
- a. 4 weeks I.U.L.
 - b. 4 months I.U.L.
 - c. 8 weeks I.U.L.
 - d. 5 months I.U.L.
- 71. Eruption of the deciduous upper and lower second molars occur respectively at the age of:**
- a. 24 and 20 months.
 - b. 24 and 22 months.
 - c. 20 and 18 months.
 - d. Two years and 22 months.
- 72. Beginning of calcification of the permanent second molars occurs at:**
- a. At birth.
 - b. 1 year,
 - c. 2- 3 years.
 - d. 4-5 years.
- 73. Beginning of root resorption of the deciduous canines takes place at:**
- a. 5 years.
 - b. 6 years,
 - c. 7 years.
 - d. 8 years.

MCQ IN DENTAL ANATOMY & HISTOLOGY

74. During the fetal period, the two halves of the mandible are united just by fibrous tissue at the symphysis menti which get completely ossified:

- a. At birth.
- b. Six months after birth.
- c. By the end of the first year.
- d. By the end of the second year.

75. In old age the mandibular angle is:

- a. 140° .
- b. 150° .
- c. 160° .
- d. 170° .

76. Marked reduction in the height of the body of the mandible in old age places the mental foramen & the mandibular canal:

- a. Near the lower border of the mandible.
- b. Close to the upper border.
- c. Midway between upper & lower border.
- d. In the middle 3rd of the body of the mandible.

77. In the adult period the mandibular angle measures :

- a. $13(1-140^\circ)$.
- b. $150-160^\circ$.
- c. $11(1-130^\circ)$.
- d. $160-170^\circ$.

78. What is the geometric form of the anterior teeth when they are viewed from the proximal (mesial or distal) aspect?

- a. rhomboid
- b. trapezoid
- c. square
- d. triangular

79. The trapezoid shape of the facial and lingual surfaces of the teeth provides:

- a. Narrow occlusal surfaces to be easily forced through the food.
- b. Self cleansing ability of the teeth.
- c. Positive contact between the neighboring teeth,
- d. Wide escapement spaces.

80. The trapezoid form of the proximal surfaces of upper posterior teeth provides:

- a. Wide interproximal spaces.
- b. Proper embrasures or spillways.
- c. Narrow occlusal surfaces leading to decrease in the forces of mastication applied on the teeth.
- d. Positive contact between the teeth.

MCQ IN DENTAL ANATOMY & HISTOLOGY

81. The rhomboidal shape of the proximal surfaces of the lower posterior teeth provides the following except:

- a. Making the axes of the crown and roots of the teeth of the arches parallel to each others.
- b. Lingual inclination of the crown bringing the cusps into proper occlusion with the cusps of their maxillary opponents.
- c. Prevents clashing between the-cusps of the lower teeth with the cusps of their maxillary opponents.
- d. Create proper compensating curvatures of the dental arches.

82. The interproximal spaces are triangular in shape, the base, side walls and apex of the triangle respectively is formed of:

- a. Alveolar process, proximal surfaces of adjacent teeth and contact areas.
- b. Contact areas, proximal surfaces of adjacent teeth and alveolar process.
- c. Cervical lines of adjacent teeth, contact areas and alveolar process
- d. Alveolar process, facial surfaces of adjacent teeth and contact areas.

83. proximal contact areas are:

- a. Narrow in the posterior teeth.
- b. Are centered buccolingually in the posterior teeth.
- c. Situated more to the lingual in anterior teeth.
- d. Situated more to the buccal in posterior teeth.

84. The divergence of two proximal surfaces from the area of contact form spaces facially, occlusally or incisally, and lingually, known as:

- a. into proximal spaces.
- b. gingival attachments.
- c. contact areas
- d. embrasures.

85. The lingual embrasure of anterior teeth is:

- a. Equal o the labial embrasure.
- b. Smaller than the labial embrasure.
- c. Larger than the labial embrasure,
- d. variable in size.

MCQ IN DENTAL ANATOMY & HISTOLOGY

86. Underdeveloped facial and lingual contours of the teeth lead to:

- a. Gingival recession.
- b. Injury to the attachment epithelium
- c. Excessive frictional irritation of the gingiva
- d. All of the above.

87. The curvature of the cervical line is important for the health of the periodontium because:

- a. It is greater in the anterior than in the posterior teeth.
- b. It is greater mesially than distally.
- c. It tends to be straight in the distal surface of some of the molars.
- d. It is the site of attachment of the gingiva to the tooth by the attachment epithelium.

88. The permanent dental arch is considered physiologically completely developed by the end of:

- a. The fifth phase.
- b. The sixth phase,
- c. The fourth phase.
- d. The third phase

89. In an ideal occlusion, the lower canine contacts what teeth in the upper arch?

- a. the upper canine and the upper lateral incisor.
- b. the upper canine and the upper first premolar.
- c. just the upper canine.
- d. the upper central incisor and the upper lateral incisor.

90. Curve of Wilson is deeper in the:

- a. Molar segment
- b. Premolar segment,
- c. Canine segment
- d. Incisors segment

91. The permanent 1st molar is the key of occlusion because of all the following reasons except:

- a. It is the last permanent tooth to develop and erupt
- b. It is the largest permanent tooth.
- c. Its eruption is guided to its correct position just distal to the deciduous molars,
- d. Its eruption can never be disturbed since it has no predecessor.

MCQ IN DENTAL ANATOMY & HISTOLOGY

92. The horizontal overlap (over jet) is very important for:

- a. Stability of the dental arches.
- b. Proper oral hygiene.
- c. Protection of the lips and cheeks against biting during mastication
- d. Providing room for the proper function of the masseter muscle.

93. Each tooth in one dental arch contacts two teeth in the opposing dental arch in centric occlusion except lower 1 and upper 8 This relation is important because:- .

- a. It causes elongation or overeruption
- b. It causes displacement of teeth through lack of antagonism
- c. It causes increase in forces over larger number of teeth.
- d. It prevents elongation or overeruption

94. The parabolic curve is formed of ;

- a. Three segments that overlap each other.
- b. Three segments that meet at a point
- c. Two segments that meet at the midline.
- d. Two segments that overlap each other.

95. The overhanging of the maxillary teeth when they are in centric occlusion with the mandibular teeth is called:

- a. Overjet
- b. Overbite.
- c. Open bite.
- d. Deep bite.

96. Bon will described the mandibular dental arch to be adapted to a 4 inches equilateral triangle, its angles are located as follows:

- a. The angles of the base are located at the anterior border of the condyles while the apex at the mesial contact areas of the lower central incisors.
- b. The angles of the base at the centers of the condyles while the apex at the mesial contact areas of the lower central incisors.
- c. The angles of the base at the center of the condyles and the apex at the cingulum of the lower central incisors.
- d. The angles of the base at the coronoid while the apex at the mesial contact areas of the lower central incisors,

MCQ IN DENTAL ANATOMY & HISTOLOGY

97. In centric occlusion the contact relation between cusp and fossa occurs when:

- a. The buccal cusps of upper molars rest in fossae of the occlusal surfaces of lower molars.
- b. Lingual cusps of lower molars rest, in fossae of the occlusal surfaces of upper molars.
- c. Lingual cusps of upper molars rest in fossae of the occlusal surfaces of lower molars.
- d. Buccal cusps of the lower molars contact the marginal ridges of upper molars.

98. Normal intact interproximal Spaces are:

- a. Spherical.
- b. Triangular.
- c. Rectangular.
- d. Square.

99. Which of the following is the sequence of eruption of permanent teeth normally seen in the maxilla?

- a. 1,6,2,4,3,5,7,8.
- b. 1,6,2,3,4,5,7,8.
- c. 6,1,2,4,3,5,7,8.
- d. 6,1,2,3,4,5,7,8.

100. Curve of Spee:

- a. Is located within coronal plane.
- b. Is located within sagittal plane.
- c. Is located within coronal and sagittal plane
- d. Ends at the posterior border of the condyle.

101. Which of the following is not considered one of tooth tissue?

- a- Enamel
- b-Dentin
- c-Pulp
- d- Periodontal ligament

102. Using the Palmer notation system, lower right first deciduous molar is represented by:

- a. DI
- b. D
- c. D|
- d. E

MCQ IN DENTAL ANATOMY & HISTOLOGY

103. Which of the following is not true regarding surfaces of the teeth?

- a. Mesial surfaces are those toward the median line of the face
- b. Distal surfaces are those most distant from the median line of the face
- c. The lingual surfaces are those towards the lip
- d. The buccal surfaces on posterior teeth are those towards the cheek.

104. All permanent teeth have lingual convergence except:

- a- Upper 6₁, upper 7₁ (4 cusp type) and lower 5 (3 cusp type),
- b- Lower molars and upper 5.
- c- Upper 4₁ and lower 4₁
- d- Lower 4₁ and lower 5 (2 cusp type).

105. Which of the following deciduous tooth is generally the last to erupt ?

- a- Central incisor.
- b- First molar,
- c- Canines.
- d- Second molar.

106. Which of the following are not succedaneous teeth?

- a- Permanent incisors.
- b- Premolars,
- c- Permanent molars.
- d- Permanent canines.

107. The occlusal surface of maxillary deciduous first molar J3 may resemble which one of the following maxillary permanent teeth?

- a- First premolar.
- b- Second premolar,
- c. First molar.
- d- Second molar.

108. Which of the following is true regarding the pulp cavity of the deciduous teeth?

- a- The pulp horns are both low and small,
- b- The pulp cavity is proportionally large,
- c- The root canals of deciduous molars are markedly narrow,
- d- The pulp chambers are very small.

109. The mixed dentition period begins with the emergence of which of the following permanent teeth?

- a- First molars.
- b- Lower canine,
- c- Lateral incisors.
- d- First premolars.

MCQ IN DENTAL ANATOMY & HISTOLOGY

110. The maxillary permanent lateral incisor is smaller than the maxillary permanent central incisor in all aspects except:
- a- Crown length.
 - b- Crown width mesio-distally.
 - c- Root length.
 - d- Crown thickness labio-lingually
111. The largest cusp found on the crown of lower first permanent molar (H5) is:
- a- Mesio buccal cusp.
 - b- Disto- buccal cusp,
 - c- Mesio- lingual cusp.
 - d- Disto- lingual cusp.
112. Which of the following is true regarding the root portion of mandibular permanent canine?
- a- It is conical in shape.
 - b- It is narrow labio-lingually.
 - c- A bifurcated root is not uncommon.
 - d- If bifurcated, the roots are mesial and distal.
113. The maxillary first premolar (4) will typically demonstrate which of the following pulpal configurations?
- a- A single root with one root canal,
 - b- A double root with two root canals,
 - c- A double root with three root canals,
 - d- One root with three root canals.
114. The root of the mandibular second premolar (5) is completed between:
- a- 9 to 10 years of age.
 - b- 10 to 12 years of age.
 - c- 13 to 15 years of age.
 - d- 16 to 17 years of age.
115. The most common form of the mandibular second premolar is:
- a- Two cusps, one buccal and one lingual
 - b- Three cusps, buccal, mesio-lingual and disto-lingual.
 - c- Three cusps two buccal and one lingual
 - d- Two cusps, one mesial and one distal
116. Which of the following teeth develop from five lobes?
- a. Upper first permanent molar & upper second deciduous molar (6 & E).
 - b. Upper first premolar & mandibular second premolar (4 & 5).
 - c. Upper & lower permanent second molar (7 & 7).
 - d. Lower first permanent molar, lower second deciduous molar & Iowa* second premolar three cusps type (6, E, 5).

117. The oblique ridge of deciduous maxillary second molar connects the:
- a- Mesio-buccal and mesio-lingual cusps,
 - b- Disto-buccal and disto-lingual cusps,
 - c- Disto-buccal and mesio-lingual cusps,
 - d- Mesio-buccal and disto-lingual cusps.
118. According to the Palmer notation system, tooth 1C represents:
- a- Deciduous lower right lateral incisor,
 - b- Deciduous upper right lateral incisor
 - c- Deciduous upper right canine,
 - d- Deciduous upper left canine.
119. Which one of the following deciduous teeth has a crown unlike that of any permanent tooth?
- a- Maxillary first molar.
 - b- Maxillary second molar,
 - c- Mandibular first molar.
 - d- Mandibular second molar.
120. The buccal cusp of upper first premolar (4₁) has:
- a- Mesial and distal cusp slopes of equal length,
 - b- Mesial cusp slope longer than the distal cusp slope,
 - c- Distal cusp slope longer than the mesial cusp slope,
 - d- Cusp tip located toward the mesial aspect of the crown.
121. Using the universal numbering system, tooth # 1d represents:
- a- Deciduous upper right second molar,
 - b- Deciduous lower right second molar,
 - c- Deciduous upper right first molar,
 - d- Deciduous lower left first molar
122. The transverse ridge results from the union of:
- a- Two marginal ridges.
 - b- Two triangular ridges,
 - c- Two cervical ridges.
 - d- A buccal and a lingual ridges.
123. A prominent concavity on the mesial surface of the crown is found in:
- a- Maxillary first premolar,
 - b- Maxillary second premolar
 - c- Mandibular first premolar,
 - d- Mandibular second premolar.

MCQ IN DENTAL ANATOMY & HISTOLOGY

124. Which of the following teeth has usually two root canals in its mesial root?

- a- Upper second permanent molar
- b- Upper first premolar
- c- Lower permanent canine
- d- Lower first permanent molar

125. In normal centric occlusion, the mesiobuccal cusp off the permanent mandibular first molar contacts the :

- a- Distal triangular fossa of the maxillary first premolar (4).
- b- Distal marginal ridge of the maxillary second premolar (5). and the mesial marginal ridge of maxillary first molar (6).
- c- Mesial triangular fossa of the maxillary second premolar (5).
- d- Central sulcus of the maxillary first molar (6).

126. Lower first premolar is characterized by the presence of:

- a- Mesial developmental groove.
- b- Mesial marginal developmental groove.
- c- Mesiolingual developmental groove
- d- Mesiobuccal groove.

127. Contact areas between teeth are significant in:

- a- Providing self cleaning action for teeth
- b- Allowing gingival massage,
- c- Preventing food impaction,
- d- Providing enough space for interdental papilla

128. According to the universal numbering system, the permanent tooth which is located second from the midline in the mandibular left quadrant is:

- | | |
|-------|-------|
| a-#7 | b-#23 |
| c-#25 | d-#26 |

129. The geometric outline of the facial and lingual surfaces of all teeth provides.

- | | |
|---------------------------------|--------------------------------|
| a- Proper occlusion | b- Proper deflection of food, |
| c- Proper contact between teeth | d- Proper angulations of teeth |

130. The roots of deciduous tooth are widely divergent to:

- a- Provide maximum support to the tooth
- b- Facilitate their resorption
- c- Allow more blood supply to their supporting tissue,
- d- Allow enough room for their successors.

MCQ IN DENTAL ANATOMY & HISTOLOGY

131. The maxillary molar primary cusp triangle is formed of:

- a- The two buccal cusps and the mesio- lingual cusp,
- b- The two mesial cusps and the disto- lingual cusp,
- c- The two distal cusps and the mesio-lingual cusp,
- d- The two lingual cusps and the disto- buccal cusp.

132. Which of the following growing teeth are present in their crypts at birth?

- a- The occlusal surface of D and the mesiolingual cusp of Iowa* 6.
- b- The incisal edge of A and 1/2 of the crown of C.
- c- The completed crown of A & B.
- d- The occlusal surface of E and 1A of the crown of D.

133. In centric occlusion, each tooth is in contact with two antagonists of the opposing dental arch except.

- a- 1 and 8
- b- 1 and 8
- c- J and JL
- d- 1 and 8.

134. In adult age the mandibular angle measurement is about.

- a-140°.
- b-120°.
- c-170°.
- d.160°.

135. In old age and after extraction of teeth, the mental foramen and mandibular canal become:

- a- Near to the lower border.
- b- Midway between the upper and lower border.
- c- Close to the upper border.
- d- Obliterated by bone deposition.

136. Which of the following is an indirect factor in protecting the prelodontium?

- a- Contact areas.
- b- Embrasures,
- c- Interproximal space.
- d- Crown and root form

137. During centric occlusion the labio-incisal ridge of lower permanent lateral incisor strikes:

- a- Upper central incisor (1) only.
- b- Upper central & lateral incisors (1& 2).
- c- Upper lateral incisor (2) only,
- d- Upper lateral incisor & upper canine (2 & 3).

138. At birth the two halves of the mandible are united to each other in the median plane by:

- a- Compact bone,
- b- Cartilage,
- c- Spongy bone
- d- Fibrous tissue

139. At birth, mandibular canal runs:

- a- Near the posterior border of the ramus.
- b- Near the lower border of the body of the mandible.
- c- Midway between the upper and lower borders of the body of the mandible.
- d- Near the upper border of the body of the mandible.

140. At birth the mandibular angle measures:

- a. 120
- b. 110
- c. 140
- d. 170

141. The condyle of the mandible is located at a lower level to the coronoid process;

- a- In old age after loss of teeth,
- b- In old age before loss of teeth
- c- At adult
- d- After eruption of deciduous teeth.

142. The geometric outline form of anterior teeth proximalBSy is:

- a- Rhomboidal.
- b- Trapezoidal.
- c- Square.
- d- Triangular.

143. The rhomboidal shape of the proximal surfaces of lower posterior teeth provides the following except:

- a. Making the axis of the crown and root of teeth of the two arches parallel to each others.
- b. Creating proper compensating curvature of the dental arch,
- c. Lingual inclination of the crown for proper occlusion,
- d- Prevention of clashing between upper and lower cusps.

144. Palatal pit is frequently present in the depth of the lingual fossa of:

- a- Upper permanent central incisor,
- b- Lower permanent lateral incisor,
- c- Upper permanent lateral incisor
- d- Lower permanent central incisor.

MCQ IN DENTAL ANATOMY & HISTOLOGY

145. According to the two digit system of teeth numbering the 21

is:

- a- Maxillary permanent left central incisor
- b- Maxillary deciduous right lateral incisor,
- c- Maxillary deciduous left central incisor,
- d- Maxillary permanent right central incisor.

**146. The geometric outline of lower permanent first molar (6)
from mesial surface is:**

- a. Hexagonal.
- b. Rhomboidal and wider occlusally
- c. Heart shape.
- d. Rhomboidal with occlusal constriction.

147. Calcification of upper permanent lateral incisor(2) begins:

- a. At birth,
- b. Before birth.
- c. At 10-12 months,
- d. At 4 months.

148. Two root canals may be found in:

- a- Upper central incisor,
- b- Upper lateral incisor,
- c- Upper canine,
- d- Lower canine.

Good Luck

Answer Key Dental Anatomy & Physiology

1-B	26-D	51-C	76-B	101-D	126-C
2-D	27-B	52-D	77-C	102-C	127-C
3-D	28-D	53-D	78-D	103-C	128-B
4-C	29-B	54-B	79-C	104-A	129-C
5-D	30-C	55-A	80-C	105-D	130-D
6-B	31-B	56-D	81-D	106-C	131-A,
7-D	32-A	57-C	82-A	107-A	132-A
8-D	33-A	58-B	83-D	108-B	133-A
9-C	34-A	59-A	84-D	109-A	134-B
10-D	35-A	60-D	85-A	110-C	135-C
11-D	36-C	61-B	86-D	111-A	136-D
12-C	37-B	62-D	87-D	112-C	137-B
13-D	38-C	63-C	88-C	113-B	138-D
14-A	39-C	64-C	89-A	114-C	139-B
15-C	40-C	65-D	90-A	115-B	140-D
16-B	41-D	66-B	91-A	116-D	141-A
17-D	42-B	67-A	92-C	117-C	142-D
18-C	43-C	68-D	93-D	118-D	143-B
19-C	44-D	69-B	94-A	119-C	144-C
20-D	45-C	70-B	95-D	120-B	145-A
21-D	46-D	71-A	96-B	121-A	146-D
22-C	47-A	72-C	97-C	122-B	147-C
23-B	48-C	73-D	98-B	123-A	148-D
24-D	49-B	74-C	99-B	124-D	
25-B	50-B	75-A	100-C	125-B	