





Incisors Revision

1

1) Labial aspect	Upper central incisor	Upper lateral incisor	Lower central incisor	Lower lateral incisor
N.B	The largest (widest and broadest) crown in all anterior teeth -it's the most prominent in mouth.	- it's smaller than the central incisor except in root length . -This tooth shows great variation + may show lingual pit + it may be peg shape due to absence of middle lobe + it may be completely absent .	-The smallest tooth in the permanent dentition. -It's the most symmetrical tooth in the permanent dentition	- it's very similar to the lower central incisor. - it is slightly larger than the lower central incisor.
GEOL	Trapezoid with the smallest uneven side cervically and the largest side incisally		Same + They have narrow neck giving the crown fan shaped appearance .	
All outlines	-Mesial: sl. Convex -Distal: more convex -Incisal: newly erupted incisors show 3 mamelons , these mamelons soon undergo attrition , giving a regular and straight incisal edge -Mamelons in upper central is well developed than in the upper lateral incisor -Cervical line : convex root wise		-Mesial: straight -Distal: straight -Incisal: straight and it's nearly a right angle to the long axis of the tooth -Mamelons: in lower incisors they are less developed than the upper incisors -Cervical line : convex root wise	The same but with small differences :- The lateral incisor has more fan shaped appearance than central incisor -the mesial outline is often longer than the distal outline , this is due to the incisal ridge slopes downward in a distal direction -Therefore it is not symmetrical as the lower central incisor
Mesial Contact area	in the incisal 1/3 near the mesio-incisal angle	Near the junction between incisal and middle one third	MCA and DCA are in the same level and exist in the incisal 1/3	near the junction between the incisal and middle third.
Distal Contact area	more cervical , Near the junction between incisal and middle one third	More cervical		DCA: more cervically.
Mesio-Incisal angle	sharp	rounded	sharp	sharp
Disto-Incisal angle	rounded	more rounded so incisal outline is rounded	sharp	rounded
1)Elevation	-The labial surface is smoothly convex with the maximum convexity at the cervical 1/3 (cervical ridge). -Mamelons : well developed in upper central and less developed in lateral incisor .		-The labial surface is smoothly convex with the maximum convexity at the cervical 1/3 (cervical ridge) -Mamelons : they are prominent in in newly erupted teeth .	
2)Depressions	-2 shallow developmental grooves which divide the labial surface into 3 lobes. -The 2 developmental grooves is well developed in upper central and less developed in upper lateral incisor .		-2 shallow developmental grooves which divide the labial surface into 3 lobes -The 2 developmental grooves is well developed in newly erupted teeth	
ROOT	Single centralized root , cone shaped , with blunt apex. So This tooth can be extracted by rotation	Single root curved distally , tapers and end with a pointed apex.	Single centralized root , the labial surface is narrow and convex. The apex is pointed and curved distally	Single root curved distally , but slightly longer than the lower central incisor



2) Lingual aspect	Upper central incisor	Upper lateral incisor	Lower central incisor	Lower lateral incisor
N.B	Both crown and root are narrower lingually than labially due to (lingual convergence)			
Elevation	-Cingulum : is well developed -Mesial and distal marginal ridges : well developed that extend from the cingulum to the incisal ridge 		-Cingulum : is less developed -Mesial and distal marginal ridges: less developed than upper central incisor. -The incisal ridge is narrow and rounded or flat 	-Cingulum: less developed and shifted distally -Mesial and distal marginal ridges : less developed , however they still more prominent in lower central incisor. 
Depressions	-the lingual fossa is a large concavity lies between the marginal ridges , the cingulum and the incisal ridge . 	- the lingual fossa is more concave and circumscribed - palatal pit may be found and enclosed by the cingulum , and prone to decay . 	-the lingual fossa is shallow	
ROOT	Root is narrower than the labial surface due to lingual convergence			
3) Mesial aspect	Upper central incisor	Upper lateral incisor	Lower central incisor	Lower lateral incisor
GEOL	Triangular in shape , the base towards the cervical line and the apex towards the incisal edge .			
The outline form	-Labial outline : convex in the cervical third (cervical ridge) And is sl. Convex or nearl flat in the middle and the incisal thirds . - Lingual outline : convex in the cervical third (cingulum) then concave in the middle (fossa) and sl. Convex in the incisal ridge . - Cervical line : curves incisally to a marked degree .			
N.B.	The incisal ridge of the crown is on a line with the center of the root . 		The incisal ridge of the crown is lingual to the center of the root , due to lingual inclination . 	
Contact areas	Centralized (labio-lingual) and in the incisal third near the mesio-incisal angle			
Root	It's cone shaped and apex is centralized and blunt			
4) Distal aspect	It differs from mesial surface in :- - the crown is somewhat thicker toward the incisal third . - the curvature of the cervical line is less distally than mesially -the distal contact areas is still centralized but is located more cervically -the surface of the root is convex and does not have a depression .			

Incisal aspect	Upper central incisor	Upper lateral incisor	Lower central incisor	Lower lateral incisor
GEOL	Triangular , the base towards the labial surface and the apex towards lingual (lingual convergence)``		Diamond , and the crown converges lingually (lingual convergence)	
	Width> thickness		Thickness > width	
N.B	-the labial surface is relatively broad and flat -the incisal edge is centrally situated in relation to thickness . -the lingual fossa is seen as broad concavity between the two marginal ridges and the cingulum -there is lingual convergency 	- the incisal aspect is similar to that of upper central incisor except in size . -the incisal outline is more rounded labially and lingually than that of upper central incisor . -when palatal pit is present , it's located in the depth of lingual fossa 	the labial and the lingual outlines is convex -the incisal edge is lingually situated (due to lingual inclination) and almost at right angles to a line bisecting the crown labio-lingually which is the characteristic feature of lower central incisor. -the lingual fossa is seen as shallow concavity between the two marginal ridges and the cingulum 	The most important feature is the appearance of the crown to be slightly twisted distolingually on its root . this is to allow the incisive ridge to follow the curvature of the dental arch . The incisal ridge is not at a right angle to a line bisecting the crown labio-lingually 
Pulp cavity	Upper central incisor	Upper Lateral incisor	Lower central incisor	Lower lateral incisor
	Mesio-distally : trapezoid while width >thickness + 3 pulp horns confirm to the mamelons . Labio-lingually: rough;y triangular with pointed incisal edge. 	-Similar but smaller than upper central incisor . -It has a single root canal . -There is no 3 pulp horns as there are less prominent mamelons in this tooth	t's similar to that of upper central incisor as it has a single root canal and 3 pulp horns (in newly erupted teeth) However the pulp chamber is narrower mesiodistally than labiolingually	