

Complete the following

1. In maxillary central permanent incisors a deep lingual fossa and exaggerated marginal ridges produce a
2. In the maxillary central incisor the.....incisal angle is , while the....is round.
3. The anterior tooth which is a subject of an enormous degree of variation is.....
4. The presence of a lateral incisor..... In the depth of the palatal fossa is-common in the maxillary lateral incisor.
5. The crown of the lower lateral incisor is twisted distolingually slightly on its root in order to allow the.....ridge to follow the
6. The mesial slope of the maxillary canine is than the distal slope.
7. In the maxillary canine, the depression on the.....surface of the root is more pronounced than that on the surface.
8. of root to a labial and lingual root is the most conspicuous variation in the lower canine.
9. The lingual surface of all teeth is smaller in size than the labial due to the.....

TRUE OR FALSE:

1. In the Mandibular central incisor the-Mesio-incisal angle is sharp while the Disto-incisal angle is rounded.
2. In the Mandibular central incisor the cingulum is centered while in the mandibular lateral incisor the cingulum is deviated distal to the center of the lingual surface.
3. Proximal aspects of anteriors have triangular outline with the apex cervically and the base incisally
4. The incisal ridge of maxillary incisors is perpendicular to the long axis while in the mandibular incisors it is inclined lingually.
5. In the incisal aspect, the width is greater than the thickness in the maxillary central incisor while the thickness is greater than the width in the maxillary lateral incisor.
6. From the labial aspect the crown of the upper canine is long, narrow&thin while that of the lower canine is short, wide thick.
7. The geometric outline of the incisal aspects of both upper & lower canines is diamond but while the The cusp tip is located mesio-lingually in the upper canine it is located mesio-labially in the lower canine.
8. Proximally the Cervical outline is curved incisally however; this curvature is less on the distal than the mesial aspect.
9. The cusp tip of 3 is centralized on the long axis or inclined labially, while the 3-cusp tip is centralized or inclined lingually.
10. The two upper centrals are the thickest of all the incisors & their crowns are the shortest of all human teeth

Give reasons:

1. Extraction of upper central incisor could be done by rotation movement.
2. Mesio-distal section of upper central incisor shows that the pulp chamber has three pulp horns.
3. The lingual inclination is a characteristic feature of lower teeth.
4. In the Mandibular lateral incisor the Crown is twisted disto-lingually on its root.
5. The cusp tips of upper & lower canines are slightly mesial to the center of crown.

MCQs Anteriors & Canines

- 1-The lingual fossa of mandibular anterior teeth when compared to the maxillary anteriors is:
 - c. more concave.
 - d. less concave.
- 2-The height of contour of the facial and lingual surface of the anterior teeth occurs in the:
 - a. mesial third.
 - c. middle third.
 - b. incisal third. d cervical third
- 3-The cingulum of maxillary anterior teeth when compared with the mandibular anterior teeth is:
 - a. less prominent.
 - b. more prominent.
 - c. more prominent in the incisal third
 - d. the same.
- 4-Of the groups of anterior teeth listed below, which group always has a straight root aligned with the longitudinal axis of the tooth?
 - a. right and left maxillary central and lateral incisors
 - b. right and left mandibular central and lateral incisors
 - c. all of the mandibular anterior teeth
 - d. No answer is applicable because all anterior teeth sometimes have an apical deviation of the root.

5-The maxillary canine can be distinguished from the mandibular canine by which of the following characteristics:

- a. The mandibular canine has a less prominent cingulum.
- b. In the mandibular canine, the cusp slopes are short
- c. In the mandibular canine, the incisal edge is located lingually to the center of the tooth.
- d. All of the above.

6-Which of the following is characteristic of the root of the mandibular canine?

- a. It is longer than the root of the maxillary canine.
- b. The root is never bifurcated.
- c. It is flattened or slightly concave on the mesial and distal surfaces.
- d. All of the above.

7-Generally, on the lingual surface of the maxillary canine, there is (are).

- a. one fossa.
- b. three fossae.
- c. two fossae.
- d. four fossae.

8-The contact on the distal surface of the mandibular canine is located:

- a. at the cervical third.
- b. at the incisal third.
- c. at the junction of the incisal and middle third. cervical to the junction of the incisal and middle third.
- d. Cervical to the junction of the incisal and middle one third.

9- compared to other anterior teeth, the mandibular canine is the most likely to have: a longitudinal grooves.

- a. Longitudinal grooves
- b. A root is narrow mesiodistally
- c. Two root canals
- d. Accessory canals

10-As seen from a facial view, the cusp tip of the maxillary canine is:

- a. centered on the midline of the tooth.
- b. displaced mesially from the midline of the tooth.
- c. displaced distally from the midline of the tooth.
- d. highly variable in location.

11-From an incisal view, the cusp tip of the mandibular canine is:

- a. Inclined toward the buccal.
- b. Twisted mesio-lingually.
- c. Inclined lingually.
- d. Very straight.

12-The mandibular canine root is sometimes bifurcated into a:

- a. mesiofacial and distolingual root.
- b. labial and lingual root.
- c. mesial and distal root.
- d. None of the above.

13-Canine teeth exhibit:

- a. a labial ridge.
- b. a lingual ridge.
- c. a mesial marginal ridge.
- d. all of the above.

14-The maxillary and mandibular canines differ in all of the following aspects but one.

Which one is the exception?

- a. prominence of the cingulum
- b. location of facial and lingual heights of contour
- c. marginal ridge prominence
- d. amount of twist of the distal end of the incisal edge

15- In humans, generally, which of the following permanent teeth is the longest?

- a. maxillary central incisor
- b. mandibular canine
- c. mandibular second incisor
- d. maxillary canine

16-The crowns of which of the following teeth are approximately triangular when viewed from the proximal?

- a. premolars
- b. first molars
- c. none of the above
- d. canines

17- The incisal angles of the upper incisors are:

- a. Both mesioincisal and distoincisal angles are sharp in the central and lateral incisors.
- b. Both angles are rounded in the central and lateral incisors.
- c. The mesioincisal angle is sharp while the distoincisal angle is rounded in both the upper central and lateral incisor.
- d. The mesioincisal angle is sharp while the distoincisal is rounded in the upper central incisors, and both angles are rounded in the upper lateral incisor.

18- the mesio-incisal angle and distoincisal angles of the lower central are:

- a. Sharp.
- b. Rounded
- c. The mesioincisal angle is sharp while the distoincisal is rounded.
- d. The mesioincisal angle is rounded while the distoincisal is sharp

19- The mesial and distal contact areas are not at the same level in the following permanent teeth except:

- a. Upper central incisor.
- b. Upper lateral incisor.
- c. Lower central incisor.
- d. d_ Lower lateral incisor.

20-The mesial cusp ridge of the upper and lower permanent canines are:

- a. Shorter than the distal cusp ridges.
- b. Longer than the distal cusp ridges.
- c. Shorter than the distal cusp ridge in the lower canine.
- d. Longer than the distal cusp ridge in the upper canine while shorter than the distal cusp ridge in the lower canine.

21- The lingual ridge divides the lingual fossa into mesial and distal lingual fossae in:

- a. Upper canine.
- b. Lower canine.
- c. Both upper and lower canines.
- d. d-upper lateral incisor

22- From the proximal aspect of the upper canine, the line that bisects the cusp lies:

- a. Labial to the line that bisects the root.
- b. Lingual to the line that bisects the root.
- c. In line with the line that bisects the root.
- d. Mesial to a line bisects the root.

23- The crown of the lower canine looks much longer than that of the upper canine due to:

- a. It has the same length or slightly longer.
- b. Its mesial and distal contact areas are closer incisally to the cusp ridges.
- c. It is narrower mesiodistally.
- d. All the above.

24- The elevations found on the crown of the lower anterior teeth compared to those found on the upper anterior teeth are:

- a. As well developed.
- b. More developed.
- c. Underdeveloped.
- d. d-variable.

25- All the upper and lower anterior teeth have a pulp chamber and single root canal except one of the following that sometimes has two root canals that may open separately in two apical foramina or reunite to open in a single apical foramen, this tooth is:

- a. Upper canine.
- b. Upper lateral incisor.
- c. Lower canine.
- d. Lower lateral incisor.

26- Which one of the permanent teeth is the smallest

- a. mandibular central incisor
- b. mandibular lateral incisor
- c. maxillary-central incisor
- d. maxillary lateral incisor

27- Three of the features listed below are found on the permanent maxillary central incisor. Which of the shown below is NOT found on a maxillary central incisor

- a. cingulum
- b. marginal ridges
- c. cervical ridges
- d. triangular ridges

28-There are four teeth listed below. Which one is most likely to be variable or even

- a. maxillary central incisor
- b. maxillary lateral incisor
- c. maxillary canine
- d. maxillary first premolar

30-There are several teeth listed below. Which one is most likely to have a divided root canal

- a. maxillary central incisor
- b. maxillary canine
- c. maxillary lateral incisor
- d. mandibular canine

31-Upper permanent central incisors are wider than lateral incisors; however, lower permanent central incisors are narrower than lateral incisors

- a. This statement is true
- b. No, this statement is false

32-Which of the following incisors has its mesial and distal contact areas at the same incisocervical level?

- a) Mandibular lateral
- b) Maxillary lateral
- c) Mandibular central
- d) Maxillary central

33-Which one of the following teeth is MOST LIKELY to have a CUSP?

- a. 24
- b. 6
- c. 10
- d. 26

34-At which of the following is the mesial contact area of a maxillary canine located?

- a) the junction of the middle and cervical thirds
- b) the middle third
- c) the junction of the incisal and middle thirds
- d) the incisal third

35- When maxillary and mandibular permanent canines (in the same mouth) are compared, which one of the following statements is TRUE?

- a. Upper canines erupt before lower canines.
- b. upper canine is more bulky than the lower canine
- c. An upper canine has a less pronounced cingulum than a lower canine.
- d. In lower canines, the mesial cusp ridge is longer than the distal cusp ridges.

36-The root of a permanent tooth is completely formed in just about after eruption of that tooth into the mouth.

- a. six months
- b. one year
- c. two years
- d. three years

37-The roots of all but one anterior tooth is likely to have longitudinal grooves on the mesial and distal surface. Which tooth listed below doesn't have these grooves?

- a) Maxillary canine
- b) Maxillary lateral incisor
- c) Maxillary central incisor
- d) Mandibular lateral incisor

38- How does the distoincisor angle of most anterior teeth compare to the mesioincisor angle?

- a) It is straighter
- b) It is more rounded
- c) There is no difference

39-Which anterior tooth listed below is most likely to have a bifurcated root?

- a. Maxillary central incisor
- b. Mandibular canine
- c. Mandibular central incisor
- d. Maxillary canine

40-Which cusp ridge listed below is longest on the permanent canines?

- a. Labial
- b. Distal
- c. Lingual
- d. Mesial

41-Which permanent tooth listed below (other than the third molars) varies in form more than any other tooth?

- a. Maxillary centrals
- c. Mandibular centrals
- b. Maxillary laterals
- d. Mandibular laterals

42- Of the two mandibular incisors, which has a root that is larger in all dimensions?

- a. Mandibular central incisor
- b. Mandibular lateral incisor

43-Which tooth listed below is the smallest and the narrowest (mesial to distal) of all permanent teeth?

- a. Mandibular lateral incisor
- c. Maxillary lateral incisor
- b. Mandibular central incisor
- d. Maxillary central incisor

44- Which permanent canines are usually the last to erupt?

- a. Maxillary canines
- b. Mandibular canines

Key answers:

Complete:

1. shovel
2. Mesio- sharp- distoincisa angle
3. Maxillary lateral incisor.
4. Palatal pit
5. Incisive- curve of the dental arch
6. Shorter
7. Distal- mesial
8. Bifurcation
9. Lingual convergency.

True and False:

1- False 2-True 3- False 4- True 5-False 6-False 7 - False 8- True 9-True 10-False

MCQs Anteriors & Canines:

1	d	20	a	39	b
2	d	21	b	40	d
3	b	22	a	41	b
4	d	23	d	42	b
5	d	24	c	43	b
6	c	25	c	44	a
7	b	26	a		
8	c	27	d		
9	c	28	c		
10	a	29	b		
11	c	30	d		
12	b	31	a		
13	b	32	c		
14	b	33	b		
15	c	34	c		
16	a	35	b		
17	d	36	d		
18	a	37	c		
19	c	38	b		

Give reasons:

- 1- Extraction of upper central incisor could be done by rotation movement because the apex is centralized on the long axis.
- 2- Mesio-distal section of upper central incisor shows that the pulp chamber has three pulp horns because these horns correspond to the three mamelons.
- 3- The lingual inclination is a characteristic feature of lower teeth to facilitate proper occlusion.
- 4- The Crown is twisted disto-lingually on its root to allow the incisal ridge to follow the curve of dental arch.
- 5- The cusp tips of upper & lower canines are slightly mesial to the center of crown because the distal cusp slope is longer than the mesial cusp slope.