

AI1103 ASSIGNMENT 4

Name: MANNAM SARANDEEP, Rollno: CS20BTECH11030

Download the python code from

<https://github.com/sarandeepmannam/ASSIGNMENT4/blob/main/Assignment4.py>

and latex-tikz code from

<https://github.com/sarandeepmannam/ASSIGNMENT4/blob/main/Assignment4.tex>

1 QUESTION-CSIR UGC NET JUNE 2012,Q.50

Let X_1, X_2, \dots be i.i.d $N(1,1)$ random variables. Let $S_n = X_1^2 + X_2^2 + \dots + X_n^2$ for $n \geq 1$. Then

$$\lim_{n \rightarrow \infty} \frac{\text{Var}(S_n)}{n} =$$

- (A) 4
- (B) 6
- (C) 1
- (D) 0

2 SOLUTION-CSIR UGC NET JUNE 2012,Q.50

If two random variables X, Y are independent then X^2, Y^2 are also independent. Since X_1, X_2, X_3, \dots are mutually independent random variables therefore the random variables $X_1^2, X_2^2, X_3^2, \dots$ are also mutually independent.

$$S_n = X_1^2 + X_2^2 + X_3^2 + \dots + X_n^2 \quad (2.0.1)$$

We know that if X and Y are independent random variables $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$.

Therefore,

$$\text{Var}(S_n) = \text{Var}(X_1^2) + \text{Var}(X_2^2) + \dots + \text{Var}(X_n^2) \quad (2.0.2)$$

Since X_1, X_2, \dots, X_n are identically distributed random variables therefore the random variables $X_1^2, X_2^2, \dots, X_n^2$ are also identical. So,

$$\text{Var}(X_1^2) = \text{Var}(X_2^2) = \dots = \text{Var}(X_n^2) \quad (2.0.3)$$

Now we find the variance of X_1^2 .

$$\text{Var}(X_1^2) = E(X_1^4) - (E(X_1^2))^2 \quad (2.0.4)$$

But we don't know the values of $E(X_1^2)$ and $E(X_1^4)$. First we find the value of $E(X_1^2)$.

$$\text{Var}(X_1) = E(X_1^2) - (E(X_1))^2 \quad (2.0.5)$$

Since X_1 is a $N(1,1)$ random variable $E(X_1) = 1$ and $\text{Var}(X_1) = 1$. Therefore,

$$E(X_1^2) - (1)^2 = 1 \quad (2.0.6)$$

$$E(X_1^2) = 2 \quad (2.0.7)$$

Now we find the value of $E(X_1^4)$.

Since X_1 is a $N(1,1)$ random variable pdf of X_1 will be

$$f(X_1) = \frac{1}{\sqrt{2\pi}} \left(e^{-\frac{(X_1-1)^2}{2}} \right) \quad (2.0.8)$$




Fig. 4: PDF of X_1, X_2, \dots

Now expected value of X_1^4 will be,

$$E(X_1^4) = \int_{-\infty}^{+\infty} X_1^4 f(X_1) dX_1 \quad (2.0.9)$$

$$= \int_{-\infty}^{+\infty} X_1^4 \left(\frac{1}{\sqrt{2\pi}} \left(e^{-\frac{(X_1-1)^2}{2}} \right) \right) dX_1 \quad (2.0.10)$$

Now put $X_1 - 1 = t$,

$$E(X_1^4) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} (t+1)^4 \left(e^{-\frac{t^2}{2}} \right) dt \quad (2.0.11)$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} (t^4 + 4t^3 + 6t^2 + 4t + 1) \left(e^{-\frac{t^2}{2}} \right) dt \quad (2.0.12)$$

Since $\int_{-\infty}^{+\infty} 4t^3 \left(e^{-\frac{t^2}{2}} \right) dt$ and $\int_{-\infty}^{+\infty} 4t \left(e^{-\frac{t^2}{2}} \right) dt$ are equal to zero, the equation will get reduced to

$$E(X_1^4) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} (t^4 + 6t^2 + 1) \left(e^{-\frac{t^2}{2}} \right) dt \quad (2.0.13)$$

Now put $\frac{t}{\sqrt{2}} = x$,

$$E(X_1^4) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{+\infty} (4x^4 + 12x^2 + 1) \left(e^{-x^2} \right) dx \quad (2.0.14)$$

We know that,

$$\int_{-\infty}^{+\infty} \left(e^{-ax^2} \right) dx = \frac{\sqrt{\pi}}{\sqrt{a}} \quad (2.0.15)$$

$$\frac{d \left(\int_{-\infty}^{+\infty} \left(e^{-ax^2} \right) dx \right)}{da} = \frac{d}{da} \left(\frac{\sqrt{\pi}}{\sqrt{a}} \right) \quad (2.0.16)$$

$$\int_{-\infty}^{+\infty} x^2 \left(e^{-ax^2} \right) dx = \frac{\sqrt{\pi}}{2\sqrt{a^3}} \quad (2.0.17)$$

similarly,

$$\int_{-\infty}^{+\infty} x^4 \left(e^{-ax^2} \right) dx = \frac{3\sqrt{\pi}}{4\sqrt{a^5}} \quad (2.0.18)$$

Using (2.0.15), (2.0.17) and (2.0.18) in (2.0.14) we

get,

$$E(X_1^4) = \frac{1}{\sqrt{\pi}} \left(4 \left(\frac{3\sqrt{\pi}}{4} \right) + 12 \left(\frac{\sqrt{\pi}}{2} \right) + \sqrt{\pi} \right) \quad (2.0.19)$$

$$= 3 + 6 + 1 \quad (2.0.20)$$

$$\Rightarrow E(X_1^4) = 10 \quad (2.0.21)$$

Substituting (2.0.21), (2.0.7) in (2.0.4) we get,

$$\text{Var}(X_1^2) = 10 - 4 \quad (2.0.22)$$

$$\Rightarrow \text{Var}(X_1^2) = 6 \quad (2.0.23)$$

By the equations (2.0.23), (2.0.3) and (2.0.2), we can conclude that

$$\text{Var}(S_n) = 6n \quad (2.0.24)$$

$$\lim_{n \rightarrow \infty} \frac{\text{Var}(S_n)}{n} = \lim_{n \rightarrow \infty} \frac{6n}{n} = \lim_{n \rightarrow \infty} 6 = 6$$

Hence, option (B) is correct.