

Assignment5

Saransh Bali

Abstract—This a simple document that explains the geometry in conics.

https://github.com/saranshbali/EE5609/blob/master/Assignment1/Code/assignment1_n.ipynb

Download all latex-tikz codes from

<https://github.com/saranshbali/EE5609/blob/master/Orthogonality>

1 PROBLEM

Through what angle must the axes be turned to reduce the equation

$$\mathbf{x}^T \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix} \mathbf{x} = 1 \quad (1.0.1)$$

to the form

$$\mathbf{x}^T \begin{pmatrix} 0 & 1/2 \\ 1/2 & 0 \end{pmatrix} \mathbf{x} = c \quad (1.0.2)$$

where c is a constant.

2 SOLUTION

The general second order equation can be expressed as follows,

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.1)$$

From (1.0.1) and (2.0.1)

$$\mathbf{V} = \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.3)$$

$$f = -1 \quad (2.0.4)$$

The matrix \mathbf{V} can be decomposed as,

$$\mathbf{V} = \mathbf{P} \mathbf{D} \mathbf{P}^T \quad \mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \quad (2.0.5)$$

where λ_1 and λ_2 are the eigen values of \mathbf{V} , and \mathbf{P} contains the eigen vectors corresponding to the

eigen values λ_1 and λ_2 . The affine transformation is given by,

$$\mathbf{x} = \mathbf{P} \mathbf{y} + \mathbf{c} \quad (2.0.6)$$

where, \mathbf{P} indicates the rotation of axes and \mathbf{c} indicates the shift of origin.

Eigen values of \mathbf{V} are,

$$|\mathbf{V} - \lambda \mathbf{I}| = 0 \quad (2.0.7)$$

$$\Rightarrow \begin{vmatrix} 1 - \lambda & -1 \\ -1 & -1 - \lambda \end{vmatrix} = 0 \quad (2.0.8)$$

$$\Rightarrow (1 - \lambda)(-1 - \lambda) - 1 = 0 \quad (2.0.9)$$

$$\Rightarrow \lambda^2 - 2 = 0 \quad (2.0.10)$$

$$\Rightarrow \lambda = \pm \sqrt{2}, \quad \mathbf{D} = \begin{pmatrix} \sqrt{2} & 0 \\ 0 & -\sqrt{2} \end{pmatrix} \quad (2.0.11)$$

Eigen vector for $\lambda_1 = \sqrt{2}$,

$$\begin{aligned} \mathbf{V} - \lambda_1 \mathbf{I} &= \begin{pmatrix} 1 - \sqrt{2} & -1 \\ -1 & -1 - \sqrt{2} \end{pmatrix} \\ &\xrightarrow{r_1/1-\sqrt{2}} \begin{pmatrix} 1 & -1/1-\sqrt{2} \\ -1 & -1-\sqrt{2} \end{pmatrix} \end{aligned} \quad (2.0.12)$$

$$\begin{aligned} \mathbf{V} - \lambda_1 \mathbf{I} &= \begin{pmatrix} 1 & -1/1-\sqrt{2} \\ -1 & -1-\sqrt{2} \end{pmatrix} \\ &\xrightarrow{r_2=r_1+r_2} \begin{pmatrix} 1 & -1/1-\sqrt{2} \\ 0 & 0 \end{pmatrix} \end{aligned} \quad (2.0.13)$$

Hence,

$$\mathbf{P}_1 = \begin{pmatrix} 1 - \sqrt{2} \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} \\ 1 \end{pmatrix} \quad (2.0.14)$$

Eigen vector for $\lambda_2 = -\sqrt{2}$,

$$\begin{aligned} \mathbf{V} - \lambda_2 \mathbf{I} &= \begin{pmatrix} 1 + \sqrt{2} & -1 \\ -1 & -1 + \sqrt{2} \end{pmatrix} \\ &\xrightarrow{r_1/1+\sqrt{2}} \begin{pmatrix} 1 & -1/1+\sqrt{2} \\ -1 & -1+\sqrt{2} \end{pmatrix} \end{aligned} \quad (2.0.15)$$

$$\mathbf{V} - \lambda_2 \mathbf{I} = \begin{pmatrix} 1 & -1/1 + \sqrt{2} \\ -1 & -1 + \sqrt{2} \end{pmatrix} \\ \xleftrightarrow{r_2=r_1+r_2} \begin{pmatrix} 1 & -1/1 + \sqrt{2} \\ 0 & 0 \end{pmatrix} \quad (2.0.16)$$

Hence,

$$\mathbf{P}_2 = \begin{pmatrix} 1 + \sqrt{2} \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{1+\sqrt{2}}{4+2\sqrt{2}} \\ \frac{1}{4+2\sqrt{2}} \end{pmatrix} \quad (2.0.17)$$

Thus,

$$\mathbf{P} = \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} & \frac{1+\sqrt{2}}{4+2\sqrt{2}} \\ \frac{1}{4-2\sqrt{2}} & \frac{1}{4+2\sqrt{2}} \end{pmatrix} \quad (2.0.18)$$

Therefore \mathbf{V} can be written as,

$$\mathbf{V} = \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} & \frac{1+\sqrt{2}}{4+2\sqrt{2}} \\ \frac{1}{4-2\sqrt{2}} & \frac{1}{4+2\sqrt{2}} \end{pmatrix} \begin{pmatrix} \sqrt{2} & 0 \\ 0 & -\sqrt{2} \end{pmatrix} \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} & \frac{1}{4-2\sqrt{2}} \\ \frac{1+\sqrt{2}}{4+2\sqrt{2}} & \frac{1}{4+2\sqrt{2}} \end{pmatrix} \quad (2.0.19)$$

Now, (1.0.1) can be transformed as

$$\mathbf{x}^T \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} & \frac{1+\sqrt{2}}{4+2\sqrt{2}} \\ \frac{1}{4-2\sqrt{2}} & \frac{1}{4+2\sqrt{2}} \end{pmatrix} \begin{pmatrix} \sqrt{2} & 0 \\ 0 & -\sqrt{2} \end{pmatrix} \begin{pmatrix} \frac{1-\sqrt{2}}{4-2\sqrt{2}} & \frac{1}{4-2\sqrt{2}} \\ \frac{1+\sqrt{2}}{4+2\sqrt{2}} & \frac{1}{4+2\sqrt{2}} \end{pmatrix} \mathbf{x} = 1 \quad (2.0.20)$$