

Assignment 14

Saransh Bali

Abstract—This is a simple document about the algebra of polynomials.

Download latex-tikz from

<https://github.com/saranshbali/EE5609/blob/master/Assignment14>

1 PROBLEM

Let \mathbf{F} be a sub-field of the complex numbers and let \mathbf{D} be the transformation on $\mathbf{F}[x]$ defined by

$$\mathbf{D}\left(\sum_{i=0}^n c_i x^i\right) = \sum_{i=0}^n i c_i x^{i-1} \quad (1.0.1)$$

Show that \mathbf{D} is a linear operator on $\mathbf{F}[x]$ and find its null space.

2 DEFINITION AND RESULT USED

Linear Transformation	A linear transformation from \mathbf{V} into \mathbf{W} is a function \mathbf{T} from \mathbf{V} into \mathbf{W} such that $\mathbf{T}(c\alpha + \beta) = c\mathbf{T}(\alpha) + \mathbf{T}(\beta)$ $\forall \alpha$ and β in \mathbf{V} and \forall scalars c in \mathbf{F} .
$\mathbf{F}[x]$	Let $\mathbf{F}[x]$ be the subspace of \mathbf{F}^∞ spanned by the vectors $1, x, x^2, \dots$. An element of $\mathbf{F}[x]$ is called a polynomial over \mathbf{F} .
Differentiation Transformation	Let \mathbf{F} be a field and let \mathbf{V} be the space of polynomial functions f from \mathbf{F} into \mathbf{F} , given by $f(x) = c_0 + c_1 x + \dots + c_k x^k$ Then, $\mathbf{D}f(x) = c_1 + 2c_2 x + \dots + k c_k x^{k-1}$ is called Differentiation Transformation. The Differentiation Transformation is a Linear map because $\begin{aligned} \mathbf{D}(cf + g)(x) &= (c.c_1 + c'_1) + 2(c.c_2 + c'_2)x + \dots + k(c.c_k + c'_k)x^{k-1} \\ &= c.c_1 + 2c.c_2 x + \dots + k c.c_k x^{k-1} + c'_1 + 2c'_2 x + \dots + k c'_k x^{k-1} \\ &= c\mathbf{D}f(x) + \mathbf{D}g(x) \end{aligned}$

3 SOLUTION

Proving D is Linear	<p>From (1.0.1), clearly D is a function from $\mathbf{F}[\mathbf{x}]$ to $\mathbf{F}[\mathbf{x}]$. We must show that D is linear. Clearly D is a Differentiation Transformation, and hence is linear. In other words</p> $ \begin{aligned} \mathbf{D}\left(\sum_{i=0}^n c_i x^i + \sum_{i=0}^n c'_i x^i\right) &= \mathbf{D}\left(\sum_{i=0}^n (c_i + c'_i) x^i\right) \\ &= \sum_{i=0}^n i (c_i + c'_i) x^{i-1} \\ &= c \sum_{i=0}^n i x^{i-1} + \sum_{i=0}^n i c'_i x^{i-1} \\ &= c \mathbf{D}\left(\sum_{i=0}^n x^i\right) + \mathbf{D}\left(\sum_{i=0}^n c'_i x^i\right) \end{aligned} $ <p>Hence, D is a linear transformation.</p>
Null Space of D	<p>Let $\mathbf{N}(\mathbf{D})$ denotes the nullspace of D. Then</p> $\mathbf{N}(\mathbf{D}) = \{f \in \mathbf{F}[\mathbf{x}] : \mathbf{D}f(x) = 0\}$ <p>A polynomial is zero if and only if its every coefficient is zero. Thus, it must be such that each $c_1 = c_2 = \dots = 0$. Since, D is a Differentiation Transformation and we know that derivative of a constant polynomial is zero. Thus, the nullspace of D contains the constant polynomials. Hence,</p> $\mathbf{N}(\mathbf{D}) = \{f \in \mathbf{F}[\mathbf{x}] : f(x) = c\}$