

Large Time Behaviour and Eigenvalue Problems for Finite State Mean-Field Particle Systems

Sarath A Y, Rajesh Sundaresan

Indian Institute of Science

17 Aug 2019

System model

- ▶ N particles. Finite state space \mathcal{Z} .

System model

- ▶ N particles. Finite state space \mathcal{Z} .
- ▶ State of particle n at time t is $X_n^N(t) \in \mathcal{Z}$.

System model

- ▶ N particles. Finite state space \mathcal{Z} .
- ▶ State of particle n at time t is $X_n^N(t) \in \mathcal{Z}$.
- ▶ Empirical measure at time t :

$$\mu_N(t) = \frac{1}{N} \sum_{n=1}^N \delta_{X_n^N(t)} \in M_1(\mathcal{Z}).$$

System model

- ▶ N particles. Finite state space \mathcal{Z} .
- ▶ State of particle n at time t is $X_n^N(t) \in \mathcal{Z}$.
- ▶ Empirical measure at time t :

$$\mu_N(t) = \frac{1}{N} \sum_{n=1}^N \delta_{X_n^N(t)} \in M_1(\mathcal{Z}).$$

- ▶ Mean-field interaction: $z \rightarrow z'$ transition at rate $\lambda_{z,z'}(\mu_N(t))$.

System model

- ▶ N particles. Finite state space \mathcal{Z} .
- ▶ State of particle n at time t is $X_n^N(t) \in \mathcal{Z}$.
- ▶ Empirical measure at time t :

$$\mu_N(t) = \frac{1}{N} \sum_{n=1}^N \delta_{X_n^N(t)} \in M_1(\mathcal{Z}).$$

- ▶ Mean-field interaction: $z \rightarrow z'$ transition at rate $\lambda_{z,z'}(\mu_N(t))$.
- ▶ μ_N is a Markov process on $M_1(\mathcal{Z})$.

System model

- ▶ N particles. Finite state space \mathcal{Z} .
- ▶ State of particle n at time t is $X_n^N(t) \in \mathcal{Z}$.
- ▶ Empirical measure at time t :

$$\mu_N(t) = \frac{1}{N} \sum_{n=1}^N \delta_{X_n^N(t)} \in M_1(\mathcal{Z}).$$

- ▶ Mean-field interaction: $z \rightarrow z'$ transition at rate $\lambda_{z,z'}(\mu_N(t))$.
- ▶ μ_N is a Markov process on $M_1(\mathcal{Z})$.
- ▶ Goal: understand large time behaviour of μ_N , and convergence to stationarity.

An Example: Interaction in WiFi networks

- ▶ N nodes accessing a common wireless medium.

An Example: Interaction in WiFi networks

- ▶ N nodes accessing a common wireless medium.
- ▶ Interaction among nodes via the distributed MAC protocol.

An Example: Interaction in WiFi networks

- ▶ N nodes accessing a common wireless medium.
- ▶ Interaction among nodes via the distributed MAC protocol.
- ▶ State $X_n^N(t)$ represents aggressiveness of packet transmission.

An Example: Interaction in WiFi networks

- ▶ N nodes accessing a common wireless medium.
- ▶ Interaction among nodes via the distributed MAC protocol.
- ▶ State $X_n^N(t)$ represents aggressiveness of packet transmission.
- ▶




Figure: Set of allowed transitions in WiFi example

- ▶ State evolution:
 - ▶ Becomes less aggressive after a collision.
 - ▶ Moves to the most aggressive state after a successful packet transmission.

A sample path of μ_N in WiFi example




Figure: Evolution of states in a WiFi network under the MAC protocol

A sample path of μ_N in WiFi example




Figure: Evolution of states in a WiFi network under the MAC protocol

- Multiple stable regions in the system. Transition between two stable region occur over large time durations.

Background on large deviations of μ_N

- ▶ Let $\mu_N(0) \rightarrow \nu$. Then, over $[0, T]$, the process μ_N is close to the McKean-Vlasov equation:

$$\dot{\mu}(t) = \Lambda_{\mu(t)}^* \mu(t), \quad \mu(0) = \nu.$$

Background on large deviations of μ_N

- ▶ Let $\mu_N(0) \rightarrow \nu$. Then, over $[0, T]$, the process μ_N is close to the McKean-Vlasov equation:

$$\dot{\mu}(t) = \Lambda_{\mu(t)}^* \mu(t), \mu(0) = \nu.$$

- ▶ Large deviations of μ_N in $D([0, T], M_1(\mathcal{Z}))$.

Background on large deviations of μ_N

- ▶ Let $\mu_N(0) \rightarrow \nu$. Then, over $[0, T]$, the process μ_N is close to the McKean-Vlasov equation:

$$\dot{\mu}(t) = \Lambda_{\mu(t)}^* \mu(t), \mu(0) = \nu.$$

- ▶ Large deviations of μ_N in $D([0, T], M_1(\mathcal{Z}))$.
- ▶ Multiple ω -limit sets for this dynamics: stable attractors, limit cycles, etc.

Background on large deviations of μ_N

- ▶ Let $\mu_N(0) \rightarrow \nu$. Then, over $[0, T]$, the process μ_N is close to the McKean-Vlasov equation:

$$\dot{\mu}(t) = \Lambda_{\mu(t)}^* \mu(t), \mu(0) = \nu.$$

- ▶ Large deviations of μ_N in $D([0, T], M_1(\mathcal{Z}))$.
- ▶ Multiple ω -limit sets for this dynamics: stable attractors, limit cycles, etc.
- ▶ Freidlin-Wentzell quasipotential $\tilde{V}(K_i, K_j)$: minimum cost of moving from K_i to K_j .

Background on large deviations of μ_N

- ▶ Let $\mu_N(0) \rightarrow \nu$. Then, over $[0, T]$, the process μ_N is close to the McKean-Vlasov equation:

$$\dot{\mu}(t) = \Lambda_{\mu(t)}^* \mu(t), \mu(0) = \nu.$$

- ▶ Large deviations of μ_N in $D([0, T], M_1(\mathcal{Z}))$.
- ▶ Multiple ω -limit sets for this dynamics: stable attractors, limit cycles, etc.
- ▶ Freidlin-Wentzell quasipotential $\tilde{V}(K_i, K_j)$: minimum cost of moving from K_i to K_j .
- ▶ Obtain one-step transition probability of μ_N near the ω -limits sets:

$$P(K_i, K_j) \simeq \exp\{-N\tilde{V}(K_i, K_j)\}.$$

Main results

- ▶ Large time behaviour of μ_N (in terms of \tilde{V}):
 - ▶ Mean exit time from ω -limit sets.
 - ▶ Estimates on probability of reaching a given ω -limit set.
 - ▶ Most likely cycle of ω -limits sets.

Main results

- ▶ Large time behaviour of μ_N (in terms of \tilde{V}):
 - ▶ Mean exit time from ω -limit sets.
 - ▶ Estimates on probability of reaching a given ω -limit set.
 - ▶ Most likely cycle of ω -limits sets.
- ▶ Mixing and convergence to the invariant measure:
 - ▶ A constant $\Lambda \geq 0$ in terms of \tilde{V} .

Main results

- ▶ Large time behaviour of μ_N (in terms of \tilde{V}):
 - ▶ Mean exit time from ω -limit sets.
 - ▶ Estimates on probability of reaching a given ω -limit set.
 - ▶ Most likely cycle of ω -limits sets.
- ▶ Mixing and convergence to the invariant measure:
 - ▶ A constant $\Lambda \geq 0$ in terms of \tilde{V} .
 - ▶ If time is of the order $\exp\{N(\Lambda - O(1))\}$, a lower bound on the transition probability of μ_N .

Main results

- ▶ Large time behaviour of μ_N (in terms of \tilde{V}):
 - ▶ Mean exit time from ω -limit sets.
 - ▶ Estimates on probability of reaching a given ω -limit set.
 - ▶ Most likely cycle of ω -limits sets.
- ▶ Mixing and convergence to the invariant measure:
 - ▶ A constant $\Lambda \geq 0$ in terms of \tilde{V} .
 - ▶ If time is of the order $\exp\{N(\Lambda - O(1))\}$, a lower bound on the transition probability of μ_N .
 - ▶ If time is of the order $\exp\{N(\Lambda + O(1))\}$, μ_N is very close to its invariant measure.

Main results

- ▶ Large time behaviour of μ_N (in terms of \tilde{V}):
 - ▶ Mean exit time from ω -limit sets.
 - ▶ Estimates on probability of reaching a given ω -limit set.
 - ▶ Most likely cycle of ω -limits sets.
- ▶ Mixing and convergence to the invariant measure:
 - ▶ A constant $\Lambda \geq 0$ in terms of \tilde{V} .
 - ▶ If time is of the order $\exp\{N(\Lambda - O(1))\}$, a lower bound on the transition probability of μ_N .
 - ▶ If time is of the order $\exp\{N(\Lambda + O(1))\}$, μ_N is very close to its invariant measure.
- ▶ Scaling of the second largest eigenvalue (assuming reversibility): $\lambda_2^N \simeq \exp\{-N\Lambda\}$.
 - ▶ Consequence: μ_N mixes slowly if there are metastable states ($\Lambda > 0$).