

# Functional neuroimaging of resilience to trauma: convergent evidence and challenges for future research

Agnes Norbury<sup>1,2\*</sup>, Saren H. Seeley<sup>1\*</sup>, M. Mercedes Perez-Rodriguez<sup>1</sup>, Adriana Feder<sup>1</sup>

<sup>1</sup> *Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA*

<sup>2</sup> *Applied Computational Psychiatry Lab, Max Planck Centre for Computational Psychiatry and Ageing Research, Queen Square Institute of Neurology and Mental Health Neuroscience Department, University College London, London, UK*

Corresponding author: Adriana Feder, MD, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA. Email: [adriana.feder@mssm.edu](mailto:adriana.feder@mssm.edu)

\* Shared first authorship; authors contributed equally.

## Word counts:

Abstract: 119

Main text: 5,049 (including in-text citations)

## Acknowledgments

This work was supported by the CDC National Institute for Occupational Safety and Health (U01OH011473) and the NIH National Institute on Mental Health (T32MH122394). This content is solely the responsibility of the authors and does not necessarily represent the official views of the CDC or NIH.

## Conflict of Interest

Dr Feder is named co-inventor on a patent application in the US, and several issued patents outside the US filed by the Icahn School of Medicine at Mount Sinai related to the use of ketamine for the treatment of PTSD. This intellectual property has not been licensed. Dr. Perez-Rodriguez has received research grant funding from Neurocrine Biosciences, Inc, Millennium Pharmaceuticals, Takeda, and Al Cure. She is a consultant for Neurocrine Biosciences, Inc. and Alkermes. She has served on an advisory board for Neurocrine Biosciences Inc. Drs. Norbury and Seeley have no conflicts of interest to declare.

## **Abstract**

Resilience is broadly defined as the ability to adapt successfully following stressful life events. Here, we review functional MRI studies that investigated key psychological factors that have been consistently linked to resilience to severe adversity and trauma exposure. These domains include emotion regulation (including cognitive reappraisal), reward responsivity, and cognitive control. Further, we briefly review functional imaging evidence related to emerging areas of study that may potentially facilitate resilience: namely social cognition, active coping, and successful fear extinction. Finally, we also touch upon ongoing issues in neuroimaging study design that will need to be addressed to enable us to harness insight from such studies to improve treatments for – or, ideally, guard against the development of – debilitating post-traumatic stress syndromes.

## 1 Introduction

2 Trauma exposure does not invariably result in adverse psychiatric outcomes.  
3 Epidemiological estimates for clinically significant posttraumatic stress symptoms range from 5-  
4 22% after exposure to natural disasters, severe injuries, or assault, to 46-65% after sexual  
5 violence ((Bromet, Karam, Koenen, & Stein, 2018; Shalev, Liberzon, & Marmar, 2017; Watson,  
6 2019). Similarly, 10-25% of those exposed to significant childhood maltreatment show better-  
7 than-expected functioning (Walsh, Dawson, & Mattingly, 2010).

8 Broadly, resilience is defined as successful adaptation following trauma, adversity, or  
9 stressful life events (Feder, Fred-Torres, Southwick, & Charney, 2019; Kalisch et al., 2017;  
10 Southwick, Charney, & DePierro, 2023). Beyond vulnerability or risk for psychiatric disorders,  
11 resilience requires having experienced adversity or trauma (Yehuda & Flory, 2007) (**Box 1:**  
12 **Quantifying trauma**). Resilience research aims to identify mechanisms to improve prevention  
13 and treatment of posttraumatic stress and trauma-related disorders. In neuroimaging studies,  
14 resilience is typically defined as absence of psychopathology despite exposure to adversity or  
15 trauma, but in some studies *alternatively* as high scores on measures of trait resilience, e.g., the  
16 Connor-Davidson Resilience Scale (CD-RISC) (Connor & Davidson, 2003a) (**Box 2:**  
17 **Quantifying resilience**), notwithstanding significant variation in its operationalization and  
18 measurement (Denckla et al., 2020; Feder et al., 2019; Kalisch et al., 2017; Southwick et al.,  
19 2023).

20

## 21 Resilience-related psychological factors

22 Despite these definitional differences, decades of research have identified psychological  
23 factors that promote resilience to trauma and severe adversity. Some widely replicated and  
24 potentially modifiable factors include effective emotion regulation, positive emotionality,  
25 cognitive flexibility and control, facing fears and active coping, and ability to harness social  
26 support (reviewed in (Seeley, Boukezzi, DePierro, Charney, & Feder, In press; Southwick et al.,  
27 2023)). Effective emotion regulation is associated with higher executive control and supports  
28 adaptive coping. Facing fears, likely facilitated by successful fear extinction plus cognitive  
29 flexibility, allows for critical appraisal of threats and active coping via tackling stressors or

problem solving. Positive emotions and related reward system function also promote resilience by supporting positive reframing and the ability to harness social support, which in turn serves as a safety net and facilitates stress recovery. The ability to harness social support draws partly on competent social cognition – capacity to accurately ‘read’ and respond to others’ intentions of others – making social cognition a potential resilience-linked factor.

## **The current review: neural correlates of psychological factors in resilience**

Building on work identifying core psychological factors associated with resilience (Feder et al., 2019; Seeley et al., In press; Southwick et al., 2023) and previous neuroimaging resilience reviews in adults and/or youth (Méndez Leal & Silvers, 2021; Moreno-López et al., 2020; van der Werff, van den Berg, Pannekoek, Elzinga, & Van Der Wee, 2013), our narrative review covers human fMRI studies of neural circuitry underlying psychological factors associated with resilience to trauma and severe adversity. We organize fMRI studies around *emotion regulation*, *reward responsivity*, and *cognitive control*, e.g., (Dennison et al., 2016; Holz, Tost, & Meyer-Lindenberg, 2020; Kaldewaij et al., 2021; van der Werff et al., 2013). As described above, *active coping*, *facing fears* (and related fear extinction) (Careaga, Girardi, & Suchecki, 2016), and the ability to harness *social support* (including competent *social cognition*) (Lepore & Kliever, 2019; Stevens & Jovanovic, 2019) have also been linked to resilience, yet there are fewer resilience neuroimaging studies of them as resilience-related psychological factors. Here, we include these factors as “emerging areas”, as they might illuminate additional pathways to resilience and underlying neural circuitry.

Cross-sectional fMRI studies of resilient individuals typically focus on (1) youth and adults who have experienced childhood maltreatment or (2) adults exposed to trauma in adulthood including severe accidents, assaults, or occupation-related incidents (e.g., first-responders, military). In this narrative review, we only discuss cross-sectional studies with a key comparison group of healthy, non-trauma-exposed participants (in addition to resilient and symptomatic groups), because this allows researchers to disentangle effects of trauma exposure and psychopathology (see van der Werff et al., 2013). Also included are several studies of neural correlates of high trait resilience.

We focus on literature published after van der Werff and colleagues' 2013 review. Due to recent large cohort studies and improved reporting standards for neuroimaging, this allows us to focus on better powered work. Within each section, we first cover cross-sectional findings in resilient trauma survivors (children; adults) and high trait resilience (when available). Additionally, we discuss longitudinal studies and a few available interventional studies that contribute new insights on the neural correlates of resilience. Although most studies reviewed here focus on task-based regional brain activity, we also include relevant functional connectivity findings, including resting state fMRI.

[Box 1 around here]

[Box 2 around here]

## Functional neuroimaging of resilience mechanisms

### Emotion regulation

Successful emotion regulation is key in resilience (Troy & Mauss, 2011). Resilient individuals downregulate neural responses to stress or threat via medial prefrontal cortex (mPFC) inhibition of emotional reactivity in a network of brain regions, including the amygdala and anterior cingulate cortex (ACC); conversely, absent or low resilience is associated with hyperreactivity to emotional stimuli, both trauma-related and -unrelated (Patel, Spreng, Shin, & Girard, 2012; Whittle et al., 2013). In an earlier review, resilient adults (exposed to trauma in childhood and/or adulthood) showed greater mPFC engagement, including rostral ACC, and lesser activity in the dorsal ACC and amygdala, compared to symptomatic trauma-exposed adults across several studies (van der Werff et al., 2013).

Cross-sectional studies in youth with a history of childhood maltreatment show increased frontolimbic connectivity in resilient youth (Demers et al., 2018; Moreno-López et al., 2020); for example greater down-regulation of amygdala activity during emotion regulation (Schweizer et al., 2016). Higher-trait resilience youth living in adverse environments spent more time than lower-trait resilience peers in a dynamic resting state functional connectivity pattern of less salience network connectivity with both the central executive and default mode networks. This

could suggest lesser influence of emotional reactivity during unconstrained thought (Iadipalo et al., 2018).

Cross-sectional findings are bolstered by several recent longitudinal studies of early adversity. Among adopted children who experienced early institutional care, self-reported anxiety at three-year follow-up decreased more in children who showed reduced amygdala responses to images of their adoptive parents (versus strangers) at baseline (Callaghan et al., 2019). Similarly, previously-institutionalized youth showing stronger ventromedial (vm)PFC-hippocampal functional connectivity during an aversive learning task reported lower anxiety at two-year follow-up (Silvers et al., 2016). A longitudinal study of internalizing symptoms in older adolescents and youth with a childhood maltreatment history compared to propensity-matched controls found that greater baseline amygdala threat reactivity predicted later internalizing symptoms – independently of stressful life events and factors such as socioeconomic status, suggesting that lesser limbic reactivity may be a unique predictor of resilience to childhood maltreatment (Gerin et al., 2019).

Cross-sectional findings in adults are similar to those in youth. Healthy adults with a childhood maltreatment history demonstrated greater frontal inhibition of emotional distractors (Demers et al., 2021). In another study of resilient adults with a childhood maltreatment history, blunted amygdala response correlated with higher depression symptoms (Yamamoto et al., 2017). However, in Yamamoto and colleagues' sample of healthy individuals without clinically significant symptoms, it is notable that higher amygdala reactivity was accompanied by a compensatory increase in prefrontal functional connectivity, linked to top-down emotion regulation. Studies of adulthood trauma exposure similarly suggest lesser amygdala reactivity and greater mPFC regulation in resilient adults. Lower baseline amygdala reactivity to fearful faces at 2-3 weeks post-incident was associated with lower self-reported PTSD symptoms at one-year follow-up, in individuals recruited from a hospital emergency department (Stevens et al., 2017). Another study, using Granger causality analysis, revealed more resting state mPFC inhibition of the amygdala in resilient typhoon survivors, compared to both symptomatic survivors and non-trauma-exposed controls (Chen et al., 2018a).

Longitudinal work in adults links pre-exposure neural functioning to later outcomes. In a large prospective sample of police trainees, pre-exposure anterior PFC activity during an

emotional action control task predicted lower PTSD symptoms at follow-up. Anterior PFC activity moderated the positive relationship between previous trauma load (a risk factor for PTSD) and PTSD symptom levels at follow-up, such that cumulative number of lifetime trauma exposures was not linked to higher PTSD symptoms in trainees with higher anterior PFC activity (Kaldewaij et al., 2021). In another cohort of police trainees scanned pre- and post-trauma exposure, those with higher post-traumatic intrusion symptoms showed increased salience network resting state connectivity after stress induction (Zhang et al., 2022). Together, these studies suggest that prefrontal regulation of limbic reactivity to emotional threat cues may predict resilience to posttraumatic stress and anxiety symptoms. In Kaldewaij et al. (2021), amygdala reactivity appeared to be acquired, rather than a prospective marker of PTSD vulnerability. However, previous longitudinal studies linked stress vulnerability to higher baseline amygdala reactivity (Admon et al., 2009; Swartz, Knodt, Radtke, & Hariri, 2015). Longitudinal prospective cohort studies, such as the ongoing Advancing Understanding of Recovery After Trauma (AURORA) study (McLean et al., 2020), which follows patients scanned 2-3 weeks after visiting the emergency department post-trauma, hold promise for a more granular understanding of emotional responding in resilience. Notably, there may be multiple pathways to risk/resilience: cross-task cluster analysis in an AURORA cohort, replicated in a different cohort with a wider range of trauma exposures, identified two vulnerability-related “biotypes” characterized by heightened emotional reactivity to threat, but differential reactivity to reward, detailed in a later section (Stevens et al., 2021).

From an interventional perspective, a pilot trial of fMRI-guided real-time neurofeedback training decreased amygdala activation and increased amygdala-vmPFC connectivity in combat veterans, but was not superior to sham in reducing PTSD symptoms (Zotev et al., 2018). However, training a pre-deployment military sample to downregulate EEG-derived amygdala signal successfully decreased amygdala BOLD signal and increased amygdala-mPFC functional connectivity at follow-up (Keynan et al., 2019). Active training improved experimental emotional regulation indices, but had no effect on self-reported anxiety (Keynan et al., 2019). Neurofeedback training represents a step towards developing new interventions for at-risk individuals. However, whether this approach will be effective in preventing or mitigating post-traumatic distress requires further study.

1       **Cognitive reappraisal.** Cognitive reappraisal is a deliberate form of emotion regulation  
 2 that buffers risk for adverse outcomes following trauma exposure (Rodman, Jenness,  
 3 Weissman, Pine, & McLaughlin, 2019). The ability to re-evaluate the meaning of experiences  
 4 promotes adaptation after trauma, and represents a core part of psychological interventions  
 5 (Hofmann, Asmundson, & Beck, 2013). For example, in an ecological momentary assessment  
 6 study in individuals remitted from depression, positive appraisal (ability to “*focus on positive*  
 7 *meaning*”) and feelings of resilience were mutually reinforcing. Positive reappraisal was also  
 8 associated with lower incidence of residual depressive symptoms (Hoorelbeke, Van den Bergh,  
 9 Wichers, & Koster, 2019). Neuroimaging meta-analyses highlight ventrolateral (vl)PFC and  
 10 dorsolateral (dl)PFC, dorsal ACC, and amygdala function in effortful emotional regulation in  
 11 healthy individuals – and highlight functional differences in healthy vs. symptomatic participants  
 12 (Zilverstand, Parvaz, & Goldstein, 2017).

13       Laboratory studies of trauma-exposed individuals found inconsistent evidence for a link  
 14 between psychological symptoms and neural activity during experimental tasks instructing  
 15 participants to deliberately modulate their emotional response to affective images using cognitive  
 16 reframing. A longitudinal study in maltreated youth found a relationship between prefrontal  
 17 recruitment during effortful cognitive reappraisal (and associated downregulation of amygdala  
 18 reactivity to negative stimuli) and subsequent risk for mood symptoms, but no relation to anxiety  
 19 symptoms (Rodman et al., 2019). In a small sample of adolescent girls exposed to violent  
 20 assault, participants who responded better to trauma-focused CBT demonstrated decreased  
 21 functional connectivity between the amygdala and mid-posterior insula cortex during reappraisal  
 22 of negative images following treatment (Cisler et al., 2016). However, in a larger sample of adults  
 23 with PTSD, regional BOLD signal during a cognitive reappraisal task was not associated with  
 24 exposure-based psychotherapy outcomes (Fonzo et al., 2017). In conclusion, individual  
 25 differences in ability to successfully engage active emotion regulation strategies like reappraisal  
 26 may buffer against the deleterious effects of trauma exposure on mood, but are not necessarily  
 27 related to changes in anxiety and PTSD symptom levels.



## Reward responsivity

Responsivity to reward underlies key psychological factors linked to resilience such as optimism and positive emotionality (Feder et al., 2019). Resilient individuals show comparatively preserved ventral striatum response both when anticipating and receiving rewards, whereas PTSD and history of childhood maltreatment have been linked to anhedonia and blunted reward responses – particularly to social reward cues (Dillon et al., 2009; Hanson, Hariri, & Williamson, 2015; Hanson et al., 2016; Sailer et al., 2008; Elman et al., 2009; Nawijn et al., 2015). In a cross-sectional community-based sample of adolescents (a substantial proportion of whom had a history of severe childhood maltreatment), greater pallidal activation in response to positive images was associated with lower depression symptoms, and higher putamen activation was associated with lesser increase in depression two years later (Dennison et al., 2016). Among a large sample of university students ( $n = 820$ ) in the Duke Neurogenetics Study cohort, increased ventral striatal activation to both anticipatory and consummatory reward during a card guessing game significantly weakened the relationship between childhood trauma exposure and adult anhedonia – including when controlling for other depression symptoms and recent life stress (Corral-Frías et al., 2015).

In a cross-sectional study, trauma-exposed adults without PTSD had greater ventral striatal responses to happy faces compared to those with PTSD (Felmingham et al., 2014). Amongst individuals who felt negatively affected by the outcome of the 2016 US presidential election (and belonged to historically marginalized groups), higher nucleus accumbens BOLD signal and higher mPFC-accumbens connectivity during reward receipt attenuated the relationship between election-related distress and depression symptoms (Tashjian & Galván, 2018). In a longitudinal assessment of combat-exposed paramedics, decreased reward response in the nucleus accumbens post-, but not pre-exposure, was found to be related to self-reported PTSD symptoms – suggesting that preserved reward responses may be a marker of resilience, rather than a prospective indicator (Admon et al., 2013).

While greater reactivity to rewarding social/environmental cues could promote resilience, the resilience-reward responsivity association may be more complex. The longitudinal AURORA biotypes study (Stevens et al., 2021) identified two clusters associated with resilience in adults

1 scanned 2-3 weeks after seeking emergency department care for a serious injury or medical  
 2 condition. One (a “high reward reactivity” cluster with low threat reactivity and low inhibitory  
 3 control) did not replicate in the test cohort. The other, replicable, cluster had low reactivity to  
 4 both threat and reward, with higher hippocampal and vmPFC engagement during the inhibitory  
 5 control task, suggesting that resilience is associated with better inhibition of emotional reactivity  
 6 broadly. The association between resilience and lower general reactivity echoes an earlier small  
 7 cross-sectional study in Special Forces soldiers – considered highly resilient individuals –, who  
 8 showed lower differential nucleus accumbens and subgenual PFC responses to monetary  
 9 reward versus no-reward, compared to non-trauma-exposed civilians (Vythilingam et al., 2009).  
 10 These findings illustrate challenges for resilience research when trying to synthesize findings  
 11 across different populations – e.g., repeated occupational trauma exposures may differ from a  
 12 singular exposure, and certain occupations (like first responders) may select for traits like higher  
 13 sensation-seeking and higher reactivity threshold. For example, a recent, rigorous attempt to  
 14 replicate Stevens et al. (2021) AURORA biotypes identified threat- and reward-related clusters  
 15 in their cohort, but these were not identical to those in AURORA and not related to PTSD  
 16 vulnerability (Ben-Zion et al., 2023), potentially due to some differences in sample composition  
 17 and methods.

## 18 Cognitive flexibility and cognitive control

19 Resilient individuals engage prefrontal and hippocampal regions during executive  
 20 function tasks e.g., (Ben-Zion et al., 2018). Conversely, deficits in inhibition, working memory,  
 21 attentional control, and cognitive flexibility are observed in both adults with a history of childhood  
 22 maltreatment and individuals with a diagnosis of PTSD, not accounted for by comorbid  
 23 psychopathology, substance use, or history of traumatic brain injury (Gould et al., 2012; Scott et  
 24 al., 2015) (although the relationship between PTSD symptom severity and cognitive function  
 25 may be bidirectional; Jacob et al., 2019). In an early study, both resilient and non-trauma-  
 26 exposed healthy adults showed greater vlPFC, mPFC, and dlPFC activation during an inhibitory  
 27 control task, vs. those with PTSD (Falconer et al., 2008). Resilient individuals have also shown  
 28 greater dlPFC and superior frontal gyrus activation during an attentional control task with  
 29 emotional distractors, compared to both adults with PTSD and non-trauma-exposed controls  
 30 (Blair et al., 2013). Recently, both resilient and non-trauma-exposed adults (but not adults with

PTSD) exhibited significant decoupling between prefrontal regions and subcortical structures when attempting to suppress experimentally-induced memories (Mary et al., 2020): top-down prefrontal modulation of subcortical memory structures (hippocampus, parahippocampal gyrus, and precuneus) was associated with successful memory suppression, whereas the PTSD group showed bottom-up modulation of information flow. Indeed, greater whole-brain resting state hippocampal functional connectivity was a significant predictor of PTSD symptoms at six months in a prospective study in traumatically-injured adults recruited from the emergency department (Fitzgerald et al., 2022). Optimal prefrontal-subcortical cognitive control network function may facilitate resilience.

A longitudinal study in the Duke Neurogenetics Study cohort ( $n=120$ ) identified that greater neural responses to threat and reward predicted future increases in anxiety in university students with average or low but not *high* prefrontal activity during a working memory task, controlling for childhood maltreatment and stressful life events (Scully, Knodt, Radtke, Brigidi, & Hariri, 2019). These findings echo (Stevens et al., 2021) in suggesting that lower general reactivity and higher prefrontal inhibitory control could protect against adverse outcomes post-trauma exposure. Further, a longitudinal assessment of recently trauma-exposed adults found that increased inhibitory control-related hippocampal activation (at 1-2 months post-trauma) predicted decreased PTSD symptom severity at three and six months, and this finding was replicable (van Rooij et al., 2018). The hippocampus links cognition, memory and emotion (Speer & Delgado, 2017), thus is likely relevant to resilience to PTSD symptoms like alterations in mood, cognitive function, and intrusive thoughts and memories.

Finally, NCANDA cohort ( $n=392$ ) resting state fMRI showed that the association between childhood trauma severity and executive function difficulties in adolescents was mediated by lower functional connectivity between a network of the postcentral and precentral gyri, dorsal ACC, intraparietal sulcus, and anterior insula cortex – hub regions implicated in cognitive/behavioral control and sensorimotor integration. Preserved interconnectivity between the postcentral gyrus and dorsal ACC was related to lower reported executive dysfunction at baseline, and predicted lower likelihood of high-risk drinking 1-4 years later (Silveira et al., 2020). In addition to protecting against PTSD symptoms, preserved prefrontal function may reduce vulnerability to maladaptive coping strategies like substance use.

## Emerging areas of interest for resilience neuroimaging research

**Social support and social cognition.** Social support is vital for resilience to trauma exposure in both childhood and adulthood (van Harmelen et al., 2016; van Harmelen et al., 2017; Fritz, de Graaff, et al., 2018; Yule et al., 2019). Conversely, non-supportive social contexts like discrimination and rejection, impact emotion-responsive regions including the amygdala, insula, and ACC, even in the absence of (or after controlling for) associated psychological symptoms (Akdeniz et al., 2014; Clark, Miller, & Hegde, 2018). Supportive social environments may converge with resilience-promoting traits to upregulate perigenual ACC and other PFC regions during acute social (and non-social) threats (Holz et al., 2020). Social support also has a protective effect against prolonged stressors. For example, supportive parenting during childhood attenuates the link between childhood adversity (e.g., maltreatment, poverty) and neural responses to negative emotional faces in childhood (Wymbs et al., 2020), and resting state functional connectivity by adulthood within executive and emotion regulation networks (Brody et al., 2019). However, a longitudinal study in adolescents found no association between childhood adversity resilience (no DSM Axis-1 disorder diagnosis following significant family discord) and neural responses to rejection, although the authors suggest that the null finding might be due to low frequency of abuse and higher socioeconomic status of the sample (Fritz et al., 2019).

Higher levels of social cognitive function (ability to infer the thoughts of others and navigate the social world) may help foster adaptive outcomes following trauma, by conferring ability to recruit and maintain social ties (Stevens & Jovanovic, 2019; Lepore & Kliever, 2019). Hudson et al. (2018) administered a theory-of-mind task to women with childhood maltreatment exposure and unexposed controls. Women with history of childhood maltreatment without sub-clinical or clinical PTSD symptoms in adulthood showed greater functional connectivity between dorsomedial (dm)PFC and right temporoparietal junction (TPJ) during implicit theory of mind, whereas unexposed controls vs. women with PTSD symptoms showed similar dmPFC-right TPJ functional connectivity. Those with PTSD symptoms also demonstrated right TPJ *hypoactivation* during theory of mind, vs. unexposed controls. The TPJ and dmPFC represent nodes in the default network dorsomedial subsystem, involved in mentalizing and social processing (Spreng


& Andrews-Hanna, 2015). In summary, we need further research on proximal neural mechanisms underlying social support and social cognitive functioning in resilience. Considering the broader social environment would improve our understanding of mechanisms underlying resilience (**Box 2**; Biglan et al., 2012).

**Active coping.** Active coping focuses on approach-oriented behaviors for managing stressors, such as “*I take additional actions to get rid of the problem*” or “*I concentrate my effort on doing something about it*”), and is associated with lower cross-sectional PTSD symptoms (e.g., Stratta et al., 2015; Bistricky et al., 2019; Getnet et al., 2019). Further, there is preliminary evidence that a switch toward active coping accompanies response to cognitive-behavioral therapy for PTSD (Bourdon, El-Baalbaki, Girard, Lapointe-Blackburn, & Guay, 2019). mPFC function in resilience to prolonged stressors is implicated in individual differences in coping styles: healthy men with high early life stress but low trait rumination decreased amygdala and increased vmPFC perfusion during a mental arithmetic stress task, whereas men with high early life stress and high trait rumination showed the opposite (Wang, Paul, Stanton, Greeson, & Smoski, 2013). Healthy volunteers undergoing an acute stress paradigm exhibited initial vmPFC deactivation during early task runs, but. vmPFC signal recovery later in the task was correlated with self-reported active coping and anticorrelated with maladaptive coping behaviors (Sinha, Lacadie, Constable, & Seo, 2016). This reactivity-recovery pattern is consistent with preclinical evidence that prefrontal downregulation is adaptive during *acute* stress, but should change when stress is prolonged (Maier & Watkins, 2010; Sinha et al., 2016). However, a single laboratory session represents a very different timescale than naturalistic prolonged stress, and directionality of the relationship between mPFC recovery during chronic stress and coping style is unclear. Gender roles and social norms can also differentially reinforce active versus avoidant coping styles (McLean and Anderson, 2009; Street and Dardis, 2018; **Box 2**); interrelationships between these factors may warrant attention.

**Fear extinction.** It is important to update fear memories when transitioning from threat to safety, in order to respond appropriately to context (Lissek & van Meurs, 2015). vmPFC and hippocampus are key in fear extinction maintenance, in both human and animal studies (Milad et al., 2007; Fullana et al., 2018). Enhanced extinction learning is associated with surprise-related learning signals in the vmPFC, and amygdala connectivity with a ventral mPFC subregion (Dunsmoor et al., 2019); the same circuitry might be critical for extinction learning as

1 an interventional mechanism (Fullana et al., 2020). Indeed, greater success of prolonged  
2 exposure therapy for PTSD – which promotes extinction learning – is linked to pre-treatment  
3 neural responses to emotion, including higher vmPFC signal during emotional regulation, greater  
4 magnitude of PFC responses while viewing fearful faces, and greater inhibition of the amygdala  
5 by lateral PFC transcranial magnetic stimulation (Fonzo et al., 2017). Further, vmPFC activity  
6 has been demonstrated to be central to the success of *imagined* fear extinction, which may be  
7 particularly relevant to the psychotherapeutic context, as in-vivo exposure is not always feasible  
8 (Reddan, Wager, & Schiller, 2018). Thus, vmPFC recruitment during emotional contexts may be  
9 important to later success of exposure-based treatment strategies – and may play a role in  
10 successful maintenance of extinction learning broadly.

11 Resting state findings support the relevance of vmPFC recruitment. Occupational trauma-  
12 exposed firefighters with greater insula-vmPFC functional connectivity had fewer PTSD  
13 symptoms, although firefighters had greater insula functional connectivity with other fear circuitry  
14 regions vs. healthy non-firefighter adults, regardless of PTSD symptoms (Jeong et al., 2019).  
15 Similarly, typhoon survivors with PTSD showed greater vmPFC connectivity with the basolateral  
16 amygdala, relative to trauma-exposed controls. The basolateral amygdala has a central role in  
17 fear learning and emotional responding. This latter study illustrates the difficulty in inferring a  
18 directional relationship between brain regions from most functional neuroimaging studies.  
19 However, a resting state study, using Granger causality, in typhoon survivors and unexposed  
20 healthy adults found that both individuals with PTSD and trauma-exposed controls exhibited  
21 greater amygdala-to-mPFC effective connectivity vs. unexposed controls, but inhibitory mPFC-  
22 to-amygdala connectivity was only observed in trauma-exposed controls. (Chen et al., 2018b).



**Figure 1.** Brain regions most commonly associated with higher resilience to trauma, organized by psychological factor (including emerging areas of interest) and underlying neural circuitry, based on review of fMRI BOLD activation literature.  
*Note.* All figures use parcellations from the Desikan-Killiany Cortical Atlas parcellation or Freesurfer's automatic subcortical segmentation, implemented with `ggseg()` and `ggsegExtra()` packages for R (Mowinckel & Vidal-Piñeiro, 2019, n.d.), except for the subcortical Reward Responsivity plot (approximate location of the nucleus accumbens drawn by hand) and the cortical Social Support & Cognition plot (which uses the AAL2 parcellation). ACC = anterior cingulate, PFC = prefrontal cortex, d = dorsal, dl = dorsolateral, m = medial, l = lateral, r = rostral, v = ventral, vm = ventromedial. Upward arrows indicate regions of greater activation in resilient individuals vs. others; downward arrows indicate regions of lower activation in resilient individuals vs. others; and both arrows together indicate mixed findings.

## Discussion

Here, we summarized task-based and resting state fMRI studies of resilience to childhood and adulthood trauma, focusing on psychological factors widely linked to resilient responses to severe adversity and trauma: effective emotion regulation, reward responsiveness, and cognitive control. We also reviewed findings related to social cognition, fear extinction, and active coping, as emerging areas of interest in resilience research (**Figure 1**). Across studies, we found robust evidence for preserved mPFC function in downregulation of limbic responses to emotional stimuli in resilient individuals. This is consistent with evidence from animal and human experimental models implicating mPFC engagement in adaptive coping with prolonged stress (Maier & Watkins, 2010). Chronic social defeat stress-susceptible mice showing social

impairment exhibit selective reductions in vmPFC spike frequencies – suggesting a potential role of preserved vmPFC function in adaptive *social* behaviour following chronic stress (Abe, Okada, Nakayama, Ikegaya, & Sasaki, 2019), in line with emerging human evidence for perigenual cingulate and prefrontal function as mediators of social support's positive effects on responses to stress and pain (Holz et al., 2020). However, we lack knowledge of neural correlates of social functioning and social cognition *in resilient humans* despite their role in fostering positive outcomes (Stevens & Jovanovic, 2019). Considering the broader social environment is also critical for understanding the brain-environment interactions underlying resilience (**Box 2**).

Preserved reward signal processing in the ventral striatum has been linked to resilience to anhedonia and depression symptoms. One possibility is that preserved reward system function promotes participation in social and other rewarding experiences, thereby buffering the impacts of stress. Intact reward circuitry could also contribute to positive prospection – e.g., a hippocampal-midbrain-vmPFC circuit has recently been implicated in imagined future positive outcomes (Ikegaya et al., 2019). However, reward system findings have been mixed; resilient people may be less reactive both to reward and threat. Lesser reactivity might allow for greater stability in a changing environment, for example via lower emotional lability or greater ability to remain focused on goals by inhibiting salient yet distracting cues (e.g., Stevens et al., 2021).

Cognitive control-related brain activation predicts positive outcomes following trauma, specifically frontal (dlPFC) and hippocampal. Adult hippocampal neurogenesis plays an important role in cognitive flexibility, which may enable individuals to disengage from trauma-related memories and thoughts, and decrease cardinal PTSD symptoms such as intrusive thoughts and flashbacks (Aupperle, Melrose, Stein, & Paulus, 2012; Anacker & Hen, 2017). Prefrontal-subcortical connectivity (including hippocampal) in cognitive control appears important for suppressing unwanted memories (Mary et al., 2020). Although perhaps not specific to PTSD, activity in this circuit may represent an important factor in vulnerability to PTSD and maladaptive coping mechanisms (e.g., substance use problems) following trauma (Ersche, 2020). Recent work in this area highlights the importance of considering within-person interactions between resilience-relevant factors – specifically, how higher levels of prefrontally-mediated executive function may 'rescue' risk for anxiety in individuals who are more reactive to threat and less responsive to rewards (Scully et al., 2019). We note that the brain regions



highlighted across the resilience-linked psychological factors also broadly correspond to those described in recent models of psychological *vulnerability* to trauma (e.g., Pitman et al., 2012; Patel et al., 2012; Yehuda et al., 2015).

## Longitudinal findings

Given that resilience inherently involves a temporal relationship between stressor and response, large longitudinal studies are better able to answer key questions – for example, are the differences between resilient people and those who develop a psychiatric disorder present pre-exposure, or do the differences emerge afterward in a compensatory manner? Can we identify people at greater risk of negative consequences at the time of the event or soon after, and intervene early to prevent PTSD and related disorders? Several of the studies reviewed here illustrate the potential of longitudinal, cohort-based neuroimaging research. Though their samples differ in terms of trauma exposure type and developmental stage, the work by Scult et al. (2019) in the Duke Neurogenetics cohort, Silveira et al. (2020) in the NCANDA cohort, and Stevens et al. (2021) in the AURORA cohort suggest that greater general bottom-up neural reactivity to emotional stimuli – both positive and negative – is linked to poorer outcomes unless accompanied by inhibitory hippocampal and prefrontal engagement.

## Limitations

It is important to clarify the scope of inference we can draw from the current data. There are too few studies to differentiate results by developmental stage at the time of the traumatic event and/or the study. In addition to differences in brain development, another difference between studies of adults and youth is that the measures used in adult studies typically focus on symptoms (psychopathology), whereas youth studies tend to operationalize resilience more broadly (**Box 2**). Many studies are retrospective and cross-sectional. Only longitudinal studies can tease out whether differences seen in resilient individuals represent pre-existing vulnerabilities or adaptations. There are also methodological weaknesses, including small sample sizes and low power (Szucs & Ioannidis, 2020). Many older studies used liberal multiple comparison corrections, which increases risk for type 1 errors (Eklund, Nichols, & Knutsson, 2016; Cox, Chen, Glen, Reynolds, & Taylor, 2017). However, some key findings replicate in well-powered samples (e.g., (Corral-Frías et al., 2015; McLean et al., 2020; Silveira et al., 2020) – with the caveat that these findings may not necessarily be robust to cohort and methodological

differences between studies (Ben-Zion et al., 2023). Finally, identifying robust neuroimaging biomarkers requires establishing intrapersonal reliability of fMRI measures (Nord, Gray, Charpentier, Robinson, & Roiser, 2017; Elliott et al., 2019, 2020). Multivariate approaches exploiting the high dimensionality of neuroimaging data may be more appropriate (Dubois & Adolphs, 2016; Finn et al., 2020).

## Conclusions and future directions

Real-world settings have started to translate neuroscientific findings to interventions (Greenberg, 2006; Waugh & Koster, 2015; Keynan et al., 2019). Examining neural correlates of treatment response may illuminate neural resilience mechanisms ‘activated’ by successful treatment. Ongoing and future longitudinal cohort studies with a peritraumatic baseline, such as AURORA (McLean et al., 2020), along with additional consideration of interindividual sociodemographic differences (**Box 3**), will contribute to further understanding resilience to severe adversity and trauma.

[Box 3 around here]

### Box 1: Quantifying trauma

Assessing the severity of different traumatic experiences is not a trivial task. Indeed, to a large extent, the degree to which any event is traumatic is determined by the psychological impact on the person or people involved (Green, 1990). However, such a definition poses some difficulties for studies of resilience, which aim to investigate individual differences in response to similarly adverse events. Studying trauma-exposed samples optimally involves several considerations, including severity of trauma exposure and chronicity, and in the case of childhood maltreatment, developmental stage at the time of trauma exposure (Harpur, Polek, & van Harmelen, 2015; Dunn et al., 2019; Gee, 2020).

Events sufficient to fulfill criteria for a diagnosis of post-traumatic stress disorder (PTSD) have been codified within the American Psychiatric Association's Diagnostic and Statistical Manual (DSM) as 'Category A' events – specifically, in DSM-5, as being “exposed to death, threatened death, actual or threatened serious injury, or actual or threatened sexual violence” (American Psychiatric Association, 2013). In the World Health Organization's International Classification of Diseases (ICD-10), the requirement is simply that a person has experienced an event or situation “of exceptionally threatening or catastrophic nature, which would be likely to cause pervasive distress in almost anyone” (World Health Organization, 1993). Under either scheme, presence of such an event in an individual's history is usually probed using a structured clinical interview (e.g., Weathers et al., 2013).

Experience of childhood trauma in younger research participants may be assessed using developmentally sensitive interviews with a child's primary caregiver (e.g., Goodyer et al., 2010). In comparison, in studies involving adult participants, experience of childhood trauma is often assessed using retrospective questionnaires, such as the Childhood Trauma Questionnaire (Bernstein et al., 1994). Notably, a recent meta-analysis reported poor agreement between prospective and retrospective measures of childhood maltreatment (Baldwin, Reuben, Newbury, & Danese, 2019). Although both prospective and retrospective measures identify groups of at-risk individuals, they appear to highlight largely non-overlapping sets of people – meaning that it may not be valid to assume the same risk mechanisms apply to individuals identified using different methods.

Further, although both intensity and chronicity are important dimensions of trauma, it is not always clear how to combine these different aspects of adverse event exposure in a standardized way. Some studies have addressed this issue by restricting recruitment to individuals exposed to a specific precipitating event, for example in volunteers from the World Trade Center rescue and recovery worker cohort (Pietrzak, Feder, et al., 2014), or individuals with combat trauma (Keane et al., 1989). However, even in such samples, there is likely to be additional lifetime trauma exposure in many participants. One approach to this problem is to use a data reduction technique such as principal components analysis (PCA) to combine several different continuous measures into a single index of trauma exposure severity (e.g., to combine different aspects of childhood family experience; van Harmelen et al., 2017). It is important to bear in mind when synthesizing data in this way that there may also be trauma-specific considerations to take into account during analysis – for example when considering responses to differently gendered angry faces in people exposed to intimate partner violence (Fonzo et al., 2010).

## Box 2: Quantifying resilience

**In trauma-exposed populations.** In neurobiological studies with adult participants, resilience has often been defined based on the absence of a DSM Axis-1 disorder diagnosis following significant trauma exposure. Such categorical analyses may be complemented by dimensional approaches that relate measures to current severity scores for PTSD, depression, or anxiety symptoms. However, it is increasingly acknowledged that symptom sum scores for specific diagnoses are not the only key outcome variable in studies of resilience. Indeed, trauma-related psychopathology is highly heterogeneous, and different symptom dimensions may differentially relate to overall burden of disability (e.g., Pietrzak, el-Gabalawy, et al., 2014) – suggesting more nuanced approaches may be warranted. Conversely, studies of the effects of childhood maltreatment have tended to take a more holistic approach towards assessing resilience, including a greater focus on psychosocial outcomes, and examining functioning across multiple domains (McGloin & Widom, 2001; Cicchetti & Rogosch, 2012). Recent examples of this approach across the literature include examining resilience across different functional outcome categories in children (Burt et al., 2016), analysing disruption to work, social, and family life in addition to PTSD symptomatology in trauma-exposed adults (Horn et al., 2016), and the use of PCA-derived cross-domain psychosocial functioning scores in an adolescent cohort (A. L. van Harmelen et al., 2017).

The notion of good functioning invoked by definitions of resilience is *environment-dependent*: adaptive behaviour in violent, volatile, or resource-poor environments may not conform to normative accounts of behaviour in more stable or resource-rich environments (Luthar, Sawyer, & Brown, 2006). Further, psychosocial environments may differ across individuals within a shared wider context – e.g. for individuals who identify as different genders and/or belong to different racial or ethnic groups (Tolin & Foa, 2008; Portnoy et al., 2018; Street & Dardis, 2018; Brondolo, 2015). For example, differences in resilience between men and women have been found to be explained by differences in trauma type exposure across genders (particularly sexual violence; Tolin & Foa, 2008; Portnoy et al., 2018), and socialized gender norms may further contribute to differences in chronic environmental stress exposure and propensity to engage in less adaptive coping strategies (Street & Dardis, 2018). In particular, racism is an additional source of trauma in minoritized people that may be under-recognized by current clinical assessments (Carter, 2007; Williams, Metzger, Leins, & DeLapp, 2018). For example, previous experiences of racial discrimination has recently been shown to add significant risk for PTSD symptom development following traumatic injury (Bird et al., 2021).

Importantly, the expected level of psychosocial functioning for a given individual should take into account their lifetime trauma burden (Karam et al., 2014; Feder et al., 2016). Various researchers have therefore proposed that resilience can best be quantified by regressing metrics of psychosocial functioning against an appropriate measure of trauma exposure severity: such that positive residuals from this model represent better than expected ('resilient') outcomes, and negative residuals represent worse than expected ('vulnerable') outcomes – compared to what would be predicted based on the group as a whole (Amstadter, Myers, & Kendler, 2014; Amstadter, Maes, Sheerin, Myers, & Kendler, 2016; A. L. van Harmelen et al., 2017; Ioannidis, Askelund, Kievit, & van Harmelen, 2020). Resilience-promoting factors can then be described as any resource (biological, psychological, social/socioeconomic) that decreases the risk of poor outcomes following adverse circumstances. Resilience-promoting factors may include both protective factors that help buffer the impact of stress, and resources that are able to foster positive compensatory changes following trauma exposure (Luthar et al., 2006; Schultze-Lutter, Schimmelmann, & Schmidt, 2016). Critically, resilience-promoting factors are not independently additive, but interact in complex ways (Luthar et al., 2006; Fritz, Fried et al., 2018; Liebenberg, 2020). It is therefore vital not to study such factors in isolation, but rather in the context of each other – ideally within the same individuals.

**In the general population.** Some researchers have also examined population variation in self-reported trait resilience. For example, the Connor-Davidson Resilience Scale (CD-RISC) probes how likely individuals are to endorse statements such as "*I am able to adapt to change*" and "*I tend to bounce back after illness or hardship*" (Connor & Davidson, 2003b). There is somewhat equivocal evidence regarding the relationship between trait resilience and outcomes following trauma exposure (Daniels et al., 2012; Powers et al., 2014). Interestingly, a recent study of emergency department attendees found that a negative association between CD-RISC score close to admission and PTSD symptoms six months in the future was mediated by lower social withdrawal in higher trait resilience individuals – suggesting that the impacts of trait resilience on functional outcome may be via increased ability to maintain or recruit social support (N. J. Thompson, Fiorillo, Rothbaum, Ressler, & Michopoulos, 2018). Further, most questionnaire measures of resilience relate to a specific conceptualisation of resilience that focuses

on individual ‘grit’ or ‘hardiness’ and that may not translate well to non-Western cultural settings – particularly those that emphasise the role of communities rather than individuals in fostering resilience (Meili & Maercker, 2019; Mendenhall & Kim, 2019). An alternative approach for studies of resilience in the general population is to study the *mechanisms* underlying resilience based on experimental intervention data – for example as reflected in lower self-reported fear or physiological reactivity during stress induction paradigms.

### Box 3: Considerations for future work

Studying multiple potential resilience-promoting factors within the same individual is necessary to tease apart how putative neurobiological resilience factors interact both with each other and with other sociodemographic factors known to affect resilience to trauma-related psychopathology. We would encourage greater consideration of an individual’s social, cultural, and socioeconomic environment in future studies of neural mechanisms related to resilience.

The trade-off for high power in large cohort studies is often the selective pressure on study measures (for inclusion, brevity, and ease of administration). However, accurate assessment of both resilience and the sociodemographic factors described above requires in-depth clinical phenotyping and trauma history screening, often by clinical interview with appropriately trained study personnel. To build our understanding of the neural correlates of resilience, future work should aim to hit a sweet spot in this trade-off between appropriate power and depth of individual phenotyping.

An important next step will be to integrate neuroimaging metrics related to resilience with other biological measurements relevant to both individual differences in neural function and environmental exposure. Novel analytic approaches such as the calculation of polygenic risk scores, or quantification of gene co-expression modules may yield sufficient power in (single site) achievable imaging sample sizes to link neural correlates of psychological constructs to underlying biology (Dima & Breen, 2015; Bogdan et al., 2017). Greater integration of neuroimaging data into such pathways should improve our ability to interpret existing findings in terms of underlying molecular mechanisms. The cost of well-powered samples in neuroimaging genetics is typically prohibitively expensive for a single researcher or work group, but large-scale multi-site efforts e.g., the international ENIGMA [Enhancing NeuroImaging Genetics through Meta Analysis] Consortium have been extremely successful in facilitating robust and reproducible research, and will continue to be a tremendous resource (Thompson et al., 2020).

## 1 References

- 2 Abe, R., Okada, S., Nakayama, R., Ikegaya, Y., & Sasaki, T. (2019). Social defeat stress causes  
3 selective attenuation of neuronal activity in the ventromedial prefrontal cortex. *Scientific*  
4 *Reports*, 9(1), 9447. <https://doi.org/10.1038/s41598-019-45833-5>
- 5 Admon, R., Lubin, G., Rosenblatt, J. D., Stern, O., Kahn, I., Assaf, M., & Hendler, T. (2013).  
6 Imbalanced Neural Responsivity to Risk and Reward Indicates Stress Vulnerability in Humans.  
7 *Cerebral Cortex*, 23(1), 28–35. <https://doi.org/10.1093/cercor/bhr369>
- 8 Admon, R., Lubin, G., Stern, O., Rosenberg, K., Sela, L., Ben-Ami, H., & Hendler, T. (2009). Human  
9 vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity.  
10 *Proceedings of the National Academy of Sciences of the United States of America*, 106(33),  
11 14120–14125. <https://doi.org/10.1073/pnas.0903183106>
- 12 Akdeniz, C., Tost, H., Streit, F., Haddad, L., Wüst, S., Schäfer, A., ... Meyer-Lindenberg, A. (2014).  
13 Neuroimaging Evidence for a Role of Neural Social Stress Processing in Ethnic Minority–  
14 Associated Environmental Risk. *JAMA Psychiatry*, 71(6), 672–680.  
15 <https://doi.org/10.1001/jamapsychiatry.2014.35>
- 16 American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (Fifth  
17 Edition). Washington, DC and London, England: American Psychiatric Association.  
18 <https://doi.org/10.1176/appi.books.9780890425596>
- 19 Amstadter, A. B., Maes, H. H., Sheerin, C. M., Myers, J. M., & Kendler, K. S. (2016). The relationship  
20 between genetic and environmental influences on resilience and on common internalizing and  
21 externalizing psychiatric disorders. *Social Psychiatry and Psychiatric Epidemiology*, 51(5), 669–  
22 678. <https://doi.org/10.1007/s00127-015-1163-6>
- 23 Amstadter, A. B., Myers, J. M., & Kendler, K. S. (2014). Psychiatric resilience: Longitudinal twin study.  
24 *The British Journal of Psychiatry: The Journal of Mental Science*, 205(4), 275–280.  
25 <https://doi.org/10.1192/bjp.bp.113.130906>

- 1 Anacker, C., & Hen, R. (2017). Adult hippocampal neurogenesis and cognitive flexibility—Linking  
2 memory and mood. *Nature Reviews Neuroscience*, 18(6), 335–346.  
3 <https://doi.org/10.1038/nrn.2017.45>
- 4 Aupperle, R. L., Melrose, A. J., Stein, M. B., & Paulus, M. P. (2012). Executive function and PTSD:  
5 Disengaging from trauma. *Neuropharmacology*, 62(2), 686–694.  
6 <https://doi.org/10.1016/j.neuropharm.2011.02.008>
- 7 Baldwin, J. R., Reuben, A., Newbury, J. B., & Danese, A. (2019). Agreement Between Prospective and  
8 Retrospective Measures of Childhood Maltreatment: A Systematic Review and Meta-analysis.  
9 *JAMA Psychiatry*, 76(6), 584. <https://doi.org/10.1001/jamapsychiatry.2019.0097>
- 10 Ben-Zion, Z., Fine, N. B., Keynan, N. J., Admon, R., Green, N., Halevi, M., ... Shalev, A. Y. (2018).  
11 Cognitive Flexibility Predicts PTSD Symptoms: Observational and Interventional Studies.  
12 *Frontiers in Psychiatry*, 9, 477. <https://doi.org/10.3389/fpsy.2018.00477>
- 13 Ben-Zion, Z., Spiller, T. R., Keynan, J. N., Admon, R., Levy, I., Liberzon, I., ... Harpaz-Rotem, I. (2023).  
14 Evaluating the Evidence for Brain-Based Biotypes of Psychiatric Vulnerability in the Acute  
15 Aftermath of Trauma. *American Journal of Psychiatry*, appi.ajp.20220271.  
16 <https://doi.org/10.1176/appi.ajp.20220271>
- 17 Bernstein, D. P., Fink, L., Handelsman, L., Foote, J., Lovejoy, M., Wenzel, K., ... Ruggiero, J. (1994).  
18 Initial reliability and validity of a new retrospective measure of child abuse and neglect. *The*  
19 *American Journal of Psychiatry*, 151(8), 1132–1136. <https://doi.org/10.1176/ajp.151.8.1132>
- 20 Biglan, A., Flay, B. R., Embry, D. D., & Sandler, I. N. (2012). The Critical Role of Nurturing  
21 Environments for Promoting Human Wellbeing. *The American Psychologist*, 67(4), 257–271.  
22 <https://doi.org/10.1037/a0026796>
- 23 Bird, C. M., Webb, E. K., Schramm, A. T., Torres, L., Larson, C., & deRoos-Cassini, T. A. (2021).  
24 Racial Discrimination is Associated with Acute Posttraumatic Stress Symptoms and Predicts  
25 Future Posttraumatic Stress Disorder Symptom Severity in Trauma-Exposed Black Adults in the  
26 United States. *Journal of Traumatic Stress*. <https://doi.org/10.1002/jts.22670>

- 1 Bistricky, S. L., Long, L. J., Lai, B. S., Gallagher, M. W., Kanenberg, H., Elkins, S. R., ... Short, M. B.  
2 (2019). Surviving the storm: Avoidant coping, helping behavior, resilience and affective  
3 symptoms around a major hurricane-flood. *Journal of Affective Disorders*, 257, 297–306.  
4 <https://doi.org/10.1016/j.jad.2019.07.044>
- 5 Blair, K. S., Vythilingam, M., Crowe, S. L., McCaffrey, D. E., Ng, P., Wu, C. C., ... Blair, R. J. R. (2013).  
6 Cognitive control of attention is differentially affected in trauma-exposed individuals with and  
7 without post-traumatic stress disorder. *Psychological Medicine*, 43(1), 85–95.  
8 <https://doi.org/10.1017/S0033291712000840>
- 9 Bogdan, R., Salmeron, B. J., Carey, C. E., Agrawal, A., Calhoun, V. D., Garavan, H., ... Goldman, D.  
10 (2017). Imaging Genetics and Genomics in Psychiatry: A Critical Review of Progress and  
11 Potential. *Biological Psychiatry*, 82(3), 165–175. <https://doi.org/10.1016/j.biopsych.2016.12.030>
- 12 Bourdon, D.-É., El-Baalbaki, G., Girard, D., Lapointe-Blackburn, É., & Guay, S. (2019). Schemas and  
13 coping strategies in cognitive-behavioral therapy for PTSD: A systematic review. *European*  
14 *Journal of Trauma & Dissociation*, 3(1), 33–47. <https://doi.org/10.1016/j.ejtd.2018.09.005>
- 15 Brody, G. H., Yu, T., Nusslock, R., Barton, A. W., Miller, G. E., Chen, E., ... Sweet, L. H. (2019). The  
16 Protective Effects of Supportive Parenting on the Relationship Between Adolescent Poverty and  
17 Resting-State Functional Brain Connectivity During Adulthood. *Psychological Science*, 30(7),  
18 1040–1049. <https://doi.org/10.1177/0956797619847989>
- 19 Bromet, E. J., Karam, E. G., Koenen, K. C., & Stein, D. J. (2018). *Trauma and Posttraumatic Stress*  
20 *Disorder: Global Perspectives from the WHO World Mental Health Surveys*. Cambridge  
21 University Press.
- 22 Brondolo, E. (2015). Racial and Ethnic Disparities in Health: Examining the Contexts That Shape  
23 Resilience and Risk. *Psychosomatic Medicine*, 77(1), 2–5.  
24 <https://doi.org/10.1097/PSY.0000000000000149>



- 1 Burt, K. B., Whelan, R., Conrod, P. J., Banaschewski, T., Barker, G. J., Bokde, A. L., ... Garavan, H.  
2 (2016). Structural brain correlates of adolescent resilience. *Journal of Child Psychology and*  
3 *Psychiatry, and Allied Disciplines*, 57(11), 1287–1296. <https://doi.org/10.1111/jcpp.12552>
- 4 Callaghan, B. L., Gee, D. G., Gabard-Durnam, L., Telzer, E. H., Humphreys, K. L., Goff, B., ...  
5 Tottenham, N. (2019). Decreased Amygdala Reactivity to Parent Cues Protects Against Anxiety  
6 Following Early Adversity: An Examination Across 3 Years. *Biological Psychiatry: Cognitive*  
7 *Neuroscience and Neuroimaging*, 4(7), 664–671. <https://doi.org/10.1016/j.bpsc.2019.02.001>
- 8 Careaga, M. B. L., Girardi, C. E. N., & Suchecki, D. (2016). Understanding posttraumatic stress  
9 disorder through fear conditioning, extinction and reconsolidation. *Neuroscience and*  
10 *Biobehavioral Reviews*, 71, 48–57. <https://doi.org/10.1016/j.neubiorev.2016.08.023>
- 11 Carter, R. T. (2007). Racism and Psychological and Emotional Injury: Recognizing and Assessing  
12 Race-Based Traumatic Stress. *The Counseling Psychologist*, 35(1), 13–105.  
13 <https://doi.org/10.1177/0011000006292033>
- 14 Chen, F., Ke, J., Qi, R., Xu, Q., Zhong, Y., Liu, T., ... Lu, G. (2018a). Increased Inhibition of the  
15 Amygdala by the mPFC may Reflect a Resilience Factor in Post-traumatic Stress Disorder: A  
16 Resting-State fMRI Granger Causality Analysis. *Frontiers in Psychiatry*, 9, 516.  
17 <https://doi.org/10.3389/fpsy.2018.00516>
- 18 Chen, F., Ke, J., Qi, R., Xu, Q., Zhong, Y., Liu, T., ... Lu, G. (2018b). Increased Inhibition of the  
19 Amygdala by the mPFC may Reflect a Resilience Factor in Post-traumatic Stress Disorder: A  
20 Resting-State fMRI Granger Causality Analysis. *Frontiers in Psychiatry*, 9, 516.  
21 <https://doi.org/10.3389/fpsy.2018.00516>
- 22 Cicchetti, D., & Rogosch, F. A. (2012). Gene × Environment interaction and resilience: Effects of child  
23 maltreatment and serotonin, corticotropin releasing hormone, dopamine, and oxytocin genes.  
24 *Development and Psychopathology*, 24(2), 411–427.  
25 <https://doi.org/10.1017/S0954579412000077>

- 1 Cisler, J. M., Sigel, B. A., Steele, J. S., Smitherman, S., Vanderzee, K., Pemberton, J., ... Kilts, C. D.  
2 (2016). Changes in functional connectivity of the amygdala during cognitive reappraisal predict  
3 symptom reduction during trauma-focused cognitive-behavioral therapy among adolescent girls  
4 with post-traumatic stress disorder. *Psychological Medicine*, 46(14), 3013–3023.  
5 <https://doi.org/10.1017/S0033291716001847>
- 6 Clark, U. S., Miller, E. R., & Hegde, R. R. (2018). Experiences of Discrimination Are Associated With  
7 Greater Resting Amygdala Activity and Functional Connectivity. *Biological Psychiatry: Cognitive*  
8 *Neuroscience and Neuroimaging*, 3(4), 367–378. <https://doi.org/10.1016/j.bpsc.2017.11.011>
- 9 Connor, K. M., & Davidson, J. R. T. (2003a). Development of a new resilience scale: The Connor-  
10 Davidson Resilience Scale (CD-RISC). *Depression and Anxiety*, 18(2), 76–82.  
11 <https://doi.org/10.1002/da.10113>
- 12 Connor, K. M., & Davidson, J. R. T. (2003b). Development of a new resilience scale: The Connor-  
13 Davidson Resilience Scale (CD-RISC). *Depression and Anxiety*, 18(2), 76–82.  
14 <https://doi.org/10.1002/da.10113>
- 15 Corral-Frías, N. S., Nikolova, Y. S., Michalski, L. J., Baranger, D. A. A., Hariri, A. R., & Bogdan, R.  
16 (2015). Stress-related anhedonia is associated with ventral striatum reactivity to reward and  
17 transdiagnostic psychiatric symptomatology. *Psychological Medicine*, 45(12), 2605–2617.  
18 <https://doi.org/10.1017/S0033291715000525>
- 19 Cox, R. W., Chen, G., Glen, D. R., Reynolds, R. C., & Taylor, P. A. (2017). FMRI Clustering in AFNI:  
20 False-Positive Rates Redux. *Brain Connectivity*, 7(3), 152–171.  
21 <https://doi.org/10.1089/brain.2016.0475>
- 22 Daniels, J. K., Hegadoren, K. M., Coupland, N. J., Rowe, B. H., Densmore, M., Neufeld, R. W. J., &  
23 Lanius, R. A. (2012). Neural correlates and predictive power of trait resilience in an acutely  
24 traumatized sample: A pilot investigation. *The Journal of Clinical Psychiatry*, 73(3), 327–332.  
25 <https://doi.org/10.4088/JCP.10m06293>

- 1 Demers, L. A., Hunt, R. H., Cicchetti, D., Cohen-Gilbert, J. E., Rogosch, F. A., Toth, S. L., & Thomas,  
2 K. M. (2021). Impact of childhood maltreatment and resilience on behavioral and neural patterns  
3 of inhibitory control during emotional distraction. *Development and Psychopathology*, 1–12.  
4 <https://doi.org/10.1017/S0954579421000055>
- 5 Demers, L. A., McKenzie, K. J., Hunt, R. H., Cicchetti, D., Cowell, R. A., Rogosch, F. A., ... Thomas, K.  
6 M. (2018). Separable Effects of Childhood Maltreatment and Adult Adaptive Functioning on  
7 Amygdala Connectivity During Emotion Processing. *Biological Psychiatry: Cognitive*  
8 *Neuroscience and Neuroimaging*, 3(2), 116–124. <https://doi.org/10.1016/j.bpsc.2017.08.010>
- 9 Denckla, C. A., Cicchetti, D., Kubzansky, L. D., Seedat, S., Teicher, M. H., Williams, D. R., & Koenen,  
10 K. C. (2020). Psychological resilience: An update on definitions, a critical appraisal, and  
11 research recommendations. *European Journal of Psychotraumatology*, 11(1), 1822064.  
12 <https://doi.org/10.1080/20008198.2020.1822064>
- 13 Dennison, M. J., Sheridan, M. A., Busso, D. S., Jenness, J. L., Peverill, M., Rosen, M. L., &  
14 McLaughlin, K. A. (2016). Neurobehavioral markers of resilience to depression amongst  
15 adolescents exposed to child abuse. *Journal of Abnormal Psychology*, 125(8), 1201–1212.  
16 <https://doi.org/10.1037/abn0000215>
- 17 Dillon, D. G., Holmes, A. J., Birk, J. L., Brooks, N., Lyons-Ruth, K., & Pizzagalli, D. A. (2009). Childhood  
18 Adversity Is Associated with Left Basal Ganglia Dysfunction During Reward Anticipation in  
19 Adulthood. *Biological Psychiatry*, 66(3), 206–213.  
20 <https://doi.org/10.1016/j.biopsych.2009.02.019>
- 21 Dima, D., & Breen, G. (2015). Polygenic risk scores in imaging genetics: Usefulness and applications.  
22 *Journal of Psychopharmacology*, 29(8), 867–871. <https://doi.org/10.1177/0269881115584470>
- 23 Dubois, J., & Adolphs, R. (2016). Building a Science of Individual Differences from fMRI. *Trends in*  
24 *Cognitive Sciences*, 20(6), 425–443. <https://doi.org/10.1016/j.tics.2016.03.014>
- 25 Dunn, E. C., Soare, T. W., Zhu, Y., Simpkin, A. J., Suderman, M. J., Klengel, T., ... Relton, C. L.  
26 (2019). Sensitive Periods for the Effect of Childhood Adversity on DNA Methylation: Results

- 1 From a Prospective, Longitudinal Study. *Biological Psychiatry*, 85(10), 838–849.
- 2 <https://doi.org/10.1016/j.biopsych.2018.12.023>
- 3 Dunsmoor, J. E., Kroes, M. C. W., Li, J., Daw, N. D., Simpson, H. B., & Phelps, E. A. (2019). Role of
- 4 Human Ventromedial Prefrontal Cortex in Learning and Recall of Enhanced Extinction. *Journal*
- 5 *of Neuroscience*, 39(17), 3264–3276. <https://doi.org/10.1523/JNEUROSCI.2713-18.2019>
- 6 Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for spatial extent
- 7 have inflated false-positive rates. *Proceedings of the National Academy of Sciences*,
- 8 201602413. <https://doi.org/10.1073/pnas.1602413113>
- 9 Elliott, M. L., Knodt, A. R., Cooke, M., Kim, M. J., Melzer, T. R., Keenan, R., ... Hariri, A. R. (2019).
- 10 General functional connectivity: Shared features of resting-state and task fMRI drive reliable and
- 11 heritable individual differences in functional brain networks. *NeuroImage*, 189, 516–532.
- 12 <https://doi.org/10.1016/j.neuroimage.2019.01.068>
- 13 Elliott, M. L., Knodt, A. R., Ireland, D., Morris, M. L., Poulton, R., Ramrakha, S., ... Hariri, A. R. (2020).
- 14 What Is the Test-Retest Reliability of Common Task-Functional MRI Measures? New Empirical
- 15 Evidence and a Meta-Analysis. *Psychological Science*, 31(7), 792–806.
- 16 <https://doi.org/10.1177/0956797620916786>
- 17 Elman, I., Lowen, S., Frederick, B. B., Chi, W., Becerra, L., & Pitman, R. K. (2009). Functional
- 18 Neuroimaging of Reward Circuitry Responsivity to Monetary Gains and Losses in Posttraumatic
- 19 Stress Disorder. *Biological Psychiatry*, 66(12), 1083–1090.
- 20 <https://doi.org/10.1016/j.biopsych.2009.06.006>
- 21 Ersche, K. D. (2020). Resilience to trauma: Just a matter of control? *Science*, 367(6479), 734–735.
- 22 <https://doi.org/10.1126/science.aaz9451>
- 23 Falconer, E., Bryant, R., Felmingham, K. L., Kemp, A. H., Gordon, E., Peduto, A., ... Williams, L. M.
- 24 (2008). The neural networks of inhibitory control in posttraumatic stress disorder. *Journal of*
- 25 *Psychiatry & Neuroscience : JPN*, 33(5), 413–422.

- 1 Feder, A., Fred-Torres, S., Southwick, S. M., & Charney, D. S. (2019). The Biology of Human  
2 Resilience: Opportunities for Enhancing Resilience Across the Life Span. *Biological Psychiatry*,  
3 86(6), 443–453. <https://doi.org/10.1016/j.biopsych.2019.07.012>
- 4 Feder, A., Mota, N., Salim, R., Rodriguez, J., Singh, R., Schaffer, J., ... Pietrzak, R. H. (2016). Risk,  
5 coping and PTSD symptom trajectories in World Trade Center responders. *Journal of*  
6 *Psychiatric Research*, 82, 68–79. <https://doi.org/10.1016/j.jpsychires.2016.07.003>
- 7 Felmingham, K. L., Falconer, E. M., Williams, L., Kemp, A. H., Allen, A., Peduto, A., & Bryant, R. A.  
8 (2014). Reduced Amygdala and Ventral Striatal Activity to Happy Faces in PTSD Is Associated  
9 with Emotional Numbing. *PLOS ONE*, 9(9), e103653.  
10 <https://doi.org/10.1371/journal.pone.0103653>
- 11 Finn, E. S., Glerean, E., Khojandi, A. Y., Nielson, D., Molfese, P. J., Handwerker, D. A., & Bandettini, P.  
12 A. (2020). Idiosynchrony: From shared responses to individual differences during naturalistic  
13 neuroimaging. *NeuroImage*, 215, 116828. <https://doi.org/10.1016/j.neuroimage.2020.116828>
- 14 Fitzgerald, J. M., Webb, E. K., Weis, C. N., Huggins, A. A., Bennett, K. P., Miskovich, T. A., ... Larson,  
15 C. L. (2022). Hippocampal Resting-State Functional Connectivity Forecasts Individual  
16 Posttraumatic Stress Disorder Symptoms: A Data-Driven Approach. *Biological Psychiatry:*  
17 *Cognitive Neuroscience and Neuroimaging*, 7(2), 139–149.  
18 <https://doi.org/10.1016/j.bpsc.2021.08.007>
- 19 Fonzo, G. A., Goodkind, M. S., Oathes, D. J., Zaiko, Y. V., Harvey, M., Peng, K. K., ... Etkin, A. (2017).  
20 PTSD Psychotherapy Outcome Predicted by Brain Activation During Emotional Reactivity and  
21 Regulation. *American Journal of Psychiatry*, 174(12), 1163–1174.  
22 <https://doi.org/10.1176/appi.ajp.2017.16091072>
- 23 Fonzo, G. A., Simmons, A. N., Thorp, S. R., Norman, S. B., Paulus, M. P., & Stein, M. B. (2010).  
24 Exaggerated and Disconnected Insular-Amygdalar Blood Oxygenation Level-Dependent  
25 Response to Threat-Related Emotional Faces in Women with Intimate-Partner Violence

- 1 Posttraumatic Stress Disorder. *Biological Psychiatry*, 68(5), 433–441.
- 2 <https://doi.org/10.1016/j.biopsych.2010.04.028>
- 3 Fritz, J., de Graaff, A. M., Caisley, H., van Harmelen, A.-L., & Wilkinson, P. O. (2018). A Systematic
- 4 Review of Amenable Resilience Factors That Moderate and/or Mediate the Relationship
- 5 Between Childhood Adversity and Mental Health in Young People. *Frontiers in Psychiatry*, 9.
- 6 <https://doi.org/10.3389/fpsy.2018.00230>
- 7 Fritz, J., Fried, E. I., Goodyer, I. M., Wilkinson, P. O., & van Harmelen, A.-L. (2018). A Network Model
- 8 of Resilience Factors for Adolescents with and without Exposure to Childhood Adversity.
- 9 *Scientific Reports*, 8(1), 15774. <https://doi.org/10.1038/s41598-018-34130-2>
- 10 Fritz, J., Stretton, J., Askelund, A. D., Schweizer, S., Walsh, N. D., Elzinga, B. M., ... van Harmelen, A.-
- 11 L. (2019). Mood and neural responses to social rejection do not seem to be altered in resilient
- 12 adolescents with a history of adversity. *Development and Psychopathology*, 1–13.
- 13 <https://doi.org/10.1017/S0954579419000178>
- 14 Fullana, M. A., Albajes-Eizaguirre, A., Soriano-Mas, C., Vervliet, B., Cardoner, N., Benet, O., ...
- 15 Harrison, B. J. (2018). Fear extinction in the human brain: A meta-analysis of fMRI studies in
- 16 healthy participants. *Neuroscience & Biobehavioral Reviews*, 88, 16–25.
- 17 <https://doi.org/10.1016/j.neubiorev.2018.03.002>
- 18 Fullana, M. A., Dunsmoor, J. E., Schruers, K. R. J., Savage, H. S., Bach, D. R., & Harrison, B. J.
- 19 (2020). Human fear conditioning: From neuroscience to the clinic. *Behaviour Research and*
- 20 *Therapy*, 124, 103528. <https://doi.org/10.1016/j.brat.2019.103528>
- 21 Gee, D. G. (2020). Caregiving influences on emotional learning and regulation: Applying a sensitive
- 22 period model. *Current Opinion in Behavioral Sciences*, 36, 177–184.
- 23 <https://doi.org/10.1016/j.cobeha.2020.11.003>
- 24 Gerin, M. I., Viding, E., Pingault, J.-B., Puetz, V. B., Knodt, A. R., Radtke, S. R., ... McCrory, E. J.
- 25 (2019). Heightened amygdala reactivity and increased stress generation predict internalizing

symptoms in adults following childhood maltreatment. *Journal of Child Psychology and Psychiatry*, 60(7), 752–761. <https://doi.org/10.1111/jcpp.13041>

Getnet, B., Medhin, G., & Alem, A. (2019). Symptoms of post-traumatic stress disorder and depression among Eritrean refugees in Ethiopia: Identifying direct, mediating and moderating predictors from path analysis. *BMJ Open*, 9(1), e021142. <https://doi.org/10.1136/bmjopen-2017-021142>

Goodyer, I. M., Croudace, T., Dunn, V., Herbert, J., & Jones, P. B. (2010). Cohort Profile: Risk patterns and processes for psychopathology emerging during adolescence: the ROOTS project. *International Journal of Epidemiology*, 39(2), 361–369. <https://doi.org/10.1093/ije/dyp173>

Gould, F., Clarke, J., Heim, C., Harvey, P. D., Majer, M., & Nemeroff, C. B. (2012). The effects of child abuse and neglect on cognitive functioning in adulthood. *Journal of Psychiatric Research*, 46(4), 500–506. Scopus. <https://doi.org/10.1016/j.jpsychires.2012.01.005>

Green, B. L. (1990). Defining Trauma: Terminology and Generic Stressor Dimensions. *Journal of Applied Social Psychology*, 20(20), 1632–1642. <https://doi.org/10.1111/j.1559-1816.1990.tb01498.x>

Greenberg, M. T. (2006). Promoting Resilience in Children and Youth. *Annals of the New York Academy of Sciences*, 1094(1), 139–150. <https://doi.org/10.1196/annals.1376.013>

Hanson, J. L., Albert, D., Iselin, A.-M. R., Carré, J. M., Dodge, K. A., & Hariri, A. R. (2016). Cumulative stress in childhood is associated with blunted reward-related brain activity in adulthood. *Social Cognitive and Affective Neuroscience*, 11(3), 405–412. <https://doi.org/10.1093/scan/nsv124>

Hanson, J. L., Hariri, A. R., & Williamson, D. E. (2015). Blunted Ventral Striatum Development in Adolescence Reflects Emotional Neglect and Predicts Depressive Symptoms. *Biological Psychiatry*, 78(9), 598–605. <https://doi.org/10.1016/j.biopsych.2015.05.010>

Harpur, L. J., Polek, E., & van Harmelen, A.-L. (2015). The role of timing of maltreatment and child intelligence in pathways to low symptoms of depression and anxiety in adolescence. *Child Abuse & Neglect*, 47, 24–37. <https://doi.org/10.1016/j.chiabu.2015.05.019>

- 1 Hofmann, S. G., Asmundson, G. J. G., & Beck, A. T. (2013). The Science of Cognitive Therapy.  
2 *Behavior Therapy*, 44(2), 199–212. <https://doi.org/10.1016/j.beth.2009.01.007>
- 3 Holz, N. E., Tost, H., & Meyer-Lindenberg, A. (2020). Resilience and the brain: A key role for regulatory  
4 circuits linked to social stress and support. *Molecular Psychiatry*, 25(2), 379–396.  
5 <https://doi.org/10.1038/s41380-019-0551-9>
- 6 Hoorelbeke, K., Van den Bergh, N., Wichers, M., & Koster, E. H. W. (2019). Between vulnerability and  
7 resilience: A network analysis of fluctuations in cognitive risk and protective factors following  
8 remission from depression. *Behaviour Research and Therapy*, 116, 1–9.  
9 <https://doi.org/10.1016/j.brat.2019.01.007>
- 10 Horn, S. R., Pietrzak, R. H., Schechter, C., Bromet, E. J., Katz, C. L., Reissman, D. B., ... Feder, A.  
11 (2016). Latent typologies of posttraumatic stress disorder in World Trade Center responders.  
12 *Journal of Psychiatric Research*, 83, 151–159. <https://doi.org/10.1016/j.jpsychires.2016.08.018>
- 13 Hudson, A., Van Hamme, C., Maeyens, L., Brass, M., & Mueller, S. (2018). *Spontaneous mentalizing*  
14 *after early interpersonal trauma: Evidence for hypoactivation of the temporoparietal junction*  
15 [Preprint]. Neuroscience. <https://doi.org/10.1101/487363>
- 16 Iadipaolo, A. S., Marusak, H. A., Paulisin, S. M., Sala-Hamrick, K., Crespo, L. M., Elrahal, F., ...  
17 Rabinak, C. A. (2018). Distinct neural correlates of trait resilience within core neurocognitive  
18 networks in at-risk children and adolescents. *NeuroImage. Clinical*, 20, 24–34.  
19 <https://doi.org/10.1016/j.nicl.2018.06.026>
- 20 Iigaya, K., Hauser, T. U., Kurth-Nelson, Z., O'Doherty, J. P., Dayan, P., & Dolan, R. J. (2019).  
21 Hippocampal-midbrain circuit enhances the pleasure of anticipation in the prefrontal cortex.  
22 *BioRxiv*, 588699. <https://doi.org/10.1101/588699>
- 23 Ioannidis, K., Askelund, A. D., Kievit, R. A., & van Harmelen, A.-L. (2020). The complex neurobiology of  
24 resilient functioning after childhood maltreatment. *BMC Medicine*, 18(1), 32.  
25 <https://doi.org/10.1186/s12916-020-1490-7>



- 1 Jacob, S. N., Dodge, C. P., & Vasterling, J. J. (2019). Posttraumatic stress disorder and neurocognition:  
2 A bidirectional relationship? *Clinical Psychology Review*, 72, 101747.  
3 <https://doi.org/10.1016/j.cpr.2019.101747>
- 4 Jeong, H., Park, S., Dager, S. R., Lim, S. M., Lee, S. L., Hong, H., ... Lyoo, I. K. (2019). Altered  
5 functional connectivity in the fear network of firefighters with repeated traumatic stress. *The*  
6 *British Journal of Psychiatry*, 214(6), 347–353. <https://doi.org/10.1192/bjp.2018.260>
- 7 Kaldewaij, R., Koch, S. B. J., Hashemi, M. M., Zhang, W., Klumpers, F., & Roelofs, K. (2021). Anterior  
8 prefrontal brain activity during emotion control predicts resilience to post-traumatic stress  
9 symptoms. *Nature Human Behaviour*. <https://doi.org/10.1038/s41562-021-01055-2>
- 10 Kalisch, R., Baker, D. G., Basten, U., Boks, M. P., Bonanno, G. A., Brummelman, E., ... Kleim, B.  
11 (2017). The resilience framework as a strategy to combat stress-related disorders. *Nature*  
12 *Human Behaviour*, 1(11), 784. <https://doi.org/10.1038/s41562-017-0200-8>
- 13 Karam, E. G., Friedman, M. J., Hill, E. D., Kessler, R. C., McLaughlin, K. A., Petukhova, M., ... Koenen,  
14 K. C. (2014). Cumulative Traumas and Risk Thresholds: 12-Month PTSD in the World Mental  
15 Health (WMH) Surveys. *Depression and Anxiety*, 31(2), 130–142.  
16 <https://doi.org/10.1002/da.22169>
- 17 Keane, T. M., Fairbank, J. A., Caddell, J. M., Zimering, R. T., Taylor, K. L., & Mora, C. A. (1989).  
18 Clinical evaluation of a measure to assess combat exposure. *Psychological Assessment: A*  
19 *Journal of Consulting and Clinical Psychology*, 1(1), 53–55. [https://doi.org/10.1037/1040-](https://doi.org/10.1037/1040-3590.1.1.53)  
20 [3590.1.1.53](https://doi.org/10.1037/1040-3590.1.1.53)
- 21 Keynan, J. N., Cohen, A., Jackont, G., Green, N., Goldway, N., Davidov, A., ... Hendler, T. (2019).  
22 Electrical fingerprint of the amygdala guides neurofeedback training for stress resilience. *Nature*  
23 *Human Behaviour*, 3(1), 63. <https://doi.org/10.1038/s41562-018-0484-3>
- 24 Lepore, S. J., & Kliewer, W. (2019). Social Intelligence Attenuates Association between Peer  
25 Victimization and Depressive Symptoms among Adolescents. *Psychology of Violence*, 9(6),  
26 644–652. <https://doi.org/10.1037/vio0000234>

- 1 Liebenberg, L. (2020). Reconsidering interactive resilience processes in mental health: Implications for  
2 child and youth services. *Journal of Community Psychology*, 48(5), 1365–1380.  
3 <https://doi.org/10.1002/jcop.22331>
- 4 Lissek, S., & van Meurs, B. (2015). Learning models of PTSD: Theoretical accounts and  
5 psychobiological evidence. *International Journal of Psychophysiology*, 98(3, Part 2), 594–605.  
6 <https://doi.org/10.1016/j.ijpsycho.2014.11.006>
- 7 Luthar, S. S., Sawyer, J. A., & Brown, P. J. (2006). Conceptual Issues in Studies of Resilience. *Annals*  
8 *of the New York Academy of Sciences*, 1094(1), 105–115.  
9 <https://doi.org/10.1196/annals.1376.009>
- 10 Maier, S. F., & Watkins, L. R. (2010). Role of the medial prefrontal cortex in coping and resilience. *Brain*  
11 *Research*, 1355, 52–60. <https://doi.org/10.1016/j.brainres.2010.08.039>
- 12 Mary, A., Dayan, J., Leone, G., Postel, C., Fraisse, F., Malle, C., ... Gagnepain, P. (2020). Resilience  
13 after trauma: The role of memory suppression. *Science*, 367(6479).  
14 <https://doi.org/10.1126/science.aay8477>
- 15 McGloin, J. M., & Widom, C. S. (2001). Resilience among abused and neglected children grown up.  
16 *Development and Psychopathology*, 13(4), 1021–1038.
- 17 McLean, C. P., & Anderson, E. R. (2009). Brave men and timid women? A review of the gender  
18 differences in fear and anxiety. *Clinical Psychology Review*, 29(6), 496–505.  
19 <https://doi.org/10.1016/j.cpr.2009.05.003>
- 20 McLean, S. A., Ressler, K., Koenen, K. C., Neylan, T., Germine, L., Jovanovic, T., ... Kessler, R.  
21 (2020). The AURORA Study: A longitudinal, multimodal library of brain biology and function  
22 after traumatic stress exposure. *Molecular Psychiatry*, 25(2), 283–296.  
23 <https://doi.org/10.1038/s41380-019-0581-3>
- 24 Meili, I., & Maercker, A. (2019). Cultural perspectives on positive responses to extreme adversity: A  
25 playing field for metaphors. *Transcultural Psychiatry*, 56(5), 1056–1075.  
26 <https://doi.org/10.1177/1363461519844355>

- 1 Mendenhall, E., & Kim, A. W. (2019). How to Fail a Scale: Reflections on a Failed Attempt to Assess  
2 Resilience. *Culture, Medicine, and Psychiatry*, 43(2), 315–325. [https://doi.org/10.1007/s11013-](https://doi.org/10.1007/s11013-018-9617-4)  
3 018-9617-4
- 4 Méndez Leal, A. S., & Silvers, J. A. (2021). Neurobiological Markers of Resilience to Early-Life  
5 Adversity During Adolescence. *Biological Psychiatry: Cognitive Neuroscience and*  
6 *Neuroimaging*, 6(2), 238–247. <https://doi.org/10.1016/j.bpsc.2020.08.004>
- 7 Milad, M. R., Wright, C. I., Orr, S. P., Pitman, R. K., Quirk, G. J., & Rauch, S. L. (2007). Recall of Fear  
8 Extinction in Humans Activates the Ventromedial Prefrontal Cortex and Hippocampus in  
9 Concert. *Biological Psychiatry*, 62(5), 446–454. <https://doi.org/10.1016/j.biopsych.2006.10.011>
- 10 Moreno-López, L., Ioannidis, K., Askelund, A. D., Smith, A. J., Schueler, K., & van Harmelen, A.-L.  
11 (2020). The Resilient Emotional Brain: A Scoping Review of the Medial Prefrontal Cortex and  
12 Limbic Structure and Function in Resilient Adults With a History of Childhood Maltreatment.  
13 *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 5(4), 392–402.  
14 <https://doi.org/10.1016/j.bpsc.2019.12.008>
- 15 Mowinckel, A. M., & Vidal-Piñeiro, D. (2019, December 17). *Visualisation of Brain Statistics with R-*  
16 *packages ggseg and ggseg3d*. arXiv. <https://doi.org/10.48550/arXiv.1912.08200>
- 17 Mowinckel, A. M., & Vidal-Piñeiro, D. (n.d.). *GgsegExtra* [R]. Retrieved from  
18 <https://github.com/ggseg/ggsegExtra> (Original work published 2019)
- 19 Nawijn, L., van Zuiden, M., Frijling, J. L., Koch, S. B. J., Veltman, D. J., & Olff, M. (2015). Reward  
20 functioning in PTSD: A systematic review exploring the mechanisms underlying anhedonia.  
21 *Neuroscience & Biobehavioral Reviews*, 51, 189–204.  
22 <https://doi.org/10.1016/j.neubiorev.2015.01.019>
- 23 Nord, C. L., Gray, A., Charpentier, C. J., Robinson, O. J., & Roiser, J. P. (2017). Unreliability of putative  
24 fMRI biomarkers during emotional face processing. *NeuroImage*, 156, 119–127.  
25 <https://doi.org/10.1016/j.neuroimage.2017.05.024>

- 1 Patel, R., Spreng, R. N., Shin, L. M., & Girard, T. A. (2012). Neurocircuitry models of posttraumatic  
2 stress disorder and beyond: A meta-analysis of functional neuroimaging studies. *Neuroscience*  
3 & *Biobehavioral Reviews*, 36(9), 2130–2142. <https://doi.org/10.1016/j.neubiorev.2012.06.003>
- 4 Phelps, E. A., Delgado, M. R., Nearing, K. I., & LeDoux, J. E. (2004). Extinction Learning in Humans:  
5 Role of the Amygdala and vmPFC. *Neuron*, 43(6), 897–905.  
6 <https://doi.org/10.1016/j.neuron.2004.08.042>
- 7 Pietrzak, R. H., el-Gabalawy, R., Tsai, J., Sareen, J., Neumeister, A., & Southwick, S. M. (2014).  
8 Typologies of posttraumatic stress disorder in the U.S. adult population. *Journal of Affective*  
9 *Disorders*, 162, 102–106. <https://doi.org/10.1016/j.jad.2014.03.024>
- 10 Pietrzak, R. H., Feder, A., Singh, R., Schechter, C. B., Bromet, E. J., Katz, C. L., ... Southwick, S. M.  
11 (2014). Trajectories of PTSD risk and resilience in World Trade Center responders: An 8-year  
12 prospective cohort study. *Psychological Medicine*, 44(1), 205–219.  
13 <https://doi.org/10.1017/S0033291713000597>
- 14 Pitman, R. K., Rasmusson, A. M., Koenen, K. C., Shin, L. M., Orr, S. P., Gilbertson, M. W., ... Liberzon,  
15 I. (2012). Biological studies of post-traumatic stress disorder. *Nature Reviews Neuroscience*,  
16 13(11), 769–787. <https://doi.org/10.1038/nrn3339>
- 17 Portnoy, G. A., Relyea, M. R., Decker, S., Shamaskin-Garroway, A., Driscoll, M., Brandt, C. A., &  
18 Haskell, S. G. (2018). Understanding Gender Differences in Resilience Among Veterans:  
19 Trauma History and Social Ecology. *Journal of Traumatic Stress*, 31(6), 845–855.  
20 <https://doi.org/10.1002/jts.22341>
- 21 Powers, M. B., Warren, A. M., Rosenfield, D., Roden-Foreman, K., Bennett, M., Reynolds, M. C., ...  
22 Smits, J. A. J. (2014). Predictors of PTSD symptoms in adults admitted to a Level I trauma  
23 center: A prospective analysis. *Journal of Anxiety Disorders*, 28(3), 301–309.  
24 <https://doi.org/10.1016/j.janxdis.2014.01.003>
- 25 Reddan, M. C., Wager, T. D., & Schiller, D. (2018). Attenuating Neural Threat Expression with  
26 Imagination. *Neuron*, 100(4), 994-1005.e4. <https://doi.org/10.1016/j.neuron.2018.10.047>

- 1 Rodman, A. M., Jenness, J. L., Weissman, D. G., Pine, D. S., & McLaughlin, K. A. (2019).  
2 Neurobiological Markers of Resilience to Depression Following Childhood Maltreatment: The  
3 Role of Neural Circuits Supporting the Cognitive Control of Emotion. *Biological Psychiatry*,  
4 86(6), 464–473. <https://doi.org/10.1016/j.biopsych.2019.04.033>
- 5 Sailer, U., Robinson, S., Fischmeister, F. P. S., König, D., Oppenauer, C., Lueger-Schuster, B., ...  
6 Bauer, H. (2008). Altered reward processing in the nucleus accumbens and mesial prefrontal  
7 cortex of patients with posttraumatic stress disorder. *Neuropsychologia*, 46(11), 2836–2844.  
8 <https://doi.org/10.1016/j.neuropsychologia.2008.05.022>
- 9 Schiller, D., & Delgado, M. R. (2010). Overlapping neural systems mediating extinction, reversal and  
10 regulation of fear. *Trends in Cognitive Sciences*, 14(6), 268–276.  
11 <https://doi.org/10.1016/j.tics.2010.04.002>
- 12 Schultze-Lutter, F., Schimmelmann, B. G., & Schmidt, S. J. (2016). Resilience, risk, mental health and  
13 well-being: Associations and conceptual differences. *European Child & Adolescent Psychiatry*,  
14 25(5), 459–466. <https://doi.org/10.1007/s00787-016-0851-4>
- 15 Schweizer, S., Walsh, N. D., Stretton, J., Dunn, V. J., Goodyer, I. M., & Dalgleish, T. (2016). Enhanced  
16 emotion regulation capacity and its neural substrates in those exposed to moderate childhood  
17 adversity. *Social Cognitive and Affective Neuroscience*, 11(2), 272–281.  
18 <https://doi.org/10.1093/scan/nsv109>
- 19 Scott, J. C., Matt, G. E., Wrocklage, K. M., Crnich, C., Jordan, J., Southwick, S. M., ... Schweinsburg,  
20 B. C. (2015). A quantitative meta-analysis of neurocognitive functioning in posttraumatic stress  
21 disorder. *Psychological Bulletin*, 141(1), 105–140. <https://doi.org/10.1037/a0038039>
- 22 Scult, M. A., Knodt, A. R., Radtke, S. R., Brigidi, B. D., & Hariri, A. R. (2019). Prefrontal Executive  
23 Control Rescues Risk for Anxiety Associated with High Threat and Low Reward Brain Function.  
24 *Cerebral Cortex*, 29(1), 70–76. <https://doi.org/10.1093/cercor/bhx304>
- 25 Seeley, S. H., Boukezzzi, S., DePierro, J. M., Charney, D. S., & Feder, A. (In press). Biological  
26 mechanisms of stress resilience: Human and animal studies. In *Charney and Nestler's*

*Neurobiology of Mental Illness* (eds. Charney, D.S, Nestler, E.J., Buxbaum, J.D., Binder, E., Gordon, J.A., and Picciotto, M.) (6th ed.). Oxford University Press.

Shalev, A., Liberzon, I., & Marmar, C. (2017). Post-Traumatic Stress Disorder. *New England Journal of Medicine*, 376(25), 2459–2469. <https://doi.org/10.1056/NEJMra1612499>

Silveira, S., Shah, R., Nooner, K. B., Nagel, B. J., Tapert, S. F., de Bellis, M. D., & Mishra, J. (2020). Impact of Childhood Trauma on Executive Function in Adolescence—Mediating Functional Brain Networks and Prediction of High-Risk Drinking. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 5(5), 499–509. <https://doi.org/10.1016/j.bpsc.2020.01.011>

Silvers, J. A., Lumian, D. S., Gabard-Durnam, L., Gee, D. G., Goff, B., Fareri, D. S., ... Tottenham, N. (2016). Previous Institutionalization Is Followed by Broader Amygdala-Hippocampal-PFC Network Connectivity during Aversive Learning in Human Development. *Journal of Neuroscience*, 36(24), 6420–6430. <https://doi.org/10.1523/JNEUROSCI.0038-16.2016>

Sinha, R., Lacadie, C. M., Constable, R. T., & Seo, D. (2016). Dynamic neural activity during stress signals resilient coping. *Proceedings of the National Academy of Sciences*, 113(31), 8837–8842. <https://doi.org/10.1073/pnas.1600965113>

Southwick, S. M., Charney, D. S., & DePierro, J. M. (2023). *Resilience: The Science of Mastering Life's Greatest Challenges* (3rd ed.). Cambridge University Press.

Speer, M. E., & Delgado, M. R. (2017). Reminiscing about positive memories buffers acute stress responses. *Nature Human Behaviour*, 1(5), 0093. <https://doi.org/10.1038/s41562-017-0093>

Spreng, R. N., & Andrews-Hanna, J. (2015). The Default Network and Social Cognition. *Brain Mapping: An Encyclopedic Reference*, 3, 165–169. <https://doi.org/10.1016/B978-0-12-397025-1.00173-1>

Stevens, J. S., Harnett, N. G., Lebois, L. A. M., van Rooij, S. J. H., Ely, T. D., Roeckner, A., ... Ressler, K. J. (2021). Brain-Based Biotypes of Psychiatric Vulnerability in the Acute Aftermath of Trauma. *American Journal of Psychiatry*, 178(11), 1037–1049. <https://doi.org/10.1176/appi.ajp.2021.20101526>

- 1 Stevens, J. S., & Jovanovic, T. (2019). Role of social cognition in post-traumatic stress disorder: A  
2 review and meta-analysis. *Genes, Brain and Behavior*, 18(1), e12518.  
3 <https://doi.org/10.1111/gbb.12518>
- 4 Stevens, J. S., Kim, Y. J., Galatzer-Levy, I. R., Reddy, R., Ely, T. D., Nemeroff, C. B., ... Ressler, K. J.  
5 (2017). Amygdala Reactivity and Anterior Cingulate Habituation Predict Posttraumatic Stress  
6 Disorder Symptom Maintenance After Acute Civilian Trauma. *Biological Psychiatry*, 81(12),  
7 1023–1029. <https://doi.org/10.1016/j.biopsych.2016.11.015>
- 8 Stratta, P., Capanna, C., Dell'Osso, L., Carmassi, C., Patriarca, S., Di Emidio, G., ... Rossi, A. (2015).  
9 Resilience and coping in trauma spectrum symptoms prediction: A structural equation modeling  
10 approach. *Personality and Individual Differences*, 77, 55–61.  
11 <https://doi.org/10.1016/j.paid.2014.12.035>
- 12 Street, A. E., & Dardis, C. M. (2018). Using a social construction of gender lens to understand gender  
13 differences in posttraumatic stress disorder. *Clinical Psychology Review*, 66, 97–105.  
14 <https://doi.org/10.1016/j.cpr.2018.03.001>
- 15 Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A neural biomarker of psychological  
16 vulnerability to future life stress. *Neuron*, 85(3), 505–511.  
17 <https://doi.org/10.1016/j.neuron.2014.12.055>
- 18 Szucs, D., & Ioannidis, J. PA. (2020). Sample size evolution in neuroimaging research: An evaluation of  
19 highly-cited studies (1990–2012) and of latest practices (2017–2018) in high-impact journals.  
20 *NeuroImage*, 221, 117164. <https://doi.org/10.1016/j.neuroimage.2020.117164>
- 21 Thompson, N. J., Fiorillo, D., Rothbaum, B. O., Ressler, K. J., & Michopoulos, V. (2018). Coping  
22 strategies as mediators in relation to resilience and posttraumatic stress disorder. *Journal of*  
23 *Affective Disorders*, 225, 153–159. <https://doi.org/10.1016/j.jad.2017.08.049>
- 24 Thompson, P. M., Jahanshad, N., Ching, C. R. K., Salminen, L. E., Thomopoulos, S. I., Bright, J., ...  
25 Zelman, V. (2020). ENIGMA and global neuroscience: A decade of large-scale studies of the

- 1 brain in health and disease across more than 40 countries. *Translational Psychiatry*, 10(1), 1–  
2 28. <https://doi.org/10.1038/s41398-020-0705-1>
- 3 Tolin, D. F., & Foa, E. B. (2008). Sex differences in trauma and posttraumatic stress disorder: A  
4 quantitative review of 25 years of research. *Psychological Trauma: Theory, Research, Practice,*  
5 *and Policy*, S(1), 37–85. <https://doi.org/10.1037/1942-9681.S.1.37>
- 6 Troy, A. S., & Mauss, I. B. (2011). Resilience in the face of stress: Emotion regulation as a protective  
7 factor. In B. T. Litz, D. Charney, M. J. Friedman, & S. M. Southwick (Eds.), *Resilience and*  
8 *Mental Health: Challenges Across the Lifespan* (pp. 30–44). Cambridge: Cambridge University  
9 Press. <https://doi.org/10.1017/CBO9780511994791.004>
- 10 van der Werff, S. J. A., van den Berg, S. M., Pannekoek, J. N., Elzinga, B. M., & Van Der Wee, N. J. A.  
11 (2013). Neuroimaging resilience to stress: A review. *Frontiers in Behavioral Neuroscience*, 7.  
12 <https://doi.org/10.3389/fnbeh.2013.00039>
- 13 van Harmelen, A. L., Kievit, R. A., Ioannidis, K., Neufeld, S., Jones, P. B., Bullmore, E., ... Goodyer, I.  
14 (2017). Adolescent friendships predict later resilient functioning across psychosocial domains in  
15 a healthy community cohort. *Psychological Medicine*, 47(13), 2312–2322.  
16 <https://doi.org/10.1017/S0033291717000836>
- 17 van Harmelen, A.-L., Gibson, J. L., Clair, M. C. S., Owens, M., Brodbeck, J., Dunn, V., ... Goodyer, I.  
18 M. (2016). Friendships and Family Support Reduce Subsequent Depressive Symptoms in At-  
19 Risk Adolescents. *PLOS ONE*, 11(5), e0153715. <https://doi.org/10.1371/journal.pone.0153715>
- 20 van Rooij, S. J. H., Stevens, J. S., Ely, T. D., Hinrichs, R., Michopoulos, V., Winters, S. J., ...  
21 Jovanovic, T. (2018). The Role of the Hippocampus in Predicting Future Posttraumatic Stress  
22 Disorder Symptoms in Recently Traumatized Civilians. *Biological Psychiatry*, 84(2), 106–115.  
23 <https://doi.org/10.1016/j.biopsych.2017.09.005>
- 24 Vythilingam, M., Nelson, E. E., Scaramozza, M., Waldeck, T., Hazlett, G., Southwick, S. M., ... Ernst,  
25 M. (2009). Reward circuitry in resilience to severe trauma: An fMRI investigation of resilient



special forces soldiers. *Psychiatry Research: Neuroimaging*, 172(1), 75–77.

<https://doi.org/10.1016/j.psychres.2008.06.008>

Walsh, W. A., Dawson, J., & Mattingly, M. J. (2010). How Are We Measuring Resilience Following Childhood Maltreatment? Is the Research Adequate and Consistent? What is the Impact on Research, Practice, and Policy? *Trauma, Violence, & Abuse*, 11(1), 27–41.

<https://doi.org/10.1177/1524838009358892>

Wang, L., Paul, N., Stanton, S. J., Greeson, J. M., & Smoski, M. (2013). Loss of Sustained Activity in the Ventromedial Prefrontal Cortex in Response to Repeated Stress in Individuals with Early-Life Emotional Abuse: Implications for Depression Vulnerability. *Frontiers in Psychology*, 4.

<https://doi.org/10.3389/fpsyg.2013.00320>

Watson, P. (2019). PTSD as a Public Mental Health Priority. *Current Psychiatry Reports*, 21(7), 61.

<https://doi.org/10.1007/s11920-019-1032-1>

Waugh, C. E., & Koster, E. H. W. (2015). A resilience framework for promoting stable remission from depression. *Clinical Psychology Review*, 41, 49–60. <https://doi.org/10.1016/j.cpr.2014.05.004>

Weathers, F. W., Blake, D. D., Schnurr, P. P., Kaloupek, D. G., Marx, B. P., & Keane, T. M. (2013).

Clinician-Administered PTSD Scale for DSM-5 (CAPS-5). Retrieved October 9, 2018, from

<https://www.ptsd.va.gov/professional/assessment/adult-int/caps.asp>

Whittle, S., Dennison, M., Vijayakumar, N., Simmons, J. G., Yücel, M., Lubman, D. I., ... Allen, N. B.

(2013). Childhood Maltreatment and Psychopathology Affect Brain Development During

Adolescence. *Journal of the American Academy of Child & Adolescent Psychiatry*, 52(9), 940-

952.e1. <https://doi.org/10.1016/j.jaac.2013.06.007>

Williams, M., Metzger, I., Leins, C., & DeLapp, C. (2018). Assessing Racial Trauma Within a DSM–5

Framework: The UConn Racial/Ethnic Stress & Trauma Survey. *Practice Innovations*, 3(4),

242–260. <https://doi.org/10.1037/pri0000076>

World Health Organization. (1993). *The ICD-10 Classification of Mental and Behavioural Disorders:*

*Diagnostic Criteria for Research*. World Health Organization.

- 1 Wymbs, N. F., Orr, C., Albaugh, M. D., Althoff, R. R., O'Loughlin, K., Holbrook, H., ... Kaufman, J.  
2 (2020). Social supports moderate the effects of child adversity on neural correlates of threat  
3 processing. *Child Abuse & Neglect*, 102, 104413. <https://doi.org/10.1016/j.chiabu.2020.104413>
- 4 Yamamoto, T., Toki, S., Siegle, G. J., Takamura, M., Takaishi, Y., Yoshimura, S., ... Yamawaki, S.  
5 (2017). Increased amygdala reactivity following early life stress: A potential resilience enhancer  
6 role. *BMC Psychiatry*, 17(1), 27. <https://doi.org/10.1186/s12888-017-1201-x>
- 7 Yehuda, R., & Flory, J. D. (2007). Differentiating biological correlates of risk, PTSD, and resilience  
8 following trauma exposure. *Journal of Traumatic Stress*, 20(4), 435–447.  
9 <https://doi.org/10.1002/jts.20260>
- 10 Yehuda, R., Hoge, C. W., McFarlane, A. C., Vermetten, E., Lanius, R. A., Nievergelt, C. M., ... Hyman,  
11 S. E. (2015). Post-traumatic stress disorder. *Nature Reviews Disease Primers*, 1, 15057.  
12 <https://doi.org/10.1038/nrdp.2015.57>
- 13 Yule, K., Houston, J., & Grych, J. (2019). Resilience in Children Exposed to Violence: A Meta-analysis  
14 of Protective Factors Across Ecological Contexts. *Clinical Child and Family Psychology Review*,  
15 22(3), 406–431. <https://doi.org/10.1007/s10567-019-00293-1>
- 16 Zhang, W., Kaldewaij, R., Hashemi, M. M., Koch, S. B. J., Smit, A., van Ast, V. A., ... Roelofs, K.  
17 (2022). Acute-stress-induced change in salience network coupling prospectively predicts post-  
18 trauma symptom development. *Translational Psychiatry*, 12(1), 63.  
19 <https://doi.org/10.1038/s41398-022-01798-0>
- 20 Zilverstand, A., Parvaz, M. A., & Goldstein, R. Z. (2017). Neuroimaging cognitive reappraisal in clinical  
21 populations to define neural targets for enhancing emotion regulation. A systematic review.  
22 *NeuroImage*, 151, 105–116. <https://doi.org/10.1016/j.neuroimage.2016.06.009>
- 23 Zotev, V., Phillips, R., Misaki, M., Wong, C. K., Wurfel, B. E., Krueger, F., ... Bodurka, J. (2018). Real-  
24 time fMRI neurofeedback training of the amygdala activity with simultaneous EEG in veterans  
25 with combat-related PTSD. *NeuroImage: Clinical*, 19, 106–121.  
26 <https://doi.org/10.1016/j.nicl.2018.04.010>

