Table 1: Ordinary regression with multiple displayed covariates

	Price	Trunk space (cu. ft.)	Turn circle (ft.)
Foreign origin	3,573.1***	0.677	-1.74***
	(647.6)	(0.965)	(0.612)
	[0.001]	[0.194]	[0.006]
Weight (lb.)	5.77***	-0.001	0.002**
	(1.50)	(0.001)	(0.001)
	[0.001]	[0.220]	[0.049]
Length (inches)	-91.37*	0.167***	0.097***
	(48.18)	(0.034)	(0.026)
	[0.022]	[0.001]	[0.001]
Control mean	6,072.4	14.75	41.44
N	74	74	74

Notes: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$, using conventional p-values. Standard errors appear in round brackets. Square brackets contain sharpened q-values. Stata's 1978 automobile data. Each outcome is regressed on foreign origin, weight and length; all slopes are displayed. These covariates are not treatment arms, and no equality tests are requested. Standard errors are heteroskedasticity-robust with finite-df t inference. No weights or fixed effects. Control mean denotes the domestic-car outcome mean within each model's estimation sample. N counts cars. Observational associations, not causal estimates. Anderson/BKY sharpened q-values use the .001 grid across all three outcomes separately for each covariate. RI is inapplicable.