Table 1: Synthetic treatment-effect demonstration

	Synthetic endline outcome 1	Synthetic endline outcome 2	Synthetic endline outcome 3
Panel 1: Specification 1			
Treatment arm 1	-0.068	2.33	85.23***
	(0.234)	(2.27)	(11.76)
	[0.856]	[0.444]	[0.001]
Treatment arm 2	0.308	4.01*	122.1***
	(0.249)	(2.20)	(11.94)
	[0.119]	[0.077]	[0.001]
p-value: Treatment 1 = Treatment 2	0.081	0.376	0.004
Control mean	0.089	38.53	194.7
N	120	107	120
Panel 3: Specification 3			
Treatment arm 1	-0.002	2.24	74.26***
	(0.227)	(2.04)	(12.02)
	[0.701]	[0.379]	[0.001]
Treatment arm 2	0.387	5.35***	117.5***
	(0.263)	(1.83)	(11.50)
	[0.051]	[0.005]	[0.001]
p-value: Treatment 1 = Treatment 2	0.072	0.057	0.000
Control mean	0.055	37.92	196.8
N	110	98	110

Notes: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$, using conventional p-values. Standard errors appear in round brackets. Square brackets contain sharpened q-values. Synthetic examples, not empirical findings. Outcome definitions and data generation are in examples/stata/data.do. OLS with heteroskedasticity-robust standard errors; specification 1 has no additional controls; specification 3 controls for xmiss. No fixed effects or weights. Sharpened q-values use Anderson (2008) and BKY (2006), conventional p-values, a .001 grid and a complete outcome family for each panel and treatment effect. No RI was run. Control means use arm=0 at endline in each model's estimation sample; missing model variables are excluded model by model. Equality-test rows report unadjusted robust Wald F p-values for treatment 1 versus each other treatment, using the model covariance and residual degrees of freedom.