WOMEN

A
Mean glucose
q=0.126
Mean glucose (mg/dL)
99
98
97
96
95
Follicular
Luteal

B
CV
q=0.022 | perm p=0.009
CV (%)
14.5
14
13.5
13
12.5
12
11.5
Follicular
Luteal

C
MAGE
q=0.022 | perm p=0.015
MAGE (mg/dL)
36
34
32
30
Follicular
Luteal

D
GMI
q=0.126
GMI (%)
5.67
5.65
5.62
5.6
Follicular
Luteal

MEN

E
Mean glucose
q=0.129
Mean glucose (mg/dL)
103
102
101
100
99
98
Pseudo-follicular
Pseudo-luteal
0
25
50
75
100
Cycle progress (%)

F
CV
q=0.741
CV (%)
14
13.5
13
12.5
Pseudo-follicular
Pseudo-luteal
0
25
50
75
100
Cycle progress (%)

G
MAGE
q=0.541
MAGE (mg/dL)
38
36
34
32
Pseudo-follicular
Pseudo-luteal
0
25
50
75
100
Cycle progress (%)

H
GMI
q=0.129
GMI (%)
5.78
5.75
5.72
5.7
5.67
5.65
Pseudo-follicular
Pseudo-luteal
0
25
50
75
100
Cycle progress (%)

I
Cycle-effect significance
Mean
CV
MAGE
Range (Max-Min)
SD
MAG
SDWSH
GMI
Women
Men
q=0.05
0.0
0.5
1.0
1.5
Cycle-effect significance $-\log_{10}(q)$

J
Cyclic amplitude for women
CV
8.7% (1.15 %)
MAGE
9.5% (3.11 mg/dL)
Range (Max-Min)
8.4% (5.22 mg/dL)
SD
8.2% (1.05 mg/dL)
MAG
8.1% (0.76 mg/dL/h)
SDWSH
6.0% (0.29 mg/dL)
0
5
10
15
20
Cyclic amplitude (% of cohort cycle-average, 95% CI)

Cycle average
Moving average
Binned mean ±SEM
Cyclic spline (n.s.)
Cyclic spline (sig.)
Peak window
Nadir window