






A
Hourly CV
Follicular
Luteal
night time
Within-hour CV (%)
Hour of day



B
Daytime and overnight phase contrasts
Follicular
Luteal
*
ns
Within-hour CV (%)
Day time
(05-21 h)
Night time
(00-05 h)


C
Mean postprandial response
Follicular (n=1,881)
Luteal (n=1,934)
Δ Luteal − Follicular
glucose above baseline (mg/dL)
Δ Luteal − Follicular (mg/dL)
minutes from meal


D
Population GLMM: PPGR step by phase
Descent Rate
+11.5% *
iAUC
+5.3% ns
Peak Rise
+4.1% ns
Time-to-peak
+2.9% ns
Baseline
-0.2% ns
Luteal − Follicular step (% of follicular mean, 95% CI)


E
Phase contrasts by meal window
Descent Rate
+22.5%* (n=921)
+3.7% (n=746)
+12.6% (n=940)
iAUC
+20.4%* (n=1008)
-8.3% (n=833)
+0.6% (n=1047)
Peak Rise
+15.2%* (n=1008)
-6.2% (n=833)
+1.4% (n=1047)
Time-to-peak
+0.5% (n=1008)
+0.5% (n=833)
+3.7% (n=1047)
Baseline
+0.6% (n=1008)
+1.0% (n=833)
-0.9% (n=1047)
Breakfast
Lunch
Dinner
Luteal − follicular (%)


F
Descent Rate at breakfast
n=921 | q=0.039 | perm p=0.005
Descent Rate (mg/dL/min)
Follicular
Luteal
Cycle progress (%)


G
Peak Rise at breakfast
n=1,008 | q=0.045 | perm p=0.014
Peak Rise (mg/dL)
Follicular
Luteal
Cycle progress (%)
Cycle average
Moving average
Binned mean ±SEM
All meals (n.s.)
Breakfast spline (sig.)
Peak window
Nadir window