

Medical Applications and Biological Effects of Radiation

Sarita Kanwar

21/01/2022

Table of Contents

1. Abstract
2. Introduction
3. Sources of Radiation
4. Biological Effects of Radiation
5. Dosimetry

Abstract

With the development of radio-therapeutic oncology, computer technology and medical imaging technology, radiation therapy has made great progress[12]. Radiation's benefits were initially realised in the use of X-rays for medical diagnosis, and then later with the discovery of radiation and radioactivity. The desire to take advantage of the medical benefits led to a reasonable assessment of the hazards and resulting consequences. Only the most evident effects of large doses of radiation, such as radiation burns, were noticed in those early days, and protection efforts were concentrated on preventing them, mostly for practitioners rather than patients[3]. In this term paper we will be discussing about the various sources of radiation, medical applications of radiation and how the damage produced by ionizing radiation in biological molecules affects the body cells.

Introduction

Modern techniques in nuclear medicine have proven to be powerful tools for diagnosis and treatment of an increasing number of diseases[9]. However, they all add to the patient's radiation exposure. As a result, constant attention should be paid to reducing the radiation dose involved. The risks associated with a specific nuclear procedure should be known in order to weigh its advantage

against its disadvantages[1]. These risks depend on the dose and dose and dose-effect relationship. The discipline involved in dose calculation is called dosimetry. The combination of the results of dosimetry and dose-effect relation allows the estimation of the risk associated with a certain nuclear procedure[11].

People would appreciate that radiation has quiet and useful applications in our daily lives once they understand it. New difficulties in terms of worldwide levels of radiation exposure continue to emerge, new biological information on the impacts of radiation exposure is becoming available[13].

Sources of Radiation

As a part of living on earth, people are exposed to radiation from various sources every day. Radiation is the energy that travels through space, in the form of particles or electromagnetic waves such as radio, microwaves, infra-red, visible light, ultraviolet, alpha particles, X-rays and Gamma-rays etc[13]. According to [8] these sources of ionizing radiation could be from natural background radiation such as radon and thoron, cosmic and terrestrial radiation, or man-made radiation such as those from xray or nuclear medicine (NM) procedures.

Natural Radiation

The assessment of the natural radiation doses from natural sources in human is of particular importance natural radiation is the largest contributor to the collective dose of world population[2]. The natural radiation sources are classified into:

- External Irradiation
- Internal Irradiation

External Irradiation

1. Cosmic Radiation:

This is simply the radiation from the sun and stars. Flying based at high altitudes much frequently and for long duration will attract extra cosmic radiation exposure[5].

2. Terrestrial Radiation:

This is the radiation due to the presence of radioactive materials such as uranium, thorium, and radium that exist naturally in soil, water and rocks. Essentially air contains radon, which is responsible for the dose from natural background sources, and all organic matter (plant and animal) also contains radioactive carbon and potassium[7]. However, the dose from these sources varies in different parts of the world, but locations with higher soil concentrations of uranium and thorium generally have higher doses. Therefore, the background radiation levels vary in certain

areas due to geological differences and sometimes the exposure can be more than 200 times higher than the global average[10].

Internal Irradiation

From birth to death, this sort of radiation is caused by the internal composition of human bodies, which includes radioactive potassium-40 and carbon-14[4].

1. Artificial (Man-made) Radiation:

The following are examples of man-made radiation:

- Medical techniques such as diagnostic x-rays, nuclear medicine, and radiation therapy expose people to radiation. Consumer products, such as building materials, combustible fuels (gas and coal), television, and cell phones, are also included in this category.
- Radiation from nuclear sites, which account for less than 0.01 percent of the average annual dose, as well as exposure from radioactive materials shipment and residual fallout from nuclear weapons testing and accidents like Chernobyl.

Biological Effects of Radiation

Biological Effects of Radiation can be broken into two groups according to how the responses (symptoms or effects) relate to dose (or amount of radiation received)

1. The First Group of biological effects are Stochastic Effects
2. The Second Group of biological effects are Deterministic Effects

The first category consists of exposure to low doses of radiation over an extended period of time producing chronic or long term effects (Stochastic) while the second category represents exposure to high doses of radiation over short periods of time producing acute or short term effects (Deterministic). The high doses tend to kill cells, while low doses tend to damage or change them. High doses can kill so many cells that will lead to damage of tissues and organs. This may result to a rapid whole body response often called the Acute Radiation Syndrome (ARS)[13]. The effect of radiation is dependent on many factors including:

- The type of radiation (alpha, beta or gamma)
- The amount received
- The rate at which it is received
- Which part of the body is exposed

- Whether the exposure is chronic (regular, low doses) or acute (short time, high dose)
- The age of the irradiated person.

Dosimetry

The science of "dosimetry" is the measurement, calculation, or combination of measurement and calculation used to calculate radiation exposure. Radiation dosage is also known as "absorbed dose," which is defined as the quantity of radiation energy deposited in tissue divided by the tissue's mass. The most important physical component that impacts the response of tumours and the rest of the body to radiation is the absorbed dosage[6]. When a subject is exposed to radiation, dosimetry provides a method of calculating parameter for risk estimation. Thereby two different situations have to be distinguished: irradiation from an external source and irradiation from activity present inside the body. These two situations require a different approach.

External Source

The following aspects are involved in these calculations:

1. the strength or activity of the radioactive source;
2. the exposure of the subject to such radiation;
3. the dose absorbed by the different tissues;
4. the biological effectiveness of the type of radiation;
5. the weighting of the separate organs

These items will all be discussed subsequently in this section. Since the introduction of the Systeme internationale (SI) as the basis for quantification of scientific measurements, the units for most radiation quantities have been changed. The relation between those SI units and their corresponding conventional is indicated in Table 1

- Activity: The measure of the activity of a radioactive source is the number of disintegration per second. For many years this activity was expressed in Curie (Ci), defined as activity of one gram of radium. Since the introduction of the SI-units, the unit for radioactivity is Becquerel (Bq). It is defined as one integration per second.
- Exposure: The exposure of a subject to radiation depends on the intensity of radiation where that subject is. It is related to the source strength, the distance between the source and the subject, the type and energy of radiation and absorbing properties of the material between the source and subject. The unit for exposure is Roentgen (R) or Coulombs per kilogram of air (C.kg⁻¹).

Measurement	Description	SI unit	Conventional unit	Relationship
Radioactivity	Nuclear transformation (disintegrations per second)	Becquerel (Bq)	Curie (Ci)	$1 \text{ Bq} = 2.7 \times 10^{-11} \text{ Ci}$ $1 \text{ Ci} = 3.7 \times 10^{10} \text{ Bq}$
Exposure	Charge produced in air by γ - or X-rays	Coulombs per kilogram of air ($\text{C} \cdot \text{kg}^{-1}$)	Roentgen (R)	$1 \text{ C} \cdot \text{kg}^{-1} = 3876 \text{ R}$ $1 \text{ R} = 2.58 \times 10^{-4} \text{ C} \cdot \text{kg}^{-1}$
Exposure rate	Observed dose rate in air from a sealed source at a distance of 1m from source	$\text{C} \cdot \text{kg}^{-1} \cdot \text{s}^{-1}$	$\text{R} \cdot \text{s}^{-1}$	$1 \text{ C} \cdot \text{kg}^{-1} \cdot \text{s}^{-1} = 3876 \text{ R} \cdot \text{s}^{-1}$ $1 \text{ R} \cdot \text{s}^{-1} = 2.58 \times 10^{-4} \text{ C} \cdot \text{kg}^{-1} \cdot \text{s}^{-1}$
Absorbed dose	Amount of energy imparted to matter	Gray (Gy)	Rad	$1 \text{ Gy} = 100 \text{ Rad}$ $1 \text{ Rad} = 0.01 \text{ Gy}$
Dose equivalent	Absorbed dose multiplied by modifying factors	Sievert (Sv)	Rem	$1 \text{ Sv} = 100 \text{ Rem}$ $1 \text{ Rem} = 0.01 \text{ Sv}$

Figure 1: Comparison of common radiation terms in SI and conventional units

- **Absorbed Dose:** The absorbed dose (D) is represented by the amount of energy imparted to matter. It is proportional to the exposure and is dependent on the type of matter and the properties of radiation. The SI-units of absorbed dose is the Gray (Gy) corresponding to $\text{J} \cdot \text{kg}^{-1}$. The absorbed dose is usually calculated separately for various organs.
- **Dose Equivalent:** The effect of radiation is not only determined by the absorbed dose but also by the biological effectiveness of the radiation, given as the quality-factor Q. When this has been taken into account, the dose equivalent is obtained. The SI-unit of dose equivalent is Sievert (Sv). The biological effect of a given dose equivalent is independent of the type of the radiation.

Why it is important?

Absorbed dose determines the extent to which tumors and normal tissues are affected by radiation. The higher the absorbed dose to tumors, the more cells will be killed by radiation and the greater the likelihood of a cure. However, the higher the absorbed dose to normal tissues, the more likely and severe may be the undesirable toxic side-effects of the radiation. An important advantage of radiopharmaceutical therapy is its ability to irradiate and effectively treat tumors throughout the body; at the same time, some irradiation of normal organs is unavoidable. Therefore, the role of radiation dosimetry in targeted radionuclide therapy is to determine specifically, for each patient, the administered amount of the radiopharmaceutical that will most effectively treat the patient's disease while avoiding absorbed doses that damage normal tissues. Individualized radiation dosimetry is critical for planning the most effective and safest targeted

radionuclide therapy for each patient.

References

- [1] Manuel D Cerqueira et al. *Recommendations for reducing radiation exposure in myocardial perfusion imaging*. 2010.
- [2] United Nations Scientific Committee on the Effects of Atomic Radiation, B Annex, et al. “Exposures from natural radiation sources”. In: *cosmic rays* 9 (2000), p. 11.
- [3] FN Flakus. “Detecting and Measuring Ionizing Radiation- A Short History.” In: *IAEA bulletin* 23.4 (1982), pp. 31–36.
- [4] J Ford. “Radiation, people and the environment”. In: (2004).
- [5] MA Hapgood. “Towards a scientific understanding of the risk from extreme space weather”. In: *Advances in Space Research* 47.12 (2011), pp. 2059–2072.
- [6] TH Kirby, WF Hanson, and DA Johnston. “Uncertainty analysis of absorbed dose calculations from thermoluminescence dosimeters”. In: *Medical physics* 19.6 (1992), pp. 1427–1433.
- [7] K Kovler et al. “Basic aspects of natural radioactivity”. In: *Naturally Occurring Radioactive Materials in Construction*. Elsevier, 2017, pp. 13–36.
- [8] Mary Alice Statkiewicz Sherer et al. *Radiation protection in medical radiography*. Elsevier Health Sciences, 2013.
- [9] A Signore et al. “Molecular imaging of inflammation/infection: nuclear medicine and optical imaging agents and methods”. In: *Chemical reviews* 110.5 (2010), pp. 3112–3145.
- [10] Michalis Tzortzis et al. “Gamma-ray measurements of naturally occurring radioactive samples from Cyprus characteristic geological rocks”. In: *Radiation Measurements* 37.3 (2003), pp. 221–229.
- [11] René A de Vries et al. “The biological effects of radiation”. In: *International Journal of Risk & Safety in Medicine* 4.2 (1993), pp. 149–165.
- [12] Jin-song Wang, Hai-juan Wang, and Hai-li Qian. “Biological effects of radiation on cancer cells”. In: *Military Medical Research* 5.1 (2018), pp. 1–10.
- [13] Nasiru Imam Zakariya and MTE Kahn. “Benefits and biological effects of ionizing radiation”. In: *Sch. Acad. J. Biosci* 2.9 (2014), pp. 583–591.