

STATISTICS 321: Practice Lab

© Scott Robison 2026, all rights reserved.



Purpose

This lab is designed to prepare you for upcoming quizzes and examinations by practicing the types of reasoning expected in this course. You may use AI tools, notes, textbooks, class handouts, R, and other course resources. The goal is not simply to obtain answers, but to understand what is being expected, why a method works, when a method is appropriate, when a method becomes inefficient, and what assumptions are required.

Expectations

Show reasoning clearly. Use probability notation appropriately. Support claims with mathematical arguments. Explain why methods work and when they apply. If you disagree with a solution, identify the first step where the reasoning fails.

Question 2: Comparing Three Solutions

A fair six-sided die is rolled three times. Let X = number of sixes observed. Determine $P(X = 2)$.

Student A: Uses arrangements $(6, 6, 6^c)$, $(6, 6^c, 6)$, $(6^c, 6, 6)$; counts $\binom{3}{1} = \frac{3!}{1!2!} = 3$ $C_1 = C_1^3 = 3$ locations for the non-six and computes

$$3 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right).$$

Student B: Uses $n(S) = 6^3 = 216$ equally likely outcomes. Counts 15 favourable outcomes and computes

$$\frac{n(A)}{n(S)} = \frac{15}{216}.$$

Student C: Claims

$$P(X = 2) = 3 \left(\frac{1}{6}\right)^2 = \frac{1}{12}$$

because there are three arrangements containing two sixes.

Answer:

- (a) Verify algebraically that Students A and B obtain the same probability.
- (b) Explain precisely what $\binom{3}{1}$ counts.
- (c) Identify where the same counting idea appears in Student B's solution.
- (d) Explain the difference between arrangements and outcomes.

- (e) Evaluate the claim that Student B is more correct because actual outcomes were counted.
- (f) Determine whether Student C is correct.
- (g) Identify the first incorrect step.
- (h) Explain why Student C's answer differs from the correct answer.
- (i) Rewrite Student C's solution correctly.
- (j) Generalize to a 20-sided die rolled 15 times and $P(X = 3)$. Which strategy would you recommend?
- (k) Write the first expression produced by that strategy.
- (l) Does preferring one method imply the other is mathematically incorrect?