

Clinical Trial Protocol: Novel Therapy for Type 2 Diabetes Mellitus

1. Objective

To evaluate the safety and efficacy of GlucoX in adult patients with Type 2 Diabetes Mellitus (T2DM).

2. Inclusion Criteria

- Age between 18 and 75 years.
- Confirmed diagnosis of Type 2 Diabetes Mellitus for at least 6 months.
- HbA1c between 7.0% and 10.5% at screening.
- Stable anti-diabetic treatment regimen for at least 3 months.
- Patient has provided informed written eConsent.

3. Exclusion Criteria

- Diagnosis of Type 1 Diabetes Mellitus.
- History of diabetic ketoacidosis within the past 12 months.
- Severe renal impairment (eGFR < 30 mL/min/1.73m²).
- Significant liver disease or active hepatic dysfunction.
- Pregnant or lactating females.
- Recent cardiovascular event (myocardial infarction or stroke within 6 months).
- Known hypersensitivity to study medication.

4. Study Design

Randomized, double-blind, placebo-controlled, multicenter study lasting 24 weeks.

Participants will receive either GlucoX or placebo in addition to standard care.

5. Study Assessments

- HbA1c measurements at baseline, Week 12, and Week 24.
- Fasting plasma glucose monitoring.
- Body weight and BMI assessment.
- Blood pressure and vital signs.
- Adverse event monitoring throughout the study.

6. Primary Endpoint

- Change in HbA1c from baseline to Week 24.

7. Secondary Endpoints

- Change in fasting plasma glucose.
- Change in body weight.
- Percentage of patients achieving HbA1c < 7.0%.
- Safety and tolerability profile.