

Renvora AI - Agriculture 15 MB Test PDF

Test Page 1

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 2

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 3

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 4

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 5

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 6

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 7

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 8

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 9

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 10

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 11

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 12

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 13

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 14

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 15

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 16

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 17

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 18

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 19

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 20

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 21

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 22

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 23

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 24

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 25

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 26

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 27

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 28

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 29

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 30

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 31

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 32

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 33

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 34

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 35

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 36

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 37

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 38

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 39

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 40

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 41

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 42

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 43

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 44

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 45

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 46

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 47

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 48

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 49

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 50

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 51

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 52

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 53

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 54

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 55

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 56

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 57

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 58

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 59

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 60

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 61

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 62

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 63

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 64

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 65

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 66

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 67

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 68

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 69

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 70

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 71

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 72

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 73

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 74

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 75

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 76

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 77

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 78

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 79

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 80

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 81

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 82

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 83

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 84

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 85

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 86

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 87

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 88

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 89

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 90

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 91

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 92

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 93

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 94

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 95

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 96

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 97

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 98

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 99

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 100

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 101

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 102

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 103

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 104

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 105

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 106

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 107

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 108

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 109

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 110

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 111

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 112

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 113

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 114

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 115

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 116

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 117

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 118

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone

Renvora AI - Agriculture 15 MB Test PDF

Test Page 119

Crop: Wheat

Scientific name: *Triticum aestivum*

Recommended soil pH: 6.0 to 7.5

Typical seed rate: 100 to 125 kg per hectare

Important nutrient: Nitrogen

Common irrigation method: Furrow irrigation

Sowing season: Rabi season

Typical crop duration: 120 to 150 days

Major disease: Rust

Disease management: Use resistant varieties and recommended fungicides

Organic matter benefit: Improves soil structure and water-holding capacity

Drip irrigation benefit: Reduces water loss and delivers water near the root zone