



# Singularity Mosaic

---

“

Welcome to Singularity Mosaic, a project dedicated to exploring the enigmatic universe. We'll analyze astronomical data, focusing on black holes. Our journey will cover the history of astronomy and modern coordinate systems. We'll also dive into fascinating deep-sky objects.

---

“

# Cosmic Timeline

---

Ancient Wisdom  
to New Models

From Ptolemy's geocentric views (where the Earth was taken as the centre of the universe) to Copernicus's heliocentric model (where the Sun was taken as the centre of the universe).

Mapping the  
cosmos

There are three major celestial coordinate systems which map universe. Which are equitorial , alt-az and galactic


Modern  
Astronomical  
eyes

Modern telescopes include the EHT, Hubble Telescope, VLA, ALMA, JWST, Chandra, and others. Are our eyes in trhe vast universe

Cosmic  
phenomenon


Phenomena like quasar, pulsar, blackhole, magnetar, etc. This phenomenas are still making us wonder about our universe

# Three Key Coordinate Systems




## Alt-Az

The horizontal system uses altitude (above horizon) and azimuth (from north). Both change with location and time, making it useful for navigation and telescope pointing.



## Equatorial

Right Ascension (like longitude), Declination (like latitude). Earth-centered, fixed stars. Used for star catalogs, object identification. This system uses the Earth's equator projected onto the celestial sphere. The Ecliptic system's reference plane is Earth's orbit around the Sun.



## Galactic

Galactic longitude and galactic latitude use a Sun-centred system aligned with the Milky Way plane. These coordinates help astronomers map galaxy structure and study the distribution of stars, gas, and dust across our galaxy efficiently. And Sagittarius A is taken as the centre

# Conversions and Applications

## Complex Conversions

Converting between systems requires complex mathematical formulas.

## Streamlined Tools

Tools like Python and online calculators simplify these transformations.

## Telescope Control


Equatorial coordinates are corrected for date and time for accurate tracking using the Horizontal system.

## Mapping the Milky Way

Galactic coordinates reveal the large-scale structures within our galaxy.

## Solar System Understanding

Ecliptic coordinates are essential for tracking and understanding planet movement.



# Telescope Handling

We learned here how to use a telescope and point using it

## 1 Aligning with Polaris

We first align the polarscope on telescope with Polaris

## 2 Balancing the telescope

Balance the telescope by changing the counterweights and also shifting the tube. Such that both the RA and DEC axis are fully free

## 3 Pointing to the object

Change the RA and Dec value of the telescope to that of the object and fine-tune it using the slow motion controls

## 4 Focusing the object


To see the object clearly focus the eyepiece using focuser

# Deep-Sky Objects: Cosmic Zoo



## Neutron Stars

Incredibly dense remnants left after supernova explosions. These incredibly dense remnants form after massive star supernova. They consist mainly of neutrons. PSR J1614-2230 is twice the Sun's mass.




## White Dwarfs

The evolutionary end-stage for most low-mass stars. Small to medium stars become white dwarfs. They exhaust their nuclear fuel. Sirius B is a prime example, Earth-sized but solar mass.

# Pulsars and Quasars: Cosmic Beacons

Explore the universe's most captivating phenomena. Pulsars are rapidly rotating neutron stars, emitting precise beams of radiation. Quasars are incredibly luminous galactic nuclei powered by supermassive black holes. These cosmic beacons offer a glimpse into extreme astrophysics.



# Black Holes: Gravity's Ultimate Victory

- 1
- 2
- 3
- 4

## Gravity's Dominance

Nothing, not even light, escapes a black hole's grip. They form from collapsing massive stars.

## Event Horizon

This boundary marks the point of no return. Beyond it, escape is impossible.

## Supermassive Black Holes


Located at galactic centers, they can be billions of solar masses. Sagittarius A\* is our Milky Way's giant.

## Stellar Black Holes

These form from individual stars, ranging from 3 to 100 solar masses. Cygnus X-1 was one of the first discovered.



# Event Horizon Telescope: Seeing the Unseeable



Historic Black Hole Image  
M87\* was imaged in 2019.

Global Telescope Network  
The EHT uses Very Long Baseline  
Interferometry (VLBI).

Unprecedented Resolution  
Imaging an orange on the moon is  
now possible.

Future Cosmic Targets  
Next, imaging Sagittarius A\* at the  
Milky Way's center.

# How EHT works:

Global Synchronization  
The EHT9s telescopes  
are time-synced using  
hydrogen maser atomic  
clocks.

Central Data Correlation  
Data is flown (literally!)  
to supercomputing  
centers (MIT, Max  
Planck) for processing.

Petabytes of Data  
Each site records raw  
radio signals onto  
physical hard drives 4  
up to 4 PB total.

Image Reconstruction  
Powerful algorithms  
turn noisy, sparse data  
into a visible black hole

# Interferometry: Virtual Telescopes

Interferometry combines signals from multiple telescopes to create a single, larger virtual telescope. This technique significantly improves resolution and sensitivity. Very Long Baseline Interferometry (VLBI) is a powerful application, combining distant observatories.

---



## Signal Combination

Individual telescope signals are merged.



## Resolution Boost

Achieving finer details in celestial objects.



## Enhanced Sensitivity

Detecting fainter cosmic sources.



## VLBI in Practice

Data from ALMA and the South Pole Telescope are combined.

# UV Coverage and Fourier Transforms

## The UV Plane


This plane represents the spatial frequencies sampled by an interferometer. Each baseline provides a point.

## Fourier Transform

This mathematical tool converts raw UV data into a celestial image. It's the core of image reconstruction.

## Incomplete Coverage Challenges

When UV coverage is sparse, advanced image reconstruction techniques are necessary.



# From Frequency to Images

## Spatial Frequencies

The Fourier Transform decomposes an image into its constituent spatial frequencies.

## Convolution Theorem


This theorem simplifies complex calculations in the frequency domain. The observed image is therefore a convolution of the true sky brightness with the point spread function (PSF) or synthesized beam.

## Applications

Used in image filtering, compression, and reconstruction. MRI image formation is a prime example.


## Amplitude & Phase

Both amplitude and phase components encode vital image information.




# What is Dirty Image?

- The first image produced by a radio interferometer (array of radio telescopes).
- Created by taking the inverse Fourier transform of measured data ("visibilities").
- Not a true sky image<sup>4</sup>contains distortions and artifacts.




## Why it is Dirty?

- The image is a mix of the real sky and unwanted patterns (sidelobes, noise).
- These patterns come from the **Dirty Beam**.



# What is Dirty Beam?



- The “*point spread function*” (PSF) of the interferometer array.
- Shows how a single point in the sky would appear in the dirty image.
- Has a sharp central peak but many sidelobes (ripples and negative regions).
- Caused by incomplete and uneven sampling of the sky by the telescope array.

# **CLEAN Algorithm: Refining the Image**

CLEAN is a foundational algorithm in radio astronomy imaging, designed to remove artifacts caused by incomplete data from radio interferometers. By iteratively identifying and subtracting the effects of the instrument's response (the "dirty beam"), CLEAN enables astronomers to reconstruct much clearer and more accurate images of the radio sky.

## **Two Cycles for CLEAN Algorithm**


**Major  
Cycle**

**Minor  
Cycle**

# How CLEAN Algorithm works

Find the Peak: Locate the brightest pixel in the current residual image

Record Component: Add a fraction (gain factor) of this peak to a clean component list



**Subtract Beam:** Remove a scaled version of the dirty beam pattern centered on the peak location

**Iterate:** Repeat until stopping criteria are met (threshold or maximum iterations)

**Restore:** Convolve clean components with an idealized "clean beam" and add residuals

# Image Cleaning Process

## Minor Cycle


### Image Domain Processing:

- CLEAN components are searched for and subtracted from the image
- Only the central portion of the dirty beam is used for subtraction
- Operates on a list of the largest residuals (not the full image)
- Much faster than using the full PSF for each subtraction
- The loss in accuracy is the trade-off price we pay to cut down the computational cost.

## Major Cycle

### Visibility Domain Correction:


- The list of CLEAN components from minor cycles are processed through Fast-Fourier-Transformation.
- Components are subtracted from the original visibility data.
- A new residual image is computed from updated visibilities.
- Corrects for errors that may have accumulated during minor cycle approximations.



# CASA: Data Processing and Visualization

## CASA Installation

Installing Common Astronomy Software Applications is the first step.



The screenshot shows a terminal window titled 'Search Message' with a log of CASA startup messages. The log includes details about the Python version (3.12.4), CASA version (CASA/LITH 6.7.0.31), and various configuration parameters like 'cachadir' and 'casaconfig\_verbose'. It also mentions the loading of a configuration file from '/Applications/CASA.app/Contents/Frameworks/Python.framework/Versions/3.12/lib/python3.12/site-packages/casaconfig/config.py'. The log concludes with the loading of 'casarundata' and 'measures' tables, and information about IERS tables and TAI UTC.

```
2025-06-12 18:37:29 INFO ::casa The latest version is already installed in /Users/preetvaru/.casa/data
2025-06-12 18:37:33 INFO ::casa measures_update ... acquiring the lock ...
2025-06-12 18:37:35 INFO ::casa lock file is not empty : /Users/preetvaru/.casa/data/data_update.lock
2025-06-12 18:37:35 SEVERE ::casa The lock file at /Users/preetvaru/.casa/data is not empty.
2025-06-12 18:37:35 SEVERE ::casa Remove the lock file and set force to True with the desired version (default to most recent).
2025-06-12 18:37:35 SEVERE ::casa It may be best to completely repopulate path using pull_data and measures_update.
2025-06-12 18:37:35 SEVERE ::casa lock file is not empty : /Users/preetvaru/.casa/data/data_update.lock
2025-06-12 18:37:35 SEVERE ::casa visit https://casadocs.readthedocs.io/en/stable/notebooks/external-data.html for more information
2025-06-12 18:37:35 SEVERE ::casa
2025-06-12 18:37:35 INFO ::casa Python version 3.12.4
2025-06-12 18:37:36 INFO ::casa CASA Version CASA/LITH 6.7.0.31
2025-06-12 18:37:36 INFO ::casa optional configuration file not found, continuing CASA startup without it
2025-06-12 18:37:36 INFO ::casa loaded config file : /Applications/CASA.app/Contents/Frameworks/Python.framework/Versions/3.12/lib/python3.12/site-packages/casaconfig/config.py
2025-06-12 18:37:36 INFO ::casa
2025-06-12 18:37:36 INFO ::casa config values
2025-06-12 18:37:36 INFO ::casa agg = False
2025-06-12 18:37:36 INFO ::casa cachadir = '/Users/preetvaru/.casa'
2025-06-12 18:37:36 INFO ::casa casaconfig_verbose = 1
2025-06-12 18:37:36 INFO ::casa colors = 'Neutral'
2025-06-12 18:37:36 INFO ::casa data_auto_update = True
2025-06-12 18:37:36 INFO ::casa datapath = ['/Users/preetvaru/.casa/data']
2025-06-12 18:37:36 INFO ::casa iplot = False
2025-06-12 18:37:36 INFO ::casa ipythonfile = '/Users/preetvaru/ipython-20250612-183726.log'
2025-06-12 18:37:36 INFO ::casa logger = False
2025-06-12 18:37:36 INFO ::casa logfile = '/Users/preetvaru/casa-20250612-183726.log'
2025-06-12 18:37:36 INFO ::casa measures_auto_update = True
2025-06-12 18:37:36 INFO ::casa measurespath = '/Users/preetvaru/.casa/data'
2025-06-12 18:37:36 INFO ::casa nogui = False
2025-06-12 18:37:36 INFO ::casa nolog = False
2025-06-12 18:37:36 INFO ::casa nologger = False
2025-06-12 18:37:36 INFO ::casa pipeline = False
2025-06-12 18:37:36 INFO ::casa startupfile = '/Users/preetvaru/.casa/startup.py'
2025-06-12 18:37:36 INFO ::casa user_site = False
2025-06-12 18:37:36 INFO ::casa
2025-06-12 18:37:36 INFO ::casa casarundata version : casarundata-2025.05.15-1.tar.gz
2025-06-12 18:37:36 INFO ::casa measures version : NSRT_Measures_20250605-160001.star
2025-06-12 18:37:36 INFO ::casa
2025-06-12 18:37:36 INFO ::casa Checking Measures tables in data repository sub-directory /Users/preetvaru/.casa/data/geodetic
2025-06-12 18:37:36 INFO ::casa IERSep2000 (version date, last date in table (UTC)) : 2025/05/19/15:15, 2025/04/19/00:00:00
2025-06-12 18:37:36 INFO ::casa IERSep97 (version date, last date in table (UTC)) : 2025/05/19/15:15, 2025/04/19/00:00:00
2025-06-12 18:37:36 INFO ::casa IERSpredict (version date, last date in table (UTC)) : 2025/06/15/15, 2025/09/03/00:00:00
2025-06-12 18:37:36 INFO ::casa TAI UTC (version date, last date in table (UTC)) : 2025/05/11/15:15, 2017/01/01/00:00:00
```

## Loading EHT Data

EHT data is loaded and calibrated within CASA for analysis.

## Visualizing Black Holes

The software enables detailed visualization of black hole images and models.

# Visualising black holes

1. The ‘importuvfits’ command converts raw UVFITS files into CASA’s Measurement Set (MS) format :

```
importuvfits(fitsfile="SR1_M87_2017_095_lo_hops_netcal_StokesI.uvfits", vis  
="M87_095.ms")  
importuvfits(fitsfile="SR1_M87_2017_096_lo_hops_netcal_StokesI.uvfits", vis  
="M87_096.ms")  
importuvfits(fitsfile="SR1_M87_2017_100_lo_hops_netcal_StokesI.uvfits", vis  
="M87_100.ms")  
importuvfits(fitsfile="SR1_M87_2017_101_lo_hops_netcal_StokesI.uvfits", vis  
="M87_101.ms")  
importuvfits(fitsfile="SR1_M87_2017_095_hi_hops_netcal_StokesI.uvfits", vis  
="M87_095_hi.ms")  
importuvfits(fitsfile="SR1_M87_2017_096_hi_hops_netcal_StokesI.uvfits", vis  
="M87_096_hi.ms")  
importuvfits(fitsfile="SR1_M87_2017_100_hi_hops_netcal_StokesI.uvfits", vis  
="M87_100_hi.ms")  
importuvfits(fitsfile="SR1_M87_2017_101_hi_hops_netcal_StokesI.uvfits", vis  
="M87_101_hi.ms")
```

2. Use ‘listobs’ to examine the MS structure and metadata:

```
listobs(vis="M87_095_lo.ms")
```

3. The ‘plotms’ tool is used to visualise the UV-coverage, revealing the array geometry:

```
plotms(vis="M87_095_lo.ms", xaxis="u", yaxis="v", avgtime="1e8", avgscan=  
True, coloraxis="spw")
```

4. ‘mstransform’ is used to generate corrected MS files, resolving pointing table inconsistencies:

```
mstransform(vis="M87_2017_095_lo.ms", outputvis="M87_2017_095_lo_fixed.ms  
", datacolumn="data")  
mstransform(vis="M87_2017_096_lo.ms", outputvis="M87_2017_096_lo_fixed.ms  
", datacolumn="data")  
mstransform(vis="M87_2017_100_lo.ms", outputvis="M87_2017_100_lo_fixed.ms  
", datacolumn="data")  
mstransform(vis="M87_2017_101_lo.ms", outputvis="M87_2017_101_lo_fixed.ms  
", datacolumn="data")
```

6. Multiple MS files are merged using concat, preserving field names and time order:

```
concat(vis=[ "M87_2017_095_lo_fixed.ms", "M87_2017_096_lo_fixed.ms",  
"M87_2017_100_lo_fixed.ms", "M87_2017_101_lo_fixed.ms"],  
concatvis="M87_combined_distinct.ms",  
timesort=True, respectname=True, copypointing=False)
```

5. Unique field names are assigned before concatenation:

```
tb.open("M87_2017_095_lo_fixed.ms/FIELD", nomodify=False)  
tb.putcol("NAME", ["M87_095"])  
tb.close()  
tb.open("M87_2017_096_lo_fixed.ms/FIELD", nomodify=False)  
tb.putcol("NAME", ["M87_096"])  
tb.close()  
tb.open("M87_2017_100_lo_fixed.ms/FIELD", nomodify=False)  
tb.putcol("NAME", ["M87_100"])  
tb.close()  
tb.open("M87_2017_101_lo_fixed.ms/FIELD", nomodify=False)  
tb.putcol("NAME", ["M87_101"])  
tb.close()
```

7. To plot the final image :

```
plotms(vis="M87_combined_distinct.ms", xaxis="u", yaxis="v",  
coloraxis="field", avgchannel="9999", avgtime="9999")
```

# Python Assignment

```
[1] from google.colab import drive
drive.mount('/content/drive')

→ Mounted at /content/drive

[2] with open('/content/drive/MyDrive/Moons_and_Planets.csv','r') as f:
    lines = f.read().split('\n')
    data_moons = []
    for line in lines:
        data_moons.append(line.split(','))
    print(data_moons)

→ [[ '# Name of Moon', ' Name of Planet', ' Diameter (km)' ], [ 'Moon', 'Earth', '1737.1' ], [ 'Phobos', 'Mars', '11.1' ], [ 'Deimos', 'Mars', '6.2' ], [ 'Io', 'Ju... ]]

[3] planets = ('Earth','Mars','Jupiter','Saturn','Uranus','Neptune','Pluto')
for planet in planets:
    count = 0
    for moon in data_moons[1:]:
        if len(moon) > 1 and moon[1].strip() == planet:
            count += 1
    print(f'{planet} has {count} moons')


→ Earth has 1 moons
Mars has 2 moons
Jupiter has 79 moons
Saturn has 82 moons
Uranus has 27 moons
Neptune has 14 moons
Pluto has 5 moons

[4] dis = len(data_moons)
for i in range(1,dis):
    for j in range(1,dis-i-1):
        if float(data_moons[j][2]) < float(data_moons[j+1][2]):
            temp = data_moons[j]
            data_moons[j] = data_moons[j+1]
            data_moons[j+1] = temp
for moon in data_moons:
    print(moon)


→ [ '# Name of Moon', ' Name of Planet', ' Diameter (km)' ]
[ 'Ganymede', 'Jupiter', '2634.1' ]
[ 'Titan', 'Saturn', '2575.5' ]
[ 'Callisto', 'Jupiter', '2408.4' ]
[ 'Io', 'Jupiter', '1818.1' ]
[ 'Moon', 'Earth', '1737.1' ]
[ 'Europa', 'Jupiter', '1560.7' ]
```

# Bulla Session

Team A : Intro to Warp Drive



Team B : Information paradox



Team C: Brief info on Milky Way



**THANK YOU**