OPTI 517 — Homework #4


Saruul Nasanjargal

Hubble Space Telescope

Part 1

The Hubble Space Telescope is a Ritchey–Chrétien. Figure 1 shows the lens data. I put the object at infinity and set the aperture to an entrance pupil diameter of 2400 mm. The stop is a dummy surface in contact with the primary. The primary mirror has radius −11040 mm and conic −1.002, and the thickness after it is −4906.072 mm, so the secondary lies that far in front of the primary. The secondary has radius −1358 mm and conic −1.496. Both conics are below −1, so both mirrors are hyperboloids. A marginal-ray solve on the secondary thickness places the image 6406.090 mm after the secondary, which is 1500.018 mm behind the primary. The clear semi-diameter at the secondary is 133.628 mm.

Figure 2 shows the Cassegrain path. Collimated rays strike the primary, reflect to the secondary, and come to a focus behind the primary. The scale bar is 2×10^3 mm.



	Surface Type		Radius	Thickness	Material	Coating	Clear Semi-Dia	Conic	TCE x 1E-6
0	OBJECT	Standard	Infinity	Infinity			0.000	0.000	0.000
1		Standard	Infinity	5000.000			1200.000	0.000	0.000
2	STOP (aper)	Standard	Infinity	0.000			1200.000	0.000	0.000
3	(aper)	Standard	-1.104E+04	-4906.072	MIRROR		1200.000	-1.002	0.000
4		Standard	Infinity	0.000			135.073	0.000	0.000
5	(aper)	Standard	-1358.000	6406.090 M	MIRROR		133.628	-1.496	0.000
6	IMAGE	Standard	Infinity	-			5.194E-03	0.000	0.000

Figure 1: Hubble lens data. Entrance pupil diameter 2400 mm, stop on the primary, and both mirrors hyperbolic.




Figure 2: Hubble layout. The secondary is to the left of the primary, and the image is behind the primary.

With spherical aberration and coma removed, Figure 3 is the flat-image-surface case. The on-axis optical path difference is flat. At 0.25° both fans are strongly curved, and the plot scale is ±20 waves. The RMS wavefront error crosses the diffraction limit at about 0.027°.




Figure 3: Flat image surface. On-axis OPD is flat; at 0.25° the scale is ±20 waves. The RMS wavefront error leaves the diffraction limit near 0.027°.

The Seidel coefficients give a Petzval radius of −774.237 mm.[1] I entered that value as the image-surface radius. Figure 4 is the updated pair of plots. The off-axis fans then fit on a scale of ±5 waves, and the diffraction-limited field extends to about 0.0573°. The on-axis fan stays flat. The curved surface takes out the Petzval defocus, and the field that remains sharp is set by the residual astigmatism.




Figure 4: Image surface radius set to the Petzval radius, −774.237 mm. The off-axis OPD scale is ±5 waves, and the diffraction-limited field extends to about 0.0573°.

Part 2

Part 3

Part 4

Part 5

[1]The Petzval radius is the radius of the surface the image would lie on if astigmatism were absent. It comes only from the surface powers and the refractive indices. Putting that radius on the image surface means the detector is curved to match it. Astigmatism is still left.

Schmidt, Maksutov, and Houghton Cameras

Part 6

Part 7

Part 8

Part 9

I verify that this HW is all my work.

Signature: _____ Saruul Nasanjargal