Dimension bounds for hyperbolic Coxeter polytopes

Alexander Kolpakov

4 November 2026

Abstract

We prove that a finite-volume hyperbolic Coxeter polytope has dimension at most 57, that a compact one has dimension at most 17, and that a compact arithmetic one has dimension at most 15. The finite-volume argument compares corner weights with the geometry of vertex links. The compact arguments use projections of one vector associated with an original farthest vertex; arithmeticity further restricts the ground field and integral wall parameters. We develop the geometric reductions, reference-vector estimates and terminal arithmetic arguments in detail. The complete local inventories, remaining case proofs, computational proofs and reproduction instructions are supplied as versioned addenda in the accompanying repository.

1 Introduction

All polytopes in this article are convex, finite-sided and full-dimensional. A hyperbolic Coxeter polytope has dihedral angles that are integer submultiples of π; reflection in its facets generates a discrete group with the polytope as fundamental domain. Hyperbolic space has curvature -1.

Theorem 1.1. *Let $P \subset \mathbb{H}^n$ be a finite-volume hyperbolic Coxeter polytope. Then:*

(i) $n \leq 57$;
(ii) *if P is compact, then* $n \leq 17$;
(iii) *if P is compact and its reflection group is arithmetic, then* $n \leq 15$.

The classical finite-volume ceiling is 995, due to Prokhorov [1], using the combinatorial estimates of Khovanskii [2]. Vinberg obtained the compact ceiling 29 [3, 4]. The largest known finite-volume examples have dimension 21 [5]; the bounds here are upper bounds and make no new existence assertion. Compactness is essential in Theorem 1.1(iii).

Guo and Zheng recently obtained the general finite-volume bound 81 [6]. An earlier version of our bound 57, including its corner-weight method and higher-dimensional argument, was publicly posted on 21 October 2026 [7]. The present article condenses the three proof families assembled in the repository edition of 30 October [8].

The methods separate after the common local geometry. In the finite-volume case, triangles and quadrilaterals require corner payments that high-dimensional vertex diagrams cannot accommodate. Nonsimple ideal vertices are treated through the product structure of their links. In the compact case, local walls force quadratic lower bounds for components of a single vector of norm less than one. Arithmeticity enters only in the third argument, through integral cycles and positivity at the other real embeddings.

Detailed proofs and version. The full proof addenda [8] are fixed at repository commit `36eeaddf646eff25d4990c9ab730a20ada21f80f`. Parts I and IV contain the finite-volume argument,

Parts II and V the compact argument, and Parts III and VI the arithmetic argument. We refer to these as Addenda FV, C and A, respectively. The paper includes the common geometric lemmas, the principal local reductions and the final contradictions. The exhaustive inventories and remaining case proofs are cited at that fixed version; Section 6 gives the proof-catalogue entry points and reproduction package. The supplementary case analyses remain necessary parts of the proof, with stable references independent of subsequent research. In particular, the finite-volume local coverage and exact forest maximum, the complete compact component dispatch, and the arithmetic component inventories retain the specific online dependencies stated where they are used.

Contents

1 Introduction **1**

2 Actual walls and vertex diagrams **3**

3 The finite-volume bound **4**
- 3.1 Actual cores, extensions and whole components . . . 5
- 3.2 The finite forest algorithm . . . 7
- 3.3 The dimension-specific final local rule . . . 8
  - 3.3.1 Selected cases and arbitrary additional supporting walls . . . 8
  - 3.3.2 The affine-D12 case . . . 9
  - 3.3.3 Retention of other payments . . . 10
- 3.4 The finite maximum after the transfer . . . 10
- 3.5 Ideal vertices and positive surplus . . . 11
  - 3.5.1 The terminal domain and its exact output . . . 11
- 3.6 Shape curvature and the contradiction in dimension 58 . . . 13
- 3.7 The bounded higher-dimensional range . . . 14
  - 3.7.1 The elementary families in dimensions 98–163 . . . 14
  - 3.7.2 The remaining range 59–97 . . . 15
- 3.8 The unbounded tail: actual framing walls and two budgets . . . 16
  - 3.8.1 Outgoing edges give actual framing walls . . . 16
  - 3.8.2 The common-height Lorentz identity . . . 17
  - 3.8.3 Point potentials and the original endpoint budgets . . . 17
  - 3.8.4 Exceptional factors and ordered residual packing . . . 19
  - 3.8.5 The other height range . . . 20
  - 3.8.6 A narrow facet without a cusp . . . 21

4 The compact bound **22**
- 4.1 Farthest vertices and projections . . . 22
  - 4.1.1 Positive inverse matrices and feasible projections . . . 23
  - 4.1.2 Opposite walls and common components . . . 24
- 4.2 Quadratic estimates from reference vectors . . . 25
- 4.3 Component estimates . . . 26
  - 4.3.1 Ordinary components and exceptional actual defects . . . 26
  - 4.3.2 Proofs of the small defects and rank-two costs . . . 27
  - 4.3.3 The short directions and their exact debits . . . 27
  - 4.3.4 The two rank-two estimates . . . 28
  - 4.3.5 The A-chain branch . . . 29

4.3.6 The common B/D prefix branch . . . 29
4.4 The remaining spherical diagrams . . . 30
4.4.1 A fresh cover in exact dimension 18 . . . 30
4.4.2 The separate complete higher-dimensional tail . . . 31
4.5 The whole-B18 terminal partition . . . 32
4.6 The last case: an actual whole-D18 vertex . . . 32
4.6.1 A future point valid for every dotted head . . . 33
4.6.2 The four original reference cases . . . 33
4.7 Bounds for the head and fork edges . . . 34
4.7.1 The first three rows . . . 34
4.7.2 The weak-helper row: two separate anchor prices . . . 35
4.7.3 Both signed fork directions . . . 36
4.8 A local A18 lemma at an adjacent actual vertex . . . 37
4.9 The near/far pair and the final composition . . . 39
4.9.1 Every actual near/far fork edge is short . . . 39
4.9.2 The entire near/far family . . . 40
4.9.3 Exhausting the actual head and fork covers . . . 40
4.10 Finite algorithms and conditional verification . . . 41
4.10.1 The mathematical finite-cover calculation . . . 41

5 The compact arithmetic bound 42
5.1 The actual arithmetic field . . . 42
5.2 Paying all isolated constraints together . . . 42
5.3 The actual golden long-end star and invariant lattices . . . 43
5.4 The complete rank-seventeen reduction . . . 45
5.4.1 Shared costs and arithmetic prefix seals . . . 45
5.5 The four finite algebraic propositions . . . 52
5.5.1 Full quadratic integer rings . . . 52
5.6 The complete rank-sixteen reduction . . . 53
5.6.1 A small component calculation . . . 54
5.7 The last actual diagram . . . 56
5.7.1 Norm bounds, integer orders and valuations . . . 61
5.7.2 Three complete integer-strip lemmas . . . 62
5.8 Odd head norm and inert two . . . 62
5.8.1 Odd norm . . . 62
5.8.2 Even norm with inert two . . . 63
5.9 Even head norm with ramified two . . . 63
5.10 The complete split branch . . . 64
5.10.1 Odd trace . . . 64
5.10.2 The sole even-trace head . . . 65

6 GitHub proof catalogue and reproduction 66

2 Actual walls and vertex diagrams

We use a Lorentz space of signature $(-,+,\dots,+)$ and the future unit hyperboloid model of $\mathbb{H}^n$. Outward wall normals are spacelike; the chamber inequalities are $\langle y, e_i\rangle \le 0$. Distinct wall normals

have nonpositive products. Finite vertex diagrams are spherical. A horospherical link at an ideal vertex is a product

$$L_v = \prod_i \Delta^{r_i}, \qquad \sum_i r_i = n - 1, \tag{2.1}$$

whose factors correspond to irreducible affine diagrams of orders $r_i + 1$. These standard face and link facts are used in their actual-wall form [4, 9].

Two elementary observations account for much of the local geometry.

Lemma 2.1. *A positive-definite Gram matrix with nonpositive off-diagonal entries has entrywise nonnegative inverse. A positive-definite principal set of actual chamber walls has a feasible finite common point.*

Proof. If B is the matrix, choose s larger than its largest eigenvalue. Then $M = sI - B$ is entrywise nonnegative and has spectral radius less than s, so $B^{-1} = s^{-1} \sum_{j \geq 0} (M/s)^j$.

Take a strict interior future timelike point o and put $b_i = -\langle o, e_i \rangle > 0$ for the selected walls. The vector

$$q = o + \sum_i (B^{-1} b)_i e_i$$

is orthogonal to the selected normals and has square $q^2 = o^2 - b^T B^{-1} b < 0$. Its product with o is negative, so it is future pointing. Every other chamber inequality is strict, because the coefficients are nonnegative and all other wall products are nonpositive. Normalize q to the hyperboloid. □

An irreducible affine principal set has a positive kernel vector (u_i). The root combination $\nu = \sum_i u_i e_i$ is null, nonzero since $\langle o, \nu \rangle < 0$, and feasible against every wall. It represents an actual ideal point. Irreducibility and the positive definiteness of proper deletions identify its whole affine factor. In contrast, a kernel vector in an indefinite core represents the zero vector: that vector is null and orthogonal to a timelike vector in the core span, whose perpendicular space is positive definite. Both interpretations are needed when controlling extensions.

At an actual two-face, the supporting walls form a positive-definite diagram of rank $n-2$. Adding one side gives an edge diagram, and adding two adjacent sides gives a positive-semidefinite corner diagram. For opposite sides of a quadrilateral, the corresponding principal diagram is indefinite. A positive or affine alternative would give a feasible common finite or ideal point of those sides. Minimal indefinite subdiagrams, inverse Gram matrices and the classification of finite and affine diagrams consequently reduce the local payment questions to explicit marked configurations.

3 The finite-volume bound

The finite-volume argument has two distinct geometric tasks. In dimension 58 we compare payments at the corners of actual two-faces with the curvature of their vertex links. The exclusion of all larger dimensions uses its own dimension-specific rules and then an all-rank narrow-facet argument. In particular, an exclusion in one dimension is not passed to an arbitrary facet: the induced angles of that facet have not thereby been proved Coxeter.

We give the common geometric reductions, the change in the local rule which achieves 57, the complete concentration argument for nonsimple cusps, and the geometric construction behind the unbounded tail. The finite tables defining the full marked rule, exhaustive route classifications and exact computation certificates remain in Addendum FV. Catalogue entries FV1–FV4 specify those proof dependencies; below a computed interface is identified as such before it is used.

For a two-face F with $s(F)$ sides, set

$$\mathcal{D}_5(P) = \sum_{\dim F=2} (5 - s(F)).$$

A corner means an incidence (v, F) of a vertex and an actual two-face. A nonnegative rule assigns it a weight $w(v, F)$; write $C(v) = \sum_{F \ni v} w(v, F)$.

Lemma 3.1 (Payment comparison)**.** *Suppose every triangular face receives total payment at least 2 and every quadrilateral receives at least 1. Then*

$$\mathcal{D}_5(P) \le \sum_v C(v).$$

More generally, suppose an incidence argument gives

$$\mathcal{D}_5(P) \ge Bv_f + \sum_{v \text{ ideal}} \kappa(v) + \Delta, \qquad \Delta > 0,$$

while a simultaneous payment rule has $C(v) \le B$ at every finite vertex and $C(v) \le \kappa(v) - \varepsilon$ at every ideal vertex, where $\varepsilon \ge 0$. These hypotheses are inconsistent.

Proof. A face with s sides needs payment at least $5 - s$. This is 2 for a triangle and 1 for a quadrilateral. For $s \ge 5$ it follows from nonnegativity. Sum over faces and then over their corners. Combining the resulting upper bound with the proposed lower bound gives $0 \ge \Delta + \varepsilon v_i > 0$. □

3.1 Actual cores, extensions and whole components

We use outward norm-two roots, with diagonal Gram entries 2 and nonpositive off-diagonal entries. At an actual two-face in dimension n, the supporting white roots form a positive-definite rank $n-2$ diagram. White roots together with one black side give a strictly positive-definite edge diagram; adjacent corners are positive semidefinite. A finite corner has an elliptic forest of rank n.

These assertions refer to actual faces of the original chamber. By Lemma 2.1, a positive principal set has an actual finite common point, strict against all other walls. An irreducible affine set has a positive root relation giving a nonzero feasible null vector, and therefore an actual ideal point. Its affine component is a whole factor of that cusp: every proper principal subdiagram is positive definite. These feasibility statements are what allow us to use a formal core inside an actual two-face; see also [9, Section 1].

Lemma 3.2 (Opposite sides and minimal cores)**.** *For either opposite pair of sides of an actual quadrilateral, the supporting whites and that pair form an indefinite principal set. An inclusion-minimal indefinite subset contains both marked sides. Its order is at most ten; if all proper principal subsets are elliptic, its order is at most five.*

Proof. A positive set would have an actual common finite point by the projection argument. A positive-semidefinite singular set would have a common ideal point by its affine kernel. Opposite closed sides of a nondegenerate quadrilateral are disjoint, so neither is possible. Deleting either marked side leaves a subset of an actual finite edge diagram. Hence both marks lie in every minimal indefinite subset. The published Lannér/quasi-Lannér order bounds now apply; the classification through order ten is [10]. □

Two such opposite cores cannot be disjoint and orthogonal, since that would give two negative directions. Their overlap or a joining path supplies an adjacent-side route. The local theorem must then restore all the other supporting whites; an argument on the displayed cores alone would not justify a whole-component payment.

A whole component is a connected component of the full actual corner diagram, not merely of a displayed core. Presence conditions in a weight refer only to such whole components. This distinction is essential when additional supporting whites are restored.

Lemma 3.3 (Two ways to seal a component)**.** *Let G be the Gram matrix of an actual indefinite core.*

1. *A Gram-kernel vector z of this core gives the actual zero root relation $\sum_i z_i e_i = 0$. Every further wall therefore satisfies $\sum_i z_i(e, e_i) = 0$.*
2. *Suppose G is nonsingular, an added forest contacts it only at a position p, and $(G^{-1})_{pp} = r > 1$. If every connected component of the added forest contacts p, its total rank is at most $1/(r-1)$.*

Proof. The span of an indefinite core contains a timelike vector. The root combination associated with a Gram-kernel vector is null and orthogonal to that timelike vector. The perpendicular space is positive definite, so the combination is zero. This differs from the nonzero feasible null vector of an affine block.

For the second statement let B be the positive Gram matrix of all added supporting whites, including indirect tails, and let u be their contact column at p. Schur elimination and the one-negative- direction condition give $B - ruu^T \succeq 0$, whence $u^T B^{-1} u \le 1/r$.

For a connected square-two finite Coxeter Gram of rank k, every inverse diagonal is at least $k/(k+1)$. This follows by induction: delete the chosen node, let the remaining component ranks be k_j, and use a contact of magnitude at least one into each component. Inverse positivity gives Schur cost at least $\sum_j k_j/(k_j+1) \ge (k-1)/k$. The resulting inverse diagonal is at least $1/(2-(k-1)/k) = k/(k+1)$; the rank-one base is $1/2$. For an added forest of total rank M contacted in each component, the same bound gives $u^T B^{-1} u \ge M/(M+1)$. Thus $M/(M+1) \le 1/r$, proving $M \le 1/(r-1)$. □

The second argument takes all roots of the growing forest at once. A bound on its immediate neighbors would leave indirect extensions uncontrolled. For a radical seal, nonpositive contact signs and the displayed zero relation restrict every further root; the local short-path alternatives then force either a persistent payment or orthogonality. An indirect extension has a first contacting root, so this is the step that turns a core component into a whole one.

At an ideal vertex with affine factor ranks r_i, the link is $\prod_i \Delta^{r_i}$, with $\sum r_i = n-1$. A marked pair in factor i occurs in $\prod_{j \ne i}(r_j+1)$ actual corner fibers. Hence an affine factor envelope $(r+1)c(r)$ gives

$$C_{\mathrm{act}} \le C_{\mathrm{up}} = V \sum_i c(r_i), \qquad V = \prod_i (r_i + 1). \tag{3.1}$$

The factor-fiber theorem includes nonsimple ideal vertices. Whole affine factors used locally are established by the positive kernel and actual corner semidefiniteness, before applying this multiplicity. Indeed an edge of the product link varies in just one simplex factor. A fixed pair of its facet labels determines an edge in that factor and an arbitrary vertex in every other factor. These are precisely the $V/(r_i+1)$ fibers. The marked pair restores the varying affine factor; all other factors appear with one node deleted. Summing an unweighted copy of each affine table would miss this multiplicity.

The single intrinsic marked rule is fixed in Addendum FV, "The intrinsic corner weights" (catalogue FV1). It uses same-component weights, ordinary cross-component weights and contextual

additions at finite vertices, and the varying affine-factor weights at ideal vertices. Every guard refers to whole actual components, with the stated priority among overlapping roles. Its preliminary version restores the B_{13} endpoint unit and omits the three rank-15 changes stated below. This specification fixes which tables enter each of the two finite interfaces; changing a charge in one row without retaining the other local payments would not give a simultaneous rule.

3.2 The finite forest algorithm

For the finite calculation multiply charges by 64. Its alphabet is

$$\{A_r : 1 \le r \le 58\} \cup \{B_r : 2 \le r \le 58\} \cup \{D_r : 4 \le r \le 58\} \cup \{E_6, E_7, E_8, F_4, H_3, H_4, I_2^*\}.$$

The rank-two symbol I_2^* represents all remaining $I_2(m)$, $m \ge 5$, in the cost calculation. The local all-label theorem gives their common relevant charge and interactions; this notation is not a finite cutoff on m or an identification of their metrics. An admissible formal forest has multiplicities $m_T \ge 0$ with $\sum_T \operatorname{rank}(T) m_T = 58$. Actual finite vertices are among these forests. The calculation needs no converse realization theorem.

The ordinary cost consists of a component table $b(T)$ and a symmetric interaction table $j(T, U)$; context bonuses are separate. In particular $j(T, T)$ concerns two distinct copies of T. The cost is

$$C_{64}(m) = \sum_T b(T) m_T + \sum_T j(T,T) \binom{m_T}{2} + \sum_{T<U} j(T,U) m_T m_U + \Psi(m). \tag{3.2}$$

The tables and the context function belong to the simultaneous local rule. Their identification with actual corner charges is a geometric obligation, separate from maximizing this expression. Addendum FV, "Geometric coverage and exact local certificates" supplies the actual-face coverage and triangle arguments. Addendum FV, "Contexts in the finite-forest calculation" specifies the complete context function and the finite refinement used in this calculation.

The exact maximization fixes all context-sensitive counts before solving an additive residual problem. With positive integer ranks $r(T)$ and integer weights $w(T)$, its recurrence is

$$V(0) = 0, \qquad V(s) = \max_{r(T) \le s} \{V(s - r(T)) + w(T)\},$$

where only reachable predecessors are admitted. Induction on s proves the upper bound by removing a component and the lower bound by appending one to an attaining predecessor. For a saved certificate, every allowed last-item inequality and a tight witness at every reachable positive rank are required. Checking only an attaining path would not bound omitted forests. The complete context split, residual domains, algorithms and computational proofs are supplied by catalogue FV2, under "Contexts in the finite-forest calculation" in Addendum FV.

Proposition 3.4 (Computed finite interface)**.** *The preceding simultaneous rule in ambient 58 pays every triangle at least 2 and every quadrilateral at least 1. Every actual rank 58 finite forest has charge at most 377 except precisely the following eight formal forests: one B_{13} or D_{13} together with any multiset of three A_{15} or D_{15} components. Each of those eight has charge378. All ideal weights admit the envelope of the ideal-vertex discussion below.*

Computed interface. This is the preliminary finite proposition supplied by catalogue FV2. The proof first establishes the complete context cover and its domination of every actual vertex forest, then

verifies all residual transition inequalities just described. Reconstructing each terminal forest gives its full contextual cost, including every cross-component payment. The only formal forests above 377 have cost $78 + 3 \cdot 100 = 378$ and are exactly the eight in the statement. Their necessary-domain status suffices; no realization is asserted. The local geometric coverage and triangle proofs are catalogue FV1. □

Number A_r by its path $0, \dots, r-1$ and B_r by the same path with its last bond doubled. Number D_r by the spine $0, \dots, r-3$ with leaves $r-2, r-1$ at $r-3$; its long endpoint is 0. These conventions describe actual whole components. The intrinsic weight tables are invariant under their automorphisms.

A separate frozen B_{13} endpoint theorem removes pair $(0, 12)$, with no new coupon. Its complete selected domain has no affected image, but that absence alone is insufficient: two older affine-B_{13} singleton routes really consumed the endpoint unit. Their whole cores instead give distinct already-unit A_1/D_{12} short-leaf contributions. The other endpoint-like pair $(1, 12)$ needed by the current D_{12} theorem is retained. Thus the four B_{13} forests in Proposition 3.4 drop to 377.

3.3 The dimension-specific final local rule

Theorem 3.5 (Three-rank 15 transfer)**.** *In ambient dimension 58, make these changes to the simultaneous rule:*

1. *In a whole finite D_{13}, give pairs $(0, 11), (0, 12)$ weight $1/2$ instead of 1 only if its actual corner has three distinct whole unmarked A_{15} or D_{15} components.*

2. *Add $1/2$ to whole D_{28} pair $(0, 12)$ only if its corner has two such whole unmarked rank 15 components.*

3. *Add $1/2$ between an endpoint of whole A_{12} and the long endpoint of whole D_{16} under the same two-component presence guard.*

All ideal weights and other finite weights remain. Every triangle still pays at least 2 and every quadrilateral at least 1.

Witnesses are not counted with multiplicity. The negative guard uses $13 + 3 \cdot 15 = 58$ ranks, so the corner has exactly those four components. The two positive guards similarly force exactly $D_{28} + 2 \cdot 15$ or $A_{12} + D_{16} + 2 \cdot 15$. The theorem is not asserted in larger ambient dimensions.

3.3.1 Selected cases and arbitrary additional supporting walls

The fresh complete target domain contains 801 patterns,862 D_{13} maps, 33 affected contexts and 76 affected pair maps. The physical cores have norm-two roots; full Lorentz, white and all four edge signatures and actual corner classifications are reconstructed exactly. The native-domain completeness is inherited from the universal reduction, rather than claimed from these records alone.

There are five proper completions. Their missing fork leaf is genuinely new: old corner roots outside the displayed A_{12} are orthogonal to it. Its forced unit contact has inverse projection square 6,12,30 or 102, exceeding 2. Additional same-end contacts cannot lower the bound. Mixed-end contacts give an independently paid short path. Thus these completions are impossible.

Four radical cores, numbered 1985,2378,2575,2581, seal the entire displayed core. Indeed the whole selected D_{13} forbids contacts on its nodes; every remaining contact position has a nonzero radical coefficient. A new white must pair to zero with the radical. Nonempty contacts therefore

meet both signs and give a retained adjacent-black shortcut of length at most 8. Otherwise all contacts vanish and the unmarked A_4 is whole. It cannot coexist with the three rank 15 witnesses.

Twelve contexts similarly have a small displayed block with no permitted new contact of projection at most 2. It is whole: any indirect extension would have a first new root contacting it. The rank-capacity guard again excludes it.

In the final twelve contexts the old small factor K has rank $s = 1, 2, 3$. Its unique permitted contact position p has core-inverse square $r = 4/3, 3/2, 2$, respectively. The native gap excludes magnitudes greater than 1; positive-end inverse bounds exclude two same-end contacts. Four cases also permit an opposite contact position q, with $J_{pp} = J_{qq} = 4/3$ and $J_{pq} = -4/3$. Two new roots using p, q have residual squares $2/3$, so their full product is at most $-4/3 + 2/3 = -2/3$. Their forced native bond supplies the retained short path. Thus, without that escape, all new whites in a growing K component contact the old core only at p; some may be orthogonal to it.

Let Y contain all new whites in this whole component, including every indirect tail, and let M be its rank. Every component of Y contacts K. With contact vector u, the full Lorentz condition gives the positive semidefinite residual Schur matrix

$$G_Y - ruu^t \succeq 0, \qquad u^t G_Y^{-1} u \le 1/r.$$

The inverse bound in Lemma 3.3 implies $rM/(M+1) \le 1$, hence $M \le 3, 2, 1$ respectively. In all cases the whole K component has rank at most 4, contradicting the guard. This controls arbitrary supporting whites, not just immediate neighboring roots.

3.3.2 The affine-D_{12} case

The old opposite-fork-leaf affine-D_{12} theorem genuinely used the newly lowered whole finite D_{13} unit. It must be repaired separately. Number its affine factor F of order 13 with spine 0 through 8, short leaves 9,10 at 0 and 11,12 at 8. Choose ideal marks $a = 9, b = 11$ and opposite black roots c, d. Let Z be the elliptic spectator forest, orthogonal to F, and x, y the nonnegative contacts of c, d with Z.

An adjacent-black bond or a shared spectator neighbor is an independently paid short path. Otherwise x, y have disjoint supports. The complete all-label end-profile reduction gives each end cost at most 8; a nonextreme end has cost at most 4. Write g_c, g_d for the end costs, $h, k > 0$ for the corresponding height scales, and S for the Schur matrix in this end-profile reduction. The exact identity supplied by the reduction is

$$\frac{\det S}{h^2 k^2} = 1 - \frac{g_c + g_d}{12} + \frac{1}{12}(x/h - y/k)^t G_Z^{-1}(x/h - y/k).$$

Its determinant vanishes at the actual four-black star.

If an end is nonextreme, equality forces costs 8 and 4 and $x = y = 0$. A quiet root attached to the selected old leaf gives the retained whole A_1/D_{12} short-leaf unit. If it attaches to the mate, it gives whole D_{13}. Without the new guard its unit is unchanged. With the guard, Z consists of exactly three orthogonal whole rank 15 factors. The three nonextreme equality profiles give, at a distinct corner, respectively a retained A_1/D_{12} unit, an independent whole affine B_{13} priority, or a second guarded D_{13} half. These exhaust 24 physical cases. The current B_{13} endpoint repair uses its D_{12} cross unit, not a newly financed D_{13} route; there is no circularity.

When both ends are extreme, $h = k = 1/2$, and the determinant becomes

$$(x - y)^t G_Z^{-1}(x - y) = 1. \tag{3.3}$$

The old two-nonempty-support theorem gives retained short paths or small sealed alternatives. The changed D_{13} output belongs to the arbitrary one-sided branch, say $y = 0$. The quiet corner excludes c and retains Z orthogonally. If its D_{13} guard holds,Z is exactly three whole A_{15}/D_{15} factors.

Lemma 3.6. *Under the strict edge conditions and disjoint-support hypothesis, energy 1 on three whole A_{15}/D_{15} factors forces a single unit contact at the long endpoint of one D_{15}, by one black root. The other two factors have zero contact with both black roots.*

Proof. For A_{15} the inverse diagonal entries below 2 occur only at nodes 0,1,13,14. They are 15/16, 7/4, 7/4, and 15/16, respectively. Two native contacts give energy at least 2 and violate strict edge positivity. The complete profiles are simple contacts at these four nodes or doubled contacts at 0,14. The next native squared magnitude $(3 + \sqrt{5})/2$ is already excluded. For D_{15} only its long endpoint has inverse diagonal below 2, equal to 1, and only its simple contact fits.

Every allowed nonzero component energy is at least 15/16; no A_{15} profile has energy 1. The finite exact inverse table also checks every disjoint two-sided profile difference: all are at least 15/16 and none is 1. Equal-node contacts are forbidden by disjoint support. Two active components would have total at least $15/8 > 1$. Equation (3.3) therefore forces precisely the stated D_{15} contact. □

The two inactive components are orthogonal to the entire star and the active component. They remain whole, without indirect bridges. If c attaches to selected a, the opposite corner is whole A_{12} at an endpoint and whole D_{16} at its long endpoint. If it attaches to a's mate, that corner is whole D_{28} at $(0, 12)$. Both physical 30-root models have signature $(28, 1, 1)$, positive white rank 26, positive edge rank 27 and semidefinite corners. In either case the new half, guarded by the two inactive rank 15 factors, supplements the quiet D_{13} half at a different actual corner. This proves the required total 1.

3.3.3 Retention of other payments

The preceding repairs must be used within the single simultaneous rule. In particular every earlier route must keep its chosen positive output, or receive a proved replacement. Catalogue FV1 supplies the complete retention proof. Its low-order and protected-path outputs use distances strictly below the changed D_{13} distance eleven. Other D_{13} roles, such as its fork-center and the older affine-D_{11} outputs, remain unchanged. The affine-D_{12} case just treated is the separate case where an old route really used the changed unit.

Triangles require the complete actual-face argument, not three pairwise checks in one abstract diagram. The affine triangle-exclusion lemma in Addendum FV removes the three reduced small affine patterns. Low cores give two distinct corner paths of length at most five. High cores give paths of length at most seven or the four protected exceptional E_8 outputs. The nearby whole-leaf argument protects the three exceptional length-six outputs; length-seven and ideal contributions are unchanged. The distance-eleven transfer therefore retains their chosen payments. Together with the complete core and extension classification in catalogue FV1, this proves the local transfer theorem.

3.4 The finite maximum after the transfer

Proposition 3.7. *Every actual finite rank 58 forest under the final simultaneous rule has charge at most 377. The exact formal maximum is 377.*

Proof. The B_{13} theorem lowers four forests in Proposition 3.4 from 378 to 377. The new D_{13} guard lowers the other four by 1. Every other forest was already at most 377, except that the two new positive coupon families must be checked.

Rank capacity makes those families exact. They have respectively

$$170 + 2(100) + \tfrac{1}{2} = 370.5, \qquad 66 + 108.5 + 2(100) + 1 = 375.5.$$

The second total counts both A_{12} endpoint coupons. The previous A_{12}/D_{16} cross cost is zero. Each family has three possible multisets of two A_{15}/D_{15} factors, so all six actual forests are covered. No additional context witness can fit. Outside them the change is pointwise nonincreasing. Thus the bound 377 holds everywhere, and the eight changed forests attain it formally. A geometric realization of an attainer is neither required nor asserted. □

3.5 Ideal vertices and positive surplus

For affine ranks r_i summing to 57 set

$$H(t) = \prod_i (1 + t + \cdots + t^{r_i}) = \sum_{j=0}^{57} h_j t^j, \qquad V = \prod_i (r_i + 1),$$
$$G = \sum_j h_j \min(j, 57 - j), \qquad \kappa = \frac{5G - 57V}{2}.$$

The geometric all-family envelope is unchanged by every final finite transfer. Its complete numerical table is

$$c(r) = r/2 \quad (1 \le r \le 8), \qquad c(r) = \frac{5r + 56}{r + 1} \quad (24 \le r \le 57),$$

together with the following entries:

r	9	10	11	12	13	14	15	16
$c(r)$	71/16	53/11	31/6	11/2	40/7	367/60	411/64	227/34
r	17	18	19	20	21	22	23	
$c(r)$	119/18	128/19	137/20	146/21	155/22	163/23	85/12	

The finite arithmetic below assumes the proved factor-charge envelope (3.1); polynomial evaluation does not establish its geometric meaning.

3.5.1 The terminal domain and its exact output

Let $\mathcal{P}(n, \ell)$ be the list of nondecreasing partitions of n with parts at least ℓ. Generate it by

$$\mathcal{P}(0, \ell) = \{()\}, \qquad \mathcal{P}(n, \ell) = \bigcup_{q=\ell}^{n} \{(q, p) : p \in \mathcal{P}(n - q, q)\} \quad (n > 0).$$

The union is empty if $\ell > n$. The terminal cusp domain is

$$\mathcal{T}_{57} = \bigcup_{\ell=0}^{7} \{(\ell, p) : p \in \mathcal{P}(57 - \ell, 8)\}, \tag{3.4}$$

where a zero part is omitted. For each profile compute its polynomial by repeated convolution: start with $h_0^{(0)} = 1$, and when adjoining rank r set

$$h_j^{(k+1)} = \sum_{a=0}^{r} h_{j-a}^{(k)},$$

with out-of-range entries zero. Then compute V, G, κ from their displayed definitions and set $m = \kappa/V - \sum_i c(r_i)$.

Proposition 3.8 (Complete terminal calculation)**.** *The partition and convolution algorithms terminate, list every profile in* (3.4) *exactly once, and compute its margin by exact rational arithmetic. Their recorded complete output is 2552 profiles, with*

$$\min_{\mathcal{T}_{57}} m = \frac{87}{10240} > 0, \qquad \min_{\mathcal{T}_{57}} Vm = 36.$$

Algorithm correctness and computed output. Every recursive call lowers n by a positive integer, and every loop is finite. The unique first part of a sorted partition and its unique tail prove completeness and absence of duplicates by induction on n. Its exceptional part below 8, if present, determines the outer branch ℓ uniquely. Polynomial convolution is exactly multiplication by $1 + t + \cdots + t^r$, so induction on the factors proves every coefficient. Degrees never exceed 57. All remaining operations are finite integer or rational sums and products.

The stated cardinality and minima are the computed proposition: the complete saved certificate records every rank tuple and coefficient list, and the independent endpoint checker reconstructs the domain and every convolution before comparing them. It evaluates all 2552 margins, rather than checking only degree, symmetry or a sample of rows. The two minima are separate assertions and need not have the same attaining profile. The implementation and records are identified in the reproducibility section. □

Lemma 3.9 (Concentrating two simplex ranks)**.** *Let $1 \le a \le b$ be integers, and let U_r be uniform on $\{0, \dots, r\}$. With independent summands, $U_{a-1} + U_{b+1}$ dominates $U_a + U_b$ in convex order. Adding any common independent summand preserves this order.*

Proof. Both sums have mean $(a+b)/2$ and are symmetric about that mean. For an integer threshold t, let Δ_t be the difference of their expectations of $(X - t)_+$. Equal means allow us to use $(t - X)_+$ instead, and symmetry gives $\Delta_t = \Delta_{a+b-t}$. It suffices to take $0 \le t \le (a+b)/2$. Put $D = (a+1)(b+1)$ and $D' = a(b+2)$. For $0 \le t \le a$, direct summation of the increasing side of the two convolution triangles gives

$$\Delta_t = \binom{t+2}{3}\left(\frac{1}{D'} - \frac{1}{D}\right) \ge 0, \qquad D - D' = b - a + 1 > 0.$$

For $a \le t \le (a+b)/2 \le b$, summing over the first uniform instead gives

$$\Delta_t = \frac{t(a+b-t) - (a-1)(a+2b+2)/3}{2(b+1)(b+2)}.$$

The numerator is increasing up to $(a+b)/2$. At $t = a$ it is $(a+2)(b-a+1)/3 > 0$. Outside the common support the difference is zero; between integer thresholds it is linear. Thus all stop-loss differences are nonnegative. On the integer support the values of a convex function are an affine function plus a nonnegative sum of these stop-loss functions, proving convex order. Conditioning on a common independent summand proves the last assertion. □

Lemma 3.10 (Concentration below rank eight)**.** *Every actual ideal vertex has $\kappa - C_{\mathrm{act}} \ge 36$.*

Proof. For $1 \le a \le b \le 7$, replacing (a, b) by $(a-1, b+1)$ preserves normalized envelope cost, since all these ranks stay at most eight. Lemma 3.9, including its independent-summand assertion, gives the required convex order. Since

$$\frac{G}{V} = \frac{57 - \mathbb{E}|2S - 57|}{2},$$

the transfer does not increase normalized curvature or normalized margin. Its product decreases from $(a+1)(b+1)$ to $a(b+2)$. Iteration terminates at ranks at least 8 and at most one positive

rank below 8. Indeed the integer potential $\sum_i r_i^2$ increases by $2(b-a+1) \geq 2$ at every step and is bounded above by 57^2. Its initial value is at least 57, so at most 1596 steps occur. Rank zero is omitted. No transfer passes through the changed rank 9 envelope. The stopping domain is exactly (3.4).

Proposition 3.8 gives positive normalized terminal margin and absolute envelope surplus at least 36. If the original and terminal quantities are (V, m) and (V', m'), then $V \geq V'$ and $m \geq m' > 0$. Thus $Vm \geq V'm' \geq 36$. This multiplication uses positivity. Finally (3.1) transfers the same lower bound to actual cusp charge. □

3.6 Shape curvature and the contradiction in dimension 58

Lemma 3.11. *For a proposed ambient 58 polytope,*

$$\mathcal{D}_5 \geq 377 v_f + \sum_{v \text{ ideal}} \kappa(v) + 4205.$$

Proof. Choose a generic linear function on the bounded Klein closure of the polytope. Let E_v be the number of edges of its simple vertex figure and E_v^0 the number not cut by the induced level section. If I counts all polygon vertices with multiplicity, the published incidence identities give $I = \sum_v E_v$ and $2f_2 = \sum_v E_v^0$ [2, Theorem 9]. Hence $\mathcal{D}_5 = \sum_v (5E_v^0/2 - E_v)$.

Orient each vertex figure by its induced generic linear function. Assign an uncut edge below the section to its upper endpoint and one above the section to its lower endpoint. A vertex of index j receives j or $57 - j$ contributions, at least their minimum. The index distribution is its h-vector [2, Theorems 2 and 6], so $E_v^0 \geq G_v$. At the distinct global minimum and maximum the whole vertex figure is uncut, and symmetry gives

$$E_v - G_v = \sum_{j > 57/2} h_j(2j - 57) \geq \sum_{j=29}^{57} (2j - 57) = 841.$$

Every coefficient of a simplex or product-of-simplices h-polynomial is at least 1. Adding the two extrema and multiplying by $5/2$ gives 4205, counted once. For a finite vertex $h_j = 1$, so

$$G_0 = \sum_{j=0}^{57} \min(j, 57 - j) = 812, \qquad \frac{5G_0 - 57 \cdot 58}{2} = 377.$$

The remaining ideal terms are precisely κ. □

For faces with at least five sides,$5 - s(F) \leq 0$. The nonnegative local rule and Proposition 3.7 therefore give

$$\mathcal{D}_5 \leq 377 v_f + \sum C_{\text{act}}.$$

Subtract Lemma 3.11 and use Lemma 3.10:

$$0 \geq 4205 + \sum (\kappa - C_{\text{act}}) \geq 4205 + 36 v_i > 0.$$

This also excludes the all-finite case. Thus dimension 58 is impossible.

It remains to exclude dimensions at least 59. We now give the separate arguments used for that purpose.

3.7 The bounded higher-dimensional range

Put $d = n - 1$ and

$$\beta_n = \begin{cases} n(n-6)/8 & n \text{ even}, \\ (n-1)(n-5)/8 & n \text{ odd}. \end{cases}$$

The incidence argument of Lemma 3.11 applies at every rank:

$$\mathcal{D}_5 \geq \beta_n v_f + \sum_{v \text{ ideal}} \kappa(v) + \Delta_n, \qquad \Delta_n > 0.$$

The weaker bound $\Delta_n \geq 5(n-1)$ suffices here. For a simultaneous rule with finite bound U_n and ideal charge $C(v)$, it is enough that $U_n \leq \beta_n$ and $\kappa(v) - C(v) > 0$ at every cusp. Subtraction as in Lemma 3.1 then gives a contradiction, even if there are no ideal vertices.

For a cusp profile $\boldsymbol{r} = (r_1, \ldots, r_s)$ of total rank d, define

$$H_{\boldsymbol{r}}(t) = \prod_i (1 + \cdots + t^{r_i}) = \sum_j h_j t^j, \quad V_{\boldsymbol{r}} = \prod_i (r_i + 1), \quad G_{\boldsymbol{r}} = \sum_j h_j \min(j, d - j).$$

The normalized surplus associated with an affine-factor envelope c is

$$m_c(\boldsymbol{r}) = \frac{5G_{\boldsymbol{r}} - dV_{\boldsymbol{r}}}{2V_{\boldsymbol{r}}} - \sum_i c(r_i). \tag{3.5}$$

The factor-fiber theorem gives $C(v) \leq V_{\boldsymbol{r}} \sum_i c(r_i)$. This multiplicity is a geometric statement; a graph pair count alone would not establish it.

Lemma 3.9 permits the move $(a, b) \mapsto (a - 1, b + 1)$, $1 \leq a \leq b$, whenever $c(a-1) + c(b+1) \geq c(a) + c(b)$. The normalized curvature, and hence (3.5), does not increase. The product of the two link vertex counts decreases, since $(a+1)(b+1) - a(b+2) = b - a + 1 > 0$. Every move raises the integer sum of squared ranks by $2(b - a + 1)$, while that sum is at most d^2. Thus concentration halts. A positive terminal margin extends to every original cusp. If an absolute surplus is used, its normalized positivity is checked before multiplying by the larger original vertex count.

The sorted-partition recursion and exact convolution described above apply with total rank d. The permitted concentration moves determine the terminal multiplicity restrictions. For example, for $c(r) = \min(r/2, L/2)$, terminal profiles have arbitrary copies of L, at most one part below L and at most one part above L. A low-only reduction instead permits arbitrary parts at least L. The full domains and each exceptional rank condition are part of catalogue FV4's finite propositions. Each resulting minimum is over the entire stated domain. Finite-vertex costs use the same typed-forest recurrence with all context-sensitive data fixed. Domain completeness and all transition inequalities are proved before the numerical comparison is used.

3.7.1 The elementary families in dimensions 98–163

A feature is a node of degree at least three or an endpoint of a nonsimple bond, including a parallel bond. Distances are graph distances. A condition on a whole component always concerns the whole actual endpoint diagram.

For $98 \leq n \leq 108$, finite weights are one at distance at most 9 or in a whole component of order at most 17. Their total is at most $9n$. At a cusp, pay one at distance at most 8, in a whole component of order at most 17, or at distance 9 if a mark is within 14 steps of a feature. These are

alternative conditions for one unit, not separate charges. For an affine component of order $m \geq 18$, the exact counts are

A	$8m$
C	$\min(9m-45, 8m-4)$
B	$\min(9m-44, 8m-2)$
D	$\min(9m-43, 8m)$.

For example, a C spine contributes $8m-36$ short pairs; the extra distance-nine starts occupy the first 16 and last 16 available positions, giving $\min(m-9, 32)$. In B the spine windows have lengths 15 and 16 and the leaves add two pairs. In D both windows have length 15 and the leaves add four pairs. The union of the first a and last b positions in a list of length h has size $\min(h, a+b)$; this also covers overlapping windows. Thus these are all-rank counts. Orders at most 17 cost at most $\binom{m}{2} \leq 8m$, so $c(r) = \min(8, r/2)$.

For $k = 9, \ldots, 13$, use unit weights at distance at most k or in whole components of order at most 17, at both finite and ideal vertices. The finite bound is kn and $c(r) = \min(k, r/2)$. For $k = 14$ the distance condition alone suffices. The complete local replacement theorem for these distance families is the core-routing theorem of Addendum FV, "Geometric coverage and exact local certificates," with the corresponding retained short-path threshold. It includes overlapping cores and arbitrary additional supporting whites. The affine graph count does not prove that local theorem.

The terminal algorithm above, with $L = 2k$, gives the following exact positive minima over the indicated dimensions and their entire terminal domains:

n	U_n	$c(r)$	$\min m_c$
98 : 108	$9n$	$\min(8, r/2)$	112101/1419857
109 : 116	$9n$	$\min(9, r/2)$	45873/2476099
117 : 126	$10n$	$\min(10, r/2)$	145234/4084101
127 : 136	$11n$	$\min(11, r/2)$	40603/559682
137 : 146	$12n$	$\min(12, r/2)$	172096/1953125
147 : 156	$13n$	$\min(13, r/2)$	830407/9565938
157 : 163	$14n$	$\min(14, r/2)$	1476616/20511149

In every row $\beta_n > U_n$. This proves the exclusion of 98–163.

3.7.2 The remaining range 59–97

The smaller dimensions use refined marked roles. Catalogue FV4 supplies three complete families for 59–65, 66–97 and 98–163, with the last family described above. For the first two families the finite forest maximum U_n, the affine envelope c_n and its complete terminal profile domain are fixed separately at each stated range. They satisfy $U_n \leq \beta_n$ and $m_{c_n} > 0$ throughout their respective domains. This is a computed conjunction over the full named domains, not a numerical extrapolation from a boundary dimension.

The complete local rule and its replacement proofs are in Addendum FV, "Labelled rules for dimensions 66–97" and "The complete ambient 59–65 layer table." They include the finite and affine choices, whole-component guards and all compensating payments. The same concentration and product-multiplicity argument just proved carries positive terminal margins back to every actual cusp. The common comparison therefore excludes each dimension $59 \leq n \leq 163$.

3.8 The unbounded tail: actual framing walls and two budgets

The range $n \geq 164$ requires a proof at all ranks, not a finite list of partitions. We use Nikulin's narrow-facet theorem [11, pp. 654–655]; its unrestricted statement is also given in [12, Section 4, p. 10]. A finite-volume chamber has a facet F whose actual framing hyperplanes have pairwise distance strictly less than $\operatorname{arccosh} 7$. With outward normals of square 2 this implies

$$-(x, y) < 14 \tag{3.6}$$

for any two framing walls. This is the external narrowness input. We prove the incidence, endpoint and scale statements needed to apply it; they are not additional hypotheses on the cusp.

3.8.1 Outgoing edges give actual framing walls

Suppose first that F contains a cusp v. Its horospherical link is a product of affine Coxeter simplices. Write their ranks r_a, with $\sum_a r_a = N = n - 1$, and normalize every old root to square 2. In each factor choose two omitted nodes in advance. In the factor containing the normal α_s of F, both choices avoid s. The two choices need not be distinct. If that factor has affine rank one, use its other node twice; its modified target $2(r_a - 1)/9$ is zero. Deleting one node per factor gives a positive-definite support S of N roots and a vertex of the horospherical link. It therefore defines an actual outgoing edge in F. Finite volume gives a finite or ideal second endpoint.

At a finite endpoint choose any new wall x terminating that edge. It and α_s belong to a spherical diagram, so their rank-two Gram is positive definite. At an ideal endpoint let T_b be its affine factors. The positive-definite support S contains at most $\operatorname{rank} T_b$ nodes from each. The total rank N forces equality in every factor: exactly one node of each T_b is missing. If the factor containing α_s has rank at least 2, choose its missing node as x; every pair in such a factor has finite angle. If it is affine A_1, choose the missing node of another factor, which is orthogonal to α_s. Another factor exists in the range at issue. Thus again the pair is positive definite.

To see that this is an actual ridge, start with a strict interior timelike point z. Project it off the span of x, α_s. The projection correction is a nonnegative combination of those normals, because their Gram is a positive-definite matrix with nonpositive off-diagonal entries and nonnegative inverse. Every other wall inequality stays strict, and the projected square is even more negative. It gives a point of precisely their common face. Hence x is in the framing set controlled by (3.6). Apply the same construction to the second omission tuple and call its terminating wall y.

Neither x nor y is an old deleted cusp wall: that wall is strict along the outgoing edge away from v. Their old contact rows $p_j = -(x, \alpha_j)$ and $q_j = -(y, \alpha_j)$ are nonnegative, and every nonzero entry is at least 1. For each affine factor let $c_a = \sum_j \delta_{aj}\alpha_j$, with $\delta_{aj} > 0$ normalized by $\min_j \delta_{aj} = 1$. These are nonzero future null vectors, since their pairing with an interior point is negative. They are all positive multiples $k_a c$ of one cusp vector c. Therefore

$$\sigma_a = \delta_a \cdot p = -(x, c_a), \quad \tau_a = \delta_a \cdot q = -(y, c_a), \quad q_0 = \sigma_a/\tau_a$$

have a common positive ratio. Indeed, if a row height vanished, all its nonnegative contacts with that factor would vanish; proportionality of the c_a would force every factor height and every old contact to vanish. A square-two normal would then lie in the null orthogonal complement of the full cusp span, impossible.

For each endpoint the principal Gram $S+x$ or $S+y$ is positive semidefinite. Its Schur complement gives the separate budgets

$$\sum_a C_{0a} \leq 2, \qquad \sum_a C_{1a} \leq 2, \qquad C_{\ell a} = p^t_{\ell,S_a} G(S_a)^{-1} p_{\ell,S_a}. \tag{3.7}$$

The inverse is entrywise nonnegative. No bound on a deleted-node contact or on its Coxeter label has been used.

3.8.2 The common-height Lorentz identity

Choose a null partner d_0 with $(c, d_0) = -1$, and write $x = ac + bd_0 + u$, $y = a'c + b'd_0 + u'$ in the orthogonal Euclidean quotient. Then $b = -(x, c) > 0$, $b' = -(y, c) > 0$. Since $(x/b - y/b') \perp c$ and $x^2 = y^2 = 2$,

$$\|u/b - u'/b'\|^2 = 2/b^2 + 2/(b')^2 - 2(x, y)/(bb').$$

The quotient is the orthogonal sum of the affine factor spaces. In factor a put $E_a = \|u_a/\sigma_a - u'_a/\tau_a\|^2$. Since $\sigma_a = k_a b$ and $\tau_a = k_a b'$, the k_a^2 cancels and the preceding identity becomes

$$-(x, y) = \frac{1}{2} \sum_a \sigma_a \tau_a E_a - q_0 - q_0^{-1}. \tag{3.8}$$

This proves the scale cancellation for a product cusp; no commensurability or chosen horosphere is required. Let $M = \max(q_0, q_0^{-1})$ and $W_a = \min(\sigma_a, \tau_a)^2 E_a$. Then $\sigma_a \tau_a E_a = M W_a$ and $q_0 + q_0^{-1} \le 2M$.

The prescribed omission choices have targets

$$T_a = 2r_a/9 \quad \text{ordinarily,} \qquad T_a = 2(r_a - 1)/9 \quad \text{in the sole avoid-node factor.}$$

Thus $\sum T_a = (2N - 2)/9$. We next prove

$$\sum_a W_a \ge \begin{cases} (2N-2)/9 - 4 & 1 \le M \le 21/20, \\ (2N-2)/9 - 16/3 & M \ge 21/20. \end{cases} \tag{3.9}$$

It follows from(3.8) that $-(x, y) \ge M(N - 37)/9$ in the first range and $-(x, y) \ge M(N - 43)/9$ in the second. At $N = 163 + t$ these are at least $14 + t/9$ and $14 + 7t/60$, respectively. They contradict (3.6) for every $t \ge 0$.

3.8.3 Point potentials and the original endpoint budgets

For r Euclidean coordinates put $s_k = e_1 + \cdots + e_k$. The classical point potentials and normalized kernel squares are

	v_k	(δ_k^2)
B	$v_0 = 0,\ v_1 = e_1,\ v_k = s_k/2\ (k \ge 2)$	$(1, 1, 4, \ldots, 4, 2)$
C	$v_k = s_k,$ metric one half the usual metric	$(1, 2, \ldots, 2, 1)$
D	$v_0 = 0,\ v_1 = e_1,\ v_k = s_k/2\ (2 \le k \le r-2)$	
	$v_{r-1} = (s_{r-1} - e_r)/2,\ v_r = s_r/2$	$(1, 1, 4, \ldots, 4, 1, 1).$

Write $R(i, j) = \|v_i - v_j\|^2$. For an A cycle of order m use $\delta = 1$ and $R(i, j) = h(m - h)/m$, $h = (i - j) \bmod m$. Ordinary omissions are $(0, \lfloor m/2 \rfloor)$ in A and $(0, r)$ in B, C, D. Modified B, C use $(r - 1, 0)$ and $(1, r)$. Rotations and fork automorphisms avoid a prescribed node in A, D without weakening the ordinary geometric estimate.

Exceptional factors use the ordinary and avoid-node omission choices in Addendum FV's "Point potentials and the complete local domain." That section gives their grounded Gram matrices, positive affine kernel coefficients and the exact rational resistance formula. These fixed choices, rather than arbitrary omissions, are used in the exceptional classification and packing below.

Lemma 3.12 (Full-budget classical unit charge)**.** *For the prescribed ordinary or avoid-node omissions of a classical affine factor, let $C_0, C_1 \le 2$ be its actual retained costs, and let $W = \min(\sigma, \tau)^2 E$. If the common height ratio is at most $B = 21/20$, then*

$$W + C_0 + C_1 \ge T,$$

where $T = 2r/9$ ordinarily and $T = 2(r-1)/9$ in an avoid-node factor. This includes arbitrary real contact magnitudes with every nonzero contact at least one, and both possibilities for each omitted-node contact.

Reduction to the complete local certificates. For omissions i, j, write $R(i,j) = \|v_i - v_j\|^2$ and let w_k, z_l be the nonnegative contact weights normalized by their affine heights. They sum to one on each full row, including an omitted-node contact when present. Write their point means as $v_i + a, v_j + b$. Polarization gives

$$\begin{aligned} E = R(i,j) &- \sum_k w_k \{R(i,j) + R(i,k) - R(k,j)\} \\ &- \sum_l z_l \{R(i,j) + R(j,l) - R(l,i)\} + \|a - b\|^2. \end{aligned} \tag{3.10}$$

The omitted-node displacement is zero. Every row height is at least one, so $W \ge E$. If $c_{ik} = \delta_k^2 R(i,k)$, nonnegativity of the retained inverse and the contact gap yield

$$C_0 \ge \sum_{k \text{ retained, contacted}} c_{ik}, \qquad C_1 \ge \sum_{l \text{ retained, contacted}} c_{jl}.$$

Thus unit diagonal costs bound the weighted linear losses without an upper bound on the contact magnitudes. Substitution of the displayed B, C, D potentials, with the prescribed ordinary and avoid-node choices, gives $W + C_0 + C_1 \ge T$. The complete contact partition and its coefficient-one inequalities are the "Full-budget classical unit charge" lemma of Addendum FV. They use the original separate budgets $C_0, C_1 \le 2$, not the residual budget $3/2$.

For the A cycle of order m, the same resistance formula reduces all sufficiently large orders to polynomial inequalities. Put $\varepsilon = m \bmod 2$. Cost at most two excludes cyclic distance at least three for $m \ge 18$, because $3(m-3)/m > 2$ and resistance increases up to the antipode. A remaining contact at distance $d = 1, 2$ has linear-loss to cost ratio at most $L_m = (m + \varepsilon)/(m-2)$. Therefore

$$W + C_0 + C_1 - T \ge R(0, \lfloor m/2 \rfloor) - T - \frac{4(2+\varepsilon)}{m-2}.$$

For $m = 18 + 2u$ the right side is

$$\frac{128 + 84u + 4u^2}{36(m-2)} > 0;$$

for $m = 19 + 2u$ it is

$$\frac{360 + 1708u + 252u^2 + 8u^3}{36m(m-2)} > 0.$$

These identities hold for every $u \ge 0$. The remaining cycle orders $2 \le m \le 17$ are the complete finite convex-quadratic domain in the same addendum. Its proof includes both omitted-node states and every allowed retained support; a ratio bound on the common heights is kept throughout the finite minimization. Rotation avoids a prescribed node without weakening the ordinary target for $m \ge 3$. At $m = 2$ the repeated allowed omission has modified target zero. This completes the reduction to the finite local certificates specified in catalogue FV4. □

The complete exceptional support classification in Addendum FV identifies all factors where $T - W > C_0 + C_1$. It is applied at the original budgets $C_0, C_1 \le 2$. Its conclusion is that such a factor is an ordinary E_8 with cost floors $(8/9, 2/3)$ and target $16/9$, or an ordinary E_7/modified E_8 with floors $(1/2, 1)$ or $(1, 1/2)$ and target $14/9$. The full-budget classical lemma is required before this exceptional classification can be used; the residual bounds alone do not identify the first failing factor.

After one failing factor is removed, each remaining ordered budget is at most $3/2$. The complete finite domain at those budgets is obtained from grounded inverse matrices, every permitted contact support and both omitted-node states. Its nonnegative retained inverses bound each support by its unit-contact cost. Normalizing by the common positive heights puts the corresponding point potentials in two finite convex hulls. Exact convex-quadratic certificates bound their distance, hence their loss $T - W$, for arbitrary contact magnitudes. The finitely many exceptional and small classical matrices, and the all-rank coordinate proofs for the remaining classical ranks, are in Addendum FV, "Point potentials and the complete local domain." Their joint result is the following ordered domination table. This is the finite interface used in the next argument; the continuous local proof is not replaced by the table's row count.

3.8.4 Exceptional factors and ordered residual packing

Each positive-loss domain is dominated by a record with no larger ordered cost floors, no smaller loss bound, and the same modified flag. The complete 28-record table is:

Type	(C_0, C_1) floors	loss bound
ordinary A_1	$(1/2, 0)$	$2/9$
ordinary A_2	$(2/3, 0)$	$4/9$
ordinary A_4	$(4/5, 4/5)$	$8/9$
ordinary $B_r,\ 2 \le r \le 5$	$(0, r/4)$	$2r/9$
ordinary $D_r,\ r = 4, 5$	$(r/4, 0)$	$2r/9$
ordinary E_6	$(0, 4/3), (4/3, 0)$	$4/3$ each
ordinary E_7	$(1/2, 1), (1, 1/2)$	$14/9$ each
ordinary E_8	$(8/9, 2/3)$	$16/9$
modified $B_r,\ 3 \le r \le 7$	$((r-1)/4, 0)$	$2(r-1)/9$
modified E_6	$(0, 4/3)$	$10/9$
modified E_7	$(0, 3/2), (2/3, 5/6), (5/6, 2/3)$	$4/3$ each
modified E_8	$(1, 1/2), (8/9, 2/3)$	$14/9$ each
modified F_4	$(0, 1), (1/2, 3/4)$	$2/3$ each
modified G_2	$(0, 1/2)$	$2/9$

Catalogue FV4 supplies a dominating record for every positive-loss domain. No common geometric realization of the records is assumed.

For records with indices at least i, ordered remaining budgets p, q, and at most a modified records, the maximum residual loss is the maximum of 0 and

$$\ell_j + F(j, p - p_j, q - q_j, a - \epsilon_j), \qquad j \ge i,$$

over admissible records. Each has $p_j + q_j \ge 1/2$, so the recursion has length at most $\lfloor 2(p+q) \rfloor$. All costs have a common denominator, hence the state set is finite. Induction on remaining length proves completeness; checking every upper inequality and an equality predecessor proves the recorded maxima

$$F(0, 10/9, 4/3, 1) = 20/9, \qquad F(0, 1, 3/2, 1) = F(0, 3/2, 1, 1) = 22/9.$$

One additional complete exceptional check is needed. If $T - W > C_0 + C_1$, then it is an ordinary E_8 with cost floors $(8/9, 2/3)$ and target $16/9$, or an ordinary E_7/modified E_8 with floors $(1/2, 1)$ or $(1, 1/2)$ and target $14/9$. Its finite algorithm enumerates every grounded exceptional support below the trivial threshold $C_0 + C_1 < T$, including all inverse cross terms, and checks the resulting convex-quadratic bounds. There are 15 candidate exceptional supports. Two ordinary E_7 same-row supports at nodes 4,7 are excluded by exact convex certificates, in both omitted-node states, giving $W + 3/2 > T$. For the omissions $(3, 6)$ the four ordered full-support pairs are

$$((4, 7), (6)), \quad ((4, 7, 3), (6)), \quad ((3), (4, 7)), \quad ((3), (4, 7, 6)).$$

The coverage guard requires these four distinct keys, not merely four individually valid certificates. The remaining 13 have exactly the stated floors. This support check is separate from the residual domination table.

If no factor fails, the two budgets already bound total loss by 4. Otherwise choose any failing factor. An E_8 leaves budgets at most $(10/9, 4/3)$, and its own loss at most $16/9$ plus the residual $20/9$ is 4. In the other case the residual is at most $22/9$ and the own loss at most $14/9$, again 4. All additional failures are included in this residual calculation. Permitting one modified residual record even if the selected factor was modified only enlarges the upper bound. This proves the first line of (3.9).

3.8.5 The other height range

For the same omission choices the all-rank local inequality is

$$W + \frac{4}{3}(C_0 + C_1) \geq T.$$

To prove it, write the two normalized point potentials as $v_i + a, v_j + b$. With convex contact weights w_k, z_k,

$$E = R_{ij} - \sum_k w_k(R_{ij} + R_{ik} - R_{kj}) - \sum_k z_k(R_{ij} + R_{jk} - R_{ki}) + \|a - b\|^2.$$

A nonzero contact has magnitude at least 1, every convex weight is at most 1, and the nonnegative retained inverse makes its cost dominate the sum of unit diagonal costs. Also $W \geq E$, since both row heights are at least 1.

For B, C, D, use the coefficient-one argument in Lemma 3.12. That part of its proof uses only that both row heights are at least one, and hence is independent of the near-range ratio bound. Its complete contact table also covers modified B_2. Increasing the cost coefficient to $4/3$ preserves the inequality. The A and exceptional factors in this height range require the following separate arguments.

For A put $C = C_0 + C_1$ and $T = 2(m - 1)/9$. If $C \geq 3T/4 = (m - 1)/6$, nonnegativity of W proves the claim. Only $C < (m - 1)/6$ needs further work. For $3 \leq m \leq 6$ the smallest retained unit cost $(m - 1)/m$ already reaches this threshold, so no retained contact is possible. The literal A_1 case $m = 2$ is separate, and its modified target is zero. For $7 \leq m \leq 10$ a distance-two contact likewise reaches the threshold; only distance one remains. For $m \geq 10$ the original per-side budget two excludes every cyclic distance at least three, since $3(m - 3)/m > 2$ and resistance increases up to the antipode. For a remaining distance $d = 1, 2$, polarization leaves the margin

$$\frac{4}{3}\frac{d(m - d)}{m} - \frac{d(m + \varepsilon)}{m} = \frac{d(m - 4d - 3\varepsilon)}{3m} \geq 0.$$

The nontrivial domain starts at $m = 7$ for $d = 1$ and $m = 11$ for $d = 2$ in odd parity, where the respective margins are zero; the even cases are nonnegative too. Smaller odd ranks with a negative formal margin were already in the trivial domain. The finite exceptional omission cases enumerate every cheap retained support and both omitted-node states. Single contacts give a convex quadratic whose endpoint and stationary minima are checked exactly; the sole cheap two-contact support has energy $(1 + u^2 + v^2)/2$. These finite certificates use the explicit grounded matrices in Addendum FV, so have no missing high-rank or finite-angle branch.

Summing and using(3.7) proves the second line of (3.9). The certificate verifies the stated finite quadratic and symbolic polynomial identities. Their domains, the contact-gap argument and the coordinate identities explain why they imply the all-rank inequality.

3.8.6 A narrow facet without a cusp

For completeness we prove the compact-facet alternative rather than import a finer compact-chamber weighting. Let Q be a compact two-face. Its white normals W number $n - 2$, and every consecutive black pair with W is positive definite. A positive-definite principal block gives an actual common face by the interior-point projection already proved. A singular positive-semidefinite block has a positive null combination in an affine component. It is nonzero, future null and nonpositive against every other chamber wall, so gives an ideal point of its face.

For a triangle, a minimal indefinite star subdiagram contains all three black nodes: omitting any one puts it in a spherical corner. By the order-ten classification it has at most seven whites. In a spanning tree joining the three marks, deleting either of two outer marks leaves the other two joined. The resulting two paths avoid the third mark and have at most eight edges, so lie in two actual corner diagrams.

For a quadrilateral split opposite black pairs as U, V. The block $W + U$ is indefinite. If positive definite, its actual face would make opposite sides meet. If positive semidefinite and singular, it would give an ideal point of the compact Q. Both are impossible. Likewise $W + V$ is indefinite. Minimal indefinite subdiagrams $L \subset W + U$, $M \subset W + V$ contain their respective black pairs and at most eight whites. They cannot be orthogonal, since their negative vectors would give negative index at least two. A shortest path in their overlapping or adjacent union between U and V has only white interior nodes and length at most $8 + 1 + 8 = 17$. Its black endpoints are consecutive sides, so the path belongs to an actual corner.

Thus unit weight at ambient spherical distance at most 17 pays two on a compact triangle and one on a compact quadrilateral. Its cost at an ambient finite vertex is at most $17n$: spherical components of order at most 35 have at most $\binom{r}{2} \le 17r$ pairs, and larger components are paths or D forks with at most r pairs at each fixed distance. This uses only the finite Coxeter classification, allowing affine subdiagrams elsewhere in the chamber.

A compact facet has dimension $d = n - 1$ and is simple, because its vertices are finite vertices of the Coxeter chamber. Its two-faces are compact faces of the ambient chamber; its angles count a subset of the ambient vertex pairs. Hence their total is at most $17(d + 1)f_0$. Its nonnegative symmetric h-vector has $h_0 = h_d = 1$. Pairing the coefficients $\binom{k}{2}$ and $\binom{d-k}{2}$ gives

$$\frac{f_2}{f_0} > \begin{cases} d(d-2)/8 & d \text{ even}, \\ (d-1)^2/8 & d \text{ odd}. \end{cases}$$

The inequality is strict because the nonzero endpoint coefficients exceed the middle minimum. Consequently the average required payment $5f_2/f_0 - \binom{d}{2}$ exceeds $d(d-6)/8$ or $(d-1)(d-5)/8$. After subtracting $17(d + 1)$, the differences at $d = 143 + 2t$ and $144 + 2t$ are respectively $t^2/2 + 36t + 3/2$ and $t^2/2 + 73t/2 + 19$, positive for $t \ge 0$. Thus a compact facet forces $n \le 143$.

A finite-volume facet with no ideal vertices is compact. Apply Nikulin's theorem and split by whether the chosen facet has a cusp. The cusp branch above excludes every $n \geq 164$, and the compact-facet branch already excludes $n \geq 144$. This completes the unbounded tail without assuming that the facet itself is a Coxeter polytope.

Proposition 3.13 (Independent higher-dimensional exclusion)**.** *There is no finite-volume hyperbolic Coxeter polytope in any dimension $n \geq 59$.*

Proof. The dimension-specific families exclude $59 \leq n \leq 163$. For $n \geq 164$, the framing-wall and two-budget argument gives $-(x, y) \geq 14$, contradicting narrowness, if the narrow facet has a cusp. If it has none, the compact-facet comparison already excludes $n \geq 144$. These alternatives exhaust the facet supplied by the narrow-facet theorem. □

Together with the contradiction in dimension 58, this proves Theorem 1.1(i).

4 The compact bound

References with a prefix B in this section are to the numbered proofs in Part V of the fixed complete edition (Addendum C). Catalogue entries C1 and C2 locate those proofs and their verification records. The general projection and reference arguments, the small-defect and rank-two proofs, and the final dotted-head argument are developed here. The longer component inventories and the separately stated remaining case exclusions retain their full proofs in that addendum.

Our main result is the following dimension bound.

Theorem 4.1 (Dimension bound)**.** *Every compact real hyperbolic Coxeter polytope has dimension at most* 17*.*

Here compact means that every vertex is finite.

The central object is one vector x, obtained from an original farthest vertex p. Energies of orthogonal components of its incident diagram can be added because they are projections of this same vector. Pressures arising from different auxiliary points cannot in general be optimized separately and then added. Instead, we combine them in one reference vector and retain its complete squared norm.

The case split occurs at the finite irreducible components of the diagram at p. Small components pay uniform energy debits. A large A-component is handled in its zero-sum chain coordinates. A large B- or D-component has a long common prefix; its terminal nodes then require separate analysis. Section 4.4 records a fresh complete cover at dimension 18, and the separate higher-dimensional cover. A *cover* always means necessary possibilities, not realized examples. Section 4.5 explains the whole-B_{18} terminal partition. Sections 4.6–4.9 give the whole-D_{18} closure.

The final composition has two parts. We exclude every dimension at least 19 by its actual-rank forest cover and local consumers. At dimension 18, the complete cover leaves only a whole D_{18} diagram at the same selected farthest vertex.

4.1 Farthest vertices and projections

Use the hyperboloid model with Lorentz signature $(- + \cdots +)$. Every wall has an outward unit spacelike normal n, and the polytope is given by $\langle r, n\rangle \leq 0$. Distinct actual wall normals have nonpositive products. Fix a strict interior point z, choose an actual vertex p farthest from z, and write

$$z = E(p - x), \qquad E = -\langle z, p\rangle > 0, \qquad x \in p^{\perp}, \qquad \|x\|^2 = 1 - E^{-2} < 1. \tag{4.1}$$

The space $p^{\perp}$ is Euclidean. The incident normals at p form a positive definite spherical diagram; compact acute simplicity implies that there are exactly n of them in dimension n. The whole irreducible components T have orthogonal projections with

$$\sum_{T} e_T = \|x\|^2 < 1, \qquad e_T = \|x_T\|^2. \tag{4.2}$$

We retain these original normals and this original choice of p, z, x, E throughout an energy argument.

The function $r \mapsto -\langle z, r\rangle$ takes its maximum on a compact hyperbolic polytope at a vertex. Its restriction to a unit speed geodesic has positive second derivative equal to the function itself, so a maximum on a segment occurs at an endpoint; successively passing to proper faces proves the assertion. Consequently

$$-\langle z, r\rangle \le E \quad \text{for every actual point } r \in P. \tag{4.3}$$

At an adjacent vertex q, write $z = E_q(q - y)$. Equation (4.3) gives a comparison factor E/E_q, not a claim that q is farthest.

4.1.1 Positive inverse matrices and feasible projections

Inverse positivity is Lemma 2.1. We record its explicit projection formula in the coordinates used below.

Lemma 4.2 (An actual positive projection)**.** *Let n_j, $j \in J$, be actual wall normals with positive definite Gram matrix B_J, and let z be strict interior. Put*

$$q_0 = z - \sum_{i,j\in J} (B_J^{-1})_{ij}\langle z, n_j\rangle n_i.$$

Then q_0 is future timelike, orthogonal to all selected walls, and has negative product with every unselected actual wall. Its normalization is a point of the original polytope.

Proof. The added coefficients are nonnegative by Lemma 2.1. For an unselected wall every added product is nonpositive and the original product is negative; the selected products vanish. Moreover

$$q_0^2 = z^2 - (\langle z, n_i\rangle)_{i\in J}^{T} B_J^{-1} (\langle z, n_i\rangle)_{i\in J} < 0.$$

Since $\langle z, q_0\rangle = q_0^2 < 0$, it has future orientation. All original wall inequalities survive normalization. If fewer than n walls are selected this is a point on a face, not an asserted full-dimensional vertex. □

The same sign check applies to positive combinations of actual normals below: selected products are calculated explicitly; every unknown original wall has nonpositive product with every summand. A nonzero feasible future null vector would give a point of the ideal boundary of P, contradicting compactness. A positive affine Coxeter kernel on actual walls is an obstruction after its future sign and all remaining wall products have been checked.

Lemma 4.3 (Future-point pressure)**.** *Suppose $U \in p^{\perp}$, $\xi = Ap - U$ is future timelike with $\xi^2 = -k < 0$, and its normalization is feasible against every actual wall. Then the same original vector satisfies*

$$\langle x, U\rangle \ge A - \sqrt{k}.$$

At a different actual vertex q, use the decomposition $\xi = Aq - U$ with $U \in q^{\perp}$. With all-point comparison factor κ, the corresponding inequality is $\langle y, U\rangle \ge A - \kappa\sqrt{k}$.

Proof. The spatial hypothesis gives $A = -\langle p, \xi\rangle > 0$. Apply (4.3) to $\xi/\sqrt{k}$: $-\langle z, \xi/\sqrt{k}\rangle = E(A - \langle x, U\rangle)/\sqrt{k} \le E$. The modified comparison is the identical calculation. □

For any Euclidean subspace $S \subset p^\perp$ containing U, let x_S be the orthogonal projection of x onto S. If $U \neq 0$, then $\|U\|^2 = A^2 - k$; Cauchy therefore gives

$$\|x_S\|^2 \ge \frac{A - \sqrt{k}}{A + \sqrt{k}}. \tag{4.4}$$

The case $U = 0$ gives the same nonnegative bound directly. Normalize a future vector, not a spacelike correction. For a cap $c = Ap - v$, clear a positive definite old block J:

$$w = \sum_{i,j\in J} (B_J^{-1})_{ij}\beta_j n_i, \quad \beta_j = -\langle c, n_j\rangle \ge 0, \quad \eta = c + w, \quad \delta = \eta^2 = 1 - \beta^T B_J^{-1}\beta.$$

If $\delta < 0$, η is the future vector to normalize. If $\delta > 0$, the actual finite projection instead uses

$$p + \frac{A}{\delta}\eta, \qquad \left(p + \frac{A}{\delta}\eta\right)^2 = -1 - \frac{A^2}{\delta}.$$

When its selected products are nonpositive, its positive coefficients check all unselected walls. The pressures are

$$\langle x, v - w\rangle \ge A - \sqrt{-\delta}, \qquad \langle x, v - w\rangle \ge \frac{A}{1 + \sqrt{\delta/(A^2 + \delta)}}. \tag{4.5}$$

Old normals lie in $p^\perp$, so $v - w$ denotes the spatial part of the corresponding displayed Lorentz vector.

4.1.2 Opposite walls and common components

Let B be the original incident Gram matrix, $C = B^{-1}$, and

$$\omega_i = \sum_j C_{ij} n_j, \qquad d_i = C_{ii}, \qquad u_i = \omega_i/\sqrt{d_i}.$$

An actual opposite wall has normal

$$c = Ap - v, \quad A > 0, \quad v = \sum_i \beta_i\omega_i, \quad \beta_i = -\langle c, n_i\rangle \ge 0, \quad A^2 = \beta^T C\beta - 1.$$

Strict interiority gives

$$\langle x, v\rangle < A. \tag{4.6}$$

Along the original edge in direction $-u_i$, of length L_i, the original farthest comparison gives

$$\langle x, u_i\rangle \ge \tanh(L_i/2). \tag{4.7}$$

Whole-component projection energies sum to $\|x\|^2$.

Proposition 4.4 (Reciprocal shadows; internal geometric interface)**.** *For an actual cap put $T_i = (C\beta)_i/(A\sqrt{d_i})$. If $T_i > 1$, it is the opposite wall of edge i, and $\tanh L_i = 1/T_i$. If it is not that opposite wall, then $\tanh L_i \ge T_i$. The value $T_i = 1$ is incompatible with a finite endpoint. For distinct actual opposite walls,*

$$\tanh L_i \tanh L_j \ge K_{ij}, \qquad K_{ij} = C_{ij}/\sqrt{d_i d_j}. \tag{4.8}$$

Opposite walls of orthogonal original directions cannot coincide.

Proof. The Schur identity says that $C - C_{\cdot i}C_{i\cdot}/d_i$ vanishes on row and column i; elsewhere it is the inverse of the retained old Gram matrix. It is entrywise nonnegative. Let $d = Dp - \sum_j \gamma_j \omega_j$ be the actual opposite of edge i, distinct from c. Acuteness and the Schur identity give

$$AD \geq \beta^T C\gamma \geq (C\beta)_i (C\gamma)_i / d_i.$$

The actual endpoint equation is $\tanh L_i = D\sqrt{d_i}/(C\gamma)_i$, so $\tanh L_i \geq T_i$. If $T_i > 1$ this alternative is impossible, forcing $c = d$; its endpoint equation gives $\tanh L_i = 1/T_i$. If $T_i = 1$, the distinct alternative is impossible and the same-wall equation would give an ideal endpoint. This is excluded by compactness. For the pair inequality, first use the contacts of the actual opposite of i to get $T_j \geq K_{ij}/\tanh L_i$; the distinct alternative at edge j yields the assertion.

Finally suppose the same wall closes orthogonal directions i, j. Clearing every other old normal adds positive coefficients and gives

$$c_F = Ap - \frac{A}{r_i} u_i - \frac{A}{r_j} u_j, \qquad r_i = \tanh L_i, \quad r_j = \tanh L_j.$$

Its product with the strict interior point remains negative, so $\langle x, u_i\rangle/r_i + \langle x, u_j\rangle/r_j < 1$. But (4.7) makes this sum at least $1 + (\tau_i^2 + \tau_j^2)/2 > 1$, where $\tau_i = \tanh(L_i/2)$. This contradiction proves the orthogonal assertion. □

The displayed pair inequality concerns full tangents, not half tangents. For an opposite wall, its retained foreign component can have at most one contact: two would make a cycle in its actual finite replacement. A positive foreign cost is $q = \beta_j^2 C_{jj} \geq 1/4$. In a nontrivial component $C_{jj} \geq 4/3$, so the only cost below $1/3$ is a simple 3-contact on an A_1, of cost $1/4$. Moreover

$$q < A^2. \tag{4.9}$$

Otherwise the actual reciprocal test would close that foreign direction too, violating the orthogonal assertion, with the equality case giving a null endpoint. Set $F(t) = 2t/(1+t^2)$. For unit directions of correlation k in a component of energy at most e, positive heights satisfy

$$\tau_i + \tau_j \leq \sqrt{2e(1+k)}, \qquad F(\tau_i)F(\tau_j) \leq \frac{4s}{(1+s)^2}, \quad s = \frac{e(1+k)}{2}. \tag{4.10}$$

The second inequality follows from concavity of $\log F$ on $(0,1)$. If its upper bound is smaller than K_{ij}, Proposition 4.4 forces a common actual opposite. Its finite-white extension or positive affine kernel then gives additional actual pressures.

4.2 Quadratic estimates from reference vectors

The elementary estimate behind every certificate is

$$\|x\|^2 \geq 2\langle x, W\rangle - \|W\|^2. \tag{4.11}$$

Every term in $\langle x, W\rangle$ must be justified. Write $w = \sum_i b_i P_i$, $b_i \geq 0$, with actual pressures $\langle x, P_i\rangle > a_i$, and demand $A_w = \sum_i b_i a_i$. For a new actual pressure $\langle x, Z\rangle \geq h$, choose

$$Q = Z - \sum_i t_i P_i, \qquad W = w + \mu Q.$$

Before using pressures require $\mu \geq 0$ and check every $b_i - \mu t_i \geq 0$. Then

$$\langle x, W\rangle \geq A_w + \mu(h - \sum_i t_i a_i), \quad \|W\|^2 = \|w\|^2 + 2\mu\langle w, Q\rangle + \mu^2\|Q\|^2. \tag{4.12}$$

The available price b_i is individual to anchor i. The first inequality is strict if some strictly constrained old anchor retains a positive coefficient, or if $\mu > 0$ and the new pressure is strict. For multiple corrections every new weight must be nonnegative; their combined debit at each anchor and all mutual cross terms must be retained.

For $P_i = e_1 + \cdots + e_i$, increasing concave interpolation through $(0, 0)$ and actual floors $\langle x, P_i \rangle > f_i$ gives a reference with successive slopes as its coordinates. Its coefficients in the P_i are nonnegative slope decreases, and its demand equals its square. Deleting a knot between intervals of lengths a, b and slopes s_1, s_2 loses $ab(s_1 - s_2)^2/(a + b)$. For unit adjacent spacings this is half the squared slope difference. For the $\sqrt{i}-$ constant floors at $i \geq 5$, such a unit-neighbor loss is $< 1/3200$; nonadjacent omissions have disjoint intervals and their losses add. End omissions use their exact chord costs. An omitted floor has coefficient zero.

References are Euclidean vectors: actual feasible points prove the pressures, and nonnegative coefficients combine them in (4.11).

4.3 Component estimates

4.3.1 Ordinary components and exceptional actual defects

All bounds below concern the same original projection, with arbitrary finite spectators.

Component	Energy greater than	Component	Energy greater than
ordinary nonshort A_1	$1/10$	ordinary A_2	$319/2000$
ordinary $I_2(m), m \geq 4$	$1/7$	A_3	$1/4$
B_3, H_3	$2/7$	A_4	$2/7$
B_4, D_4, F_4	$1/3$	H_4	$11/36$
A_5	$3/8$	A_6	$3/7$
B_5, B_6	$3/7$	D_5, D_6	$5/14,\ 3/7$
B_7, D_7	$1/2$	B_8, D_8	$4/7$
E_6, E_7, E_8	$9/20,\ 5/9,\ 2/3$		

An actual isolated dotted short A_1 is denoted I. A second short-wall profile J uses two old A_1 factors whose joint energy exceeds .34. The two typed rank-two exceptions, each of energy $> .09$, are

$$N_7 = (\text{old } A_2,\ \text{new isolated native } 7), \qquad O_7 = (\text{old } I_2(7),\ \text{new isolated native } 3).$$

A bare $I_2(7)$ is not automatically O_7. There is at most one short I/J profile or one typed exception; they do not coexist. Acuteness of their actual isolated walls is essential: two on orthogonal supports would have product $-\sqrt{h_1 h_2}$ strictly between $-1/2$ and 0. Appendix B.1 proves the endpoint estimates, the J case and the ordinary rates.

Useful joint bounds, counting I only once, are

$$I + A_1 > 1/5, \quad I + 2A_1 > 3/10, \quad I + T_2 > 21/100, \quad I + T_3 > 2/7. \tag{4.13}$$

Here T_j is ordinary irreducible of rank j. Both typed exceptions force another T_2 to cost $> 1/5$. Another A_1 costs $> 1/5$ beside N_7 and $> 2/15$ beside O_7. For the latter, isolated height $h = \frac{1}{4}\csc^2(\pi/7) - 1 < 1/3$ gives squared full tangent greater than $3/7$. Squared half tangent exceeds $2/15$, since $4(2/15)/(1 + 2/15)^2 = 120/289 < 3/7$. The complete contact and defect proofs, including the actual-wall height constraints behind (4.13), are given in Appendix B.1. The uniform actual B/D floor is proved in Appendix B.2.

4.3.2 Proofs of the small defects and rank-two costs

For the following small-component calculations, let T be the actual whole component, $C_T = B_T^{-1}$ its inverse Gram matrix, and β_T the cap's contact vector on T. Write

$$q = \beta_T^T C_T \beta_T, \qquad Q = \sum_{U \neq T} \beta_U^T C_U \beta_U, \qquad h = A^2 = q + Q - 1 > 0.$$

Thus q denotes the own-component cost and Q the full foreign cost.

Lemma 4.5 (Common-wall forcing and isolation)**.** *Suppose the energy of a component is at most e, with $0 \le e < 1$. A pair for which the right side of (4.10) is below K_{ij} has one actual opposite. If one wall closes every direction of the component, then, with $v_i = (C\beta)_i/\sqrt{d_i}$ and $R_e^2 = 4e/(1+e)^2$,*

$$q - 1 \le h \le R_e^2 \min_i v_i^2. \tag{4.14}$$

If $h < 1/4$, it has no foreign contact. In a path, a wall closing every direction can have multiple contacts only at the two endpoints; in a D_4 star it has at most one.

Proof. The first assertion is (4.10). For a closed direction, $\tanh L_i = A/v_i$ and $\tau_i^2 \le e$, which gives (4.14). Every nonzero foreign cost is at least $1/4$ and is strictly below h by (4.9); hence the isolation assertion. If the old path between two contacts omits a direction closed by the wall, deleting that direction leaves a cycle in an actual finite replacement. This is impossible. The same argument applies to every two contacts in a D_4 star. Finally, a single-contact own cost $q = 1$ gives an affine principal set, impossible even in the presence of further spectators. □

4.3.3 The short directions and their exact debits

Call an A_1 direction short when $\tanh^2 L < 1/3$. For a short A_1 opposite put $b = \beta^2$, $h = A^2$, and let Q be its full foreign cost. Then

$$h = b + Q - 1, \qquad h < b/3, \qquad Q < 1 - 2b/3.$$

Equation (4.9) rules out two foreign components. With one, its cost must be the sole value below $1/3$, namely an A_1 contact 3 of cost $1/4$. Thus the only possibilities are

$$I:\ \text{isolated dotted } A_1, \quad 0 < h < 1/2; \qquad J:\ \text{dotted } A_1 + A_{1,3}, \quad 1/3 < h < 3/8.$$

For completeness, with one foreign component, $Q < h = b + Q - 1$ forces $b > 1$, and $Q < 1/3$ then forces $Q = 1/4$. For J, the heavy edge contributes more than $A/2$ to $x \cdot v$. The light shadow contributes at least $A(1 - \sqrt{1 - 1/(4h)})$; this is at least $A/2$ if $h \le 1/3$. Strict (4.6) therefore gives the asserted lower bound. Native acuteness excludes two short walls. Indeed their height product is below $1/2$, so their product must be zero. Disjoint supports give a negative product, heavy overlaps give a spatial product above $1/2$, and a sole light/light overlap of two J walls gives $1/4$, below their height product. None is zero. A J heavy direction has $\tanh^2 L > 4/13$, hence energy $> 9/100$; its light direction has $\tanh^2 L > 2/3 > 16/25$, hence energy $> 1/4$. Thus J funds both A_1 factors above their combined ordinary baseline $2/10$. Every nonshort direction pays at least $5 - 2\sqrt{6} > 1/10$, since $\sqrt{6} < 49/20$. We may conservatively discard the one short direction, but only the isolated type I can create an unfunded debit.

The rank-two exceptions are

old block / new isolated contact	$h = A^2$	energy
$A_2/7$	$(4/3)\cos^2(\pi/7) - 1 \in (0, 1/12)$	$> 9/100$
$I_2(7)/3$	$(1/4)\csc^2(\pi/7) - 1 \in (0, 1/3)$	$> 9/100.$

(4.15)

Their spatial supports lie entirely in their own blocks. Two defects in different blocks, or a defect and a short A_1, would have actual wall product $-\sqrt{h_1 h_2} \in (-1/2, 0)$. That interval is forbidden: distinct acute unit walls have product 0 or at most $-1/2$. Therefore

$$\text{at most one short } A_1, \quad \text{or one rank-two defect, but never both.} \tag{4.16}$$

This argument is independent of every other component type.

4.3.4 The two rank-two estimates

Write $\varphi = (1+\sqrt{5})/2$. For A_2, set $e_2 = 319/2000$ and $\sinh^2 L_0 = 31/50$. If both edges have length at least L_0, then their half-tangents exceed $173/500$ and energy is greater than $4 \cdot 173^2/(3 \cdot 500^2) > e_2$. A shorter edge cannot have two positive own contacts: $3/3$ is affine, while $3/4$ or larger gives $\sinh^2 L \geq 4\sqrt{2}/9 > 31/50$, increasing with either contact and with Q. If only the other node is contacted, apply the same test to that shorter contacted direction. For one contact of square b, the shorter-edge condition is

$$h = 4b/3 + Q - 1 < 124b/243, \qquad Q < 1 - 200b/243. \tag{4.17}$$

Every positive foreign cost must be below h. A label 3 gives $h < 1/4$, hence isolation and negative height. For a label 4, $h < 62/243$ forces each foreign cost to be $1/4$; the only positive possibility is $Q = 1/2$, but then $h = 1/6 < 1/4$. A label 5 has $Q < 1/2$, hence at most one foreign component, whose cost exceeds h because its own cost $\varphi^2/3$ is below one. A label 6 is affine. For a finite label $m \geq 7$, $b > .81$ gives $Q < 1/3$; thus $Q = 0$ or $Q = 1/4$. The latter is possible in (4.17) only for $b < 729/800$, namely $m = 7, 8, 9, 10$. Indeed $\cos^2(\pi/11) > 45/49 > 729/800$, using $\pi < 22/7$. A dotted contact has $Q < 1/4$ and is isolated.

For an isolated finite contact the same wall closes the mate, since its reciprocal ratio squared is $b/(4b-3) > 1$. For $m \geq 8$ that actual mate has $\tanh^2 L \geq 6\sqrt{2} - 8 > 400/841$, so its energy exceeds $4/25 > e_2$. For an isolated dotted contact, $b < 243/200$ and its distinct mate has shadow $\tanh^2 L \geq b/(4b-3) > 81/124 > 16/25$. For a mixed $m = 7, 8, 9, 10$ contact, the mate's white cost is $b + 1/4 > 1$; the distinct shadow instead gives $\tanh^2 L \geq b/(4b - 9/4) > 11/17 > 16/25$. Both give energy greater than $1/4$. Thus the only exception is isolated new 7 at old A_2. The root $t = \cos(2\pi/7)$ of $8t^3 + 4t^2 - 4t - 1$ lies in $(623/1000, 5/8)$: the polynomial changes sign there and is increasing on $[1/2, 1]$. Hence $b > 1623/2000$ and its actual mate satisfies $\tanh^2 L = 4 - 3/b > 164/541 > 3600/11881$, proving its cost $> 9/100$.

For an old $I_2(m)$, $m \geq 4$, put $k = \cos(\pi/m)$ and $d = (1-k^2)^{-1}$, so $C = d \left(\begin{smallmatrix} 1 & k \\ k & 1 \end{smallmatrix}\right)$. At energy at most $1/7$, (4.10) forces one common opposite. Two contacts give an actual $\cosh L \geq \cot(\pi/(2m))/\sqrt{3} > 11/8$, with individual energy $> 3/19 > 1/7$. For one contact β,

$$q(1 - 7k^2/16) \leq 1, \qquad q = d\beta^2. \tag{4.18}$$

Its minimum left side is $(9d+7)/64$. Since $m \geq 8$ gives $d \geq 4 + 2\sqrt{2} > 19/3$, only four old labels remain:

m	consequence of (4.18)
4	$\beta^2 \leq 16/25 < \cos^2(\pi/5)$: 3, 4; $h \leq 7/32 < 1/4$, $q \leq 1$;
5	$d > 25/9$: only 3; $h < 7/32$, $q < 1$;
6	only 3; $q = 1$;
7	$d > 5$: only 3; $Q < 3/16$.

Isolation or the affine principal excludes the first three rows. The last is isolated new 3 at old $I_2(7)$, with $0 < h < 1/3$. Its actual mate has $\cosh L = \cot(\pi/7)/\sqrt{3} > 109/91$ by the same cubic bracket; the squared comparison has positive numerator 2652. Thus its energy also exceeds $9/100$, completing (4.15) and (4.16).

4.3.5 The A-chain branch

For A_r, use the zero-sum subspace of $\mathbb{R}^{r+1}$ with roots $(e_i - e_{i+1})/\sqrt{2}$. The inverse Gram matrix is

$$C_{ij} = \frac{2\min(i,j)(r+1-\max(i,j))}{r+1}.$$

A finite-white classification of actual opposite walls gives strong dual floors except for a short list of literal helper caps. Adjacent weak helpers are excluded using their actual mutual product and a positive two-cap projection. Concave interpolation uses only strong floors; helper pressures are paid at surviving anchors. The small end variance yields stronger component bounds:

r	$r \geq 7$	$r \geq 8$	$r \geq 13$	16	17
$e(A_r) >$	1/2	4/7	5/7	199/250	6/7

$e(A_{18}) > 91/100.$

Later local refinements give $e(A_{12}) > 13/20$ and $e(A_{15}) > 413/500$, uniformly with arbitrary spectators. The shared actual feet, common-pair inventories, uniform $1/2, 4/7, 5/7$ bounds, and the $N = 17, 18, 19$ central-window and paid-reference arguments are proved in Appendix B.3. The newer $A_{16} > .86$ proof in particular implies the older .796 bound without requiring its separate historical reference. The fresh smaller-rank proofs are in Sections B.4.5 and B.4.4. Their full native-cosine inventories and individual actual-helper payments are given there.

4.3.6 The common B/D prefix branch

For B_r, use long roots $(e_i - e_{i+1})/\sqrt{2}$, $i < r$, and short root e_r. For D_r, use the long chain and the fork root $(e_{r-1} + e_r)/\sqrt{2}$. On a long initial segment, $\omega_i = \sqrt{2}P_i$, $C_{ij} = 2\min(i,j)$. In type D, $x_1 > \cdots > x_{r-1} > |x_r|$; the last coordinate is signed and the forks are distinct actual duals.

Proposition 4.6 (Generic prefix theorem)**.** *Let $6 \leq M \leq 16$. Suppose an actual component is B_r, $r \geq M+5$, or D_r, $r \geq M+6$. Put*

$$f_i = \sqrt{i} - \frac{61}{100\sqrt{2}}, \quad E_M = \frac{f_5^2}{5} + \sum_{i=5}^{M-1} (\sqrt{i+1} - \sqrt{i})^2, \quad H_M = \left\lfloor \frac{M-5}{2} \right\rfloor.$$

With arbitrary finite spectators, $\sum_{j=1}^{M+1} x_j^2 > E_M - H_M/3200$.

Proof. Appendix B.5 proves that a weak site $5 \leq i \leq M+1$ has a pure cap

$$v = \gamma\omega_i + \omega_1/2, \quad A^2 = 2i\gamma^2 + 2\gamma - \tfrac{1}{2}, \quad 101/125 < \gamma < 6/5,$$

with own label 5, finite label at least 7, or dotted own contact, and no spectator contact. Actual weak sites are nonadjacent. This is proved at these ranks, not transferred from a whole-component exclusion. If 5 is strong, choose a strong $K = M$ or $M+1$. Interpolate through actual strong knots. Each missing internal floor loses $< 1/3200$; at $K = M+1$ the extra increment $(\sqrt{M+1} - \sqrt{M})^2 > 1/(4M+2) \geq 1/66$ pays the possible extra omission.

If 5 is weak, the first-six-root point table in Appendix B.5 gives $x_1 > .5965$, and 6 is strong, with $S_6 > 2.01815$. Interpolate through $(1, .5965), (6, 2.01815)$ and a later strong endpoint. The slopes decrease: $.5965, .28433$, then $1/(\sqrt{K} + \sqrt{6}) < 1/5$. Its energy exceeds

$$L_M = .7600299945 + \frac{(\sqrt{M} - \sqrt{6})^2}{M-6},$$

the last term being zero at $M = 6$. Now $E_6 < .7$. The $k = M-6$ increments from 6 onward lie in $(1/8, 1/5)$, so their variance is at most $k(1/5 - 1/8)^2/4 \leq 9/640$. Thus $L_M - E_M > .76 - .7 - 9/640 > 0$, directly also at $M = 6$. This proves the weak-5 branch without first assuming strong 5. □

Appendix B.5 supplies the complete finite-white inventory, zero-own repair, actual two-cap nonadjacency, all four weak-5 pressures and continuous seals, with every spectator retained. Its original rank inequalities are part of the theorem. A short-tail continuation at an excluded rank still requires its own new local proof.

4.4 The remaining spherical diagrams

Proposition 4.7 (Compact rank cover and its consumers)**.** *At an original farthest vertex of a compact Coxeter polytope of dimension* 18*, the complete necessary cover is the one displayed below. Its proper-component rows and its whole A_{18} and B_{18} rows are excluded by the indicated case proofs. The higher-dimensional exclusions are separate statements at their actual ranks.*

The accounting leading to the cover is given below, followed by exact locations of the longer row proofs. None of the retained rows asserts a realized polytope.

4.4.1 A fresh cover in exact dimension 18

Call A_r, $r \geq 7$, or B_r, D_r, $r \geq 9$, dominant. Each costs $> 1/2$, so at most one is possible. With none the ordinary rank-over-14 baseline, even allowing the unique defect, exceeds 1. Exactly one is therefore present. The fresh total-rank-18 necessary cover is:

Dominant	All retained spectator alternatives
A_{12}	$I + (A_5$ or $D_5)$, or $N_7/O_7 + T_4$
A_{15}	ordinary A_3; $T_2 + A_1$; $I + T_2$; or $O_7 + A_1$
A_{16}	ordinary T_2; two ordinary A_1; $I + A_1$; or N_7/O_7
A_{17}	one ordinary nonshort A_1, or I
A_{18}	none
B_{16}	N_7/O_7
B_{17}	I
B_{18}	none
D_{16}	N_7/O_7
D_{17}	one ordinary nonshort A_1, or I
D_{18}	none

Here ordinary $T_2 = I_2(m)$, $m \geq 3$, and $T_4 \in \{A_4, B_4, D_4, F_4, H_4\}$. H_4 and native 5 are not discarded by field arguments.

For completeness, the accounting steps are as follows. A_7 leaves too many ordinary ranks; $A_8, \ldots, A_{11}$ are excluded by 4/7 plus the rank-over-14 baseline. Beside A_{12}, ordinary or J rank-six spectators cost $> 3/7$. With I, (4.13) excludes all rank-five partitions except A_5, D_5. With a typed exception, partition rank 4: two T_2's cost $> .4$, $T_3 + A_1$ costs $> 1/4 + 2/15$, and the other reducible partitions are stronger. Only irreducible T_4 remains. For A_{13}, A_{14}, the 5/7 bound and (4.13) exclude all spectators. At A_{15}, exclude B_3, H_3, three ordinary A_1's, $I + 2A_1$, $J + A_1$, and $N_7 + A_1$. At A_{16}, .796 excludes J but retains precisely the listed rank-two alternatives. At A_{17} retain the remaining A_1.

For B_r, D_r, the 9/14 bound forces spectator rank at most 5, hence $r \geq 13$. Proposition 4.6, within its rank domain, gives

r	13	14	15	16
$e(B_r), e(D_r) >$	.7349975	.7683275	.797425	.823735.

At $13, 14, 15$, every ordinary, I/J, and typed remainder exceeds the residual budget; $I + T_2 > .21$ closes rank 15. At 16, two ordinary A_1's and $I + A_1$ are excluded; ordinary T_2 and both typed

defects initially remain. The B_3, D_4, D_3 layer checks in Appendix B.6 extend helper coverage to 13, giving energy $> .8689525$; ordinary $T_2 > 1/7$ closes that case. The fresh B_{17} layers reach a strong 14/15 endpoint and energy $> .90715$, excluding a nonshort ordinary A_1. This reuses the interpolation proof after new local coverage, not Proposition 4.6 outside its legal ranks.

The first rows close with small shared consumers:

$$A_{12} + I + (A_5/D_5) > 13/20 + 5/14 = 141/140, \quad A_{12} + N_7/O_7 + T_4 > 13/20 + .09 + 2/7 = 359/350.$$

The improved 413/500 bound closes every A_{15} family: ordinary total rank three costs $> 17/70$, $I + T_2 > .21$, and beside $O_7 + A_1$ the total exceeds 787/750. Thus the older separate ordinary-rank-three consumer is unnecessary. The other internal proof homes are:

Row	Excluding argument
A_{16}	Proposition B.3.5, ordinary and $I + A_1$; Appendix B.10.1, both actual typed defects
A_{17}	Appendix B.10.2, ordinary and actual I
whole A_{18}	Appendix B.7, full two-end native-gap argument
B_{16}	Appendix B.10.3, both actual typed defects
B_{17}	Appendix B.10.5, actual I; local head and helper kernels
D_{16}	Appendix B.10.3, both actual typed defects
D_{17}	Appendices B.10.4.1 and B.10.6, ordinary and actual I
whole B_{18}	Section 4.5
whole D_{18}	Sections 4.6–4.9

Appendix B.6 proves the local layers used in this cover, including the distinct signed D_4/D_3 forks and the actual A_{17} neighbor for B_{17}. The table identifies the complete proof of every retained row, including both short rank-one cases. The independent higher-dimensional exclusions are proved in Appendices B.11–B.14. These arguments use no field, lattice or integral dotted-product hypothesis.

4.4.2 The separate complete higher-dimensional tail

The exact-18 cover does not exclude higher dimensions. The earlier internal package establishes the separate theorem:

Theorem 4.8 (Internal higher-dimensional exclusion)**.** *There is no compact hyperbolic Coxeter polytope of dimension at least* 19*.*

Appendix B.11 proves the component caps, shared windows and complete actual-rank exclusions from 29 through 20. The published ceiling is used only at dimensions at least 30. At dimension nineteen, Proposition B.11.35 leaves whole A_{19}, B_{19}, D_{19}. Their proofs occupy Appendices B.12–B.14: the A proof splits by the three weak-end patterns; the B proof splits by its two exceptional terminal walls; and the D proof splits by six head types after a common centre analysis. Each case keeps one original vector and every spectator in its cover. Section B.14.22 joins these exclusions.

The higher ranks have their own complete covers in Appendix B.11. The short B_3/B_2 and D_4/D_3 continuations are proved before the prefix estimates that consume them. Section B.14.22 combines these actual-rank arguments with the published compact ceiling. This is a logical conjunction of exclusions, not geometric dimension heredity. No arithmetic theorem or field census enters that conjunction.

4.5 The whole-B_{18} terminal partition

Classify the actual opposite of old node 17 without assuming a strong head. Section B.8.5 leaves:

(i) a pure far-three helper with finite own 5, finite own at least 7, or dotted own $1 < g < 4/3$;

(ii) near-three with own 6;

(iii) no left contact, with short white label $3, 4, 5, 6, 7, 8$.

All other isolated, left, bridge and affine/null roles are excluded in that inventory. Sections B.8.6–B.8.10 exclude the first branch, and Section B.8.11 excludes the second. In the last branch, Sections B.8.13–B.8.14 exclude distinct terminal caps and then every common-cap role. The last common cap is treated at its actual adjacent whole-A_{18} vertex, with comparison 11/8; that vertex is never assumed farthest.

Thus an original farthest whole-B_{18} vertex cannot occur. Appendix B.8 gives the complete finite-white inventory, each terminal exclusion, all original-wall checks and the individual coefficient payments. Together with the preceding row consumers, this leaves only whole D_{18} at the selected original vertex.

4.6 The last case: an actual whole-D_{18} vertex

Proposition 4.9. *The original farthest vertex cannot have whole incident diagram D_{18}.*

We give the central proof below, using the named finite-head and fork-pair case lemmas of Addendum C where indicated.

We treat the final dotted-head case using the complete inventory proved in Appendix B.9. Keep the original p, z, x and use actual old roots

$$n_i = (e_i - e_{i+1})/\sqrt{2} \quad (1 \le i \le 17), \qquad n_{18} = (e_{17} + e_{18})/\sqrt{2}.$$

Write

$$P_i = e_1 + \cdots + e_i, \quad \Omega_i = \sqrt{2}P_i \ (i \le 16), \quad \Omega_{17} = (P_{17} - e_{18})/\sqrt{2}, \quad \Omega_{18} = (P_{17} + e_{18})/\sqrt{2}.$$

In particular x_{18} is signed and $x_1 > \cdots > x_{17} > |x_{18}|$. In this part only, set

$$S_i = \langle x, P_i \rangle, \qquad f_i = \sqrt{i} - \frac{.58}{\sqrt{2}}. \tag{4.19}$$

All terminating decimals in this paper denote exact rational numbers.

The head is the actual opposite wall of the long-end edge 1. Section B.9.1 proves that its normal is

$$H = A_H p - \eta\Omega_1, \qquad A_H^2 = 2\eta^2 - 1 > 0.$$

The same section gives the complete own-contact list. Sections B.9.8 and B.9.9 prove the strong terminal floor $S_{16} > f_{16}$, its funded reference append and the complete fork-role list. The finite-head cases close in Section B.9.22. We treat every dotted head $\eta > 1$ without a parameter cutoff.

Appendix B.9 proves these earlier inputs in geometric dependency order: actual head isolation, the fresh threshold window, strong 16, finite-head closure and the every-head pair exclusions. In particular it proves the changed-threshold strong-site statements afresh rather than inheriting them unconditionally.

4.6.1 A future point valid for every dotted head

Put $u = \eta^{-2} \in (0,1)$, and consider

$$\Xi = H + n_1/\eta.$$

This is a positive combination of two actual normals. Its products with H, n_1 are respectively 0 and $-\eta + 1/\eta < 0$, and $\Xi^2 = -(1-u) < 0$. Its time is $A_H > 0$. Every other actual wall has nonpositive product with each summand. Hence its normalization is an actual future point of the original polytope, with no unknown wall omitted. Lemma 4.3 gives

$$\langle x, Z_u \rangle \geq h(u), \qquad Z_u = \frac{(2-u)e_1 + ue_2}{2}, \quad h(u) = \frac{\sqrt{2-u} - \sqrt{u(1-u)}}{\sqrt{2}}. \tag{4.20}$$

Indeed the spatial part of Ξ, divided by $\sqrt{2}\eta$, is Z_u. We will use

$$\sum_j (Z_u)_j = 1, \qquad R(u) = \|Z_u\|^2 = 1 - u + u^2/2.$$

Equation (4.20) is a point pressure, not an energy lower bound, and is valid on the entire open interval.

4.6.2 The four original reference cases

Section B.9.2 gives the following complete initial alternatives: strong 5; a weak pure far-left-three helper at 5; or an isolated dotted weak 5 with a chosen strong $J = 6$ or 7. Thus there are four reference rows. Here is why the initial classification has this precise scope. Left own 4, including own 4/white 3, gives a positive actual affine-B null on the cap and old chain $5, \ldots, 18$; the extra left contact preserves the remaining wall signs. The newly possible near-three/own-five at 5 is excluded by its actual right-H_4 point and independent old strong 11/12, before the new references are used. Other finite-white and isolated finite roles are strong. A historical fifth paid row concerned a native-five head and is outside $\eta > 1$.

We emphasize the *bare* references for the three nonhelper rows. They interpolate through the strong knots, are flat on their first J coordinates, and have endpoint sum f_{16}. Sections B.9.4 and B.9.8 give

initial case	B_0	first mean $b <$	unused P_J price $>$	J
strong 5	.9565188	.3652	.151	5
isolated dotted 5, strong 6	.9373243	.3399	.1435	6
isolated dotted 5, strong 7	.9196357	.31938	.1366	7

More explicitly, each reference w has

$$\langle x, w \rangle > A_w, \quad 2A_w - \|w\|^2 > B_0, \quad \operatorname{supp} w \subset \{1, \ldots, 16\}, \quad \langle w, P_{16} \rangle = f_{16}, \quad \beta_{16} > .127.$$

No head pressure has been paid. Their raw endpoint-15 lower bounds are $.9403998, .9212053, .9035167$. The legal endpoint-16 append adds at least $.127^2 - .00001 = .016119$; if 15 is weak, its fictitious knot is deleted before pressure. All late repairs preserve the flat first block and its individual prices.

In the isolated-5 rows there is also an unused actual pressure

$$U_5 = P_5 + \frac{v}{2}(e_6 - e_5), \quad .4 < v < 1, \qquad \langle x, U_5 \rangle > 1.8185.$$

For

$$Q_H = Z_u - P_J/J, \qquad Q_5 = U_5 - (5/J)P_J$$

the exact identities are

$$\|Q_H\|^2 = R - 1/J, \quad \langle Q_H, Q_5\rangle = 1 - 5/J, \quad \|Q_5\|^2 < \begin{cases} 77/150, & J = 6, \\ 194/175, & J = 7. \end{cases} \tag{4.21}$$

Both residuals have zero product with the bare flat reference. The positive mutual cross in (4.21) will not be discarded.

For the fourth row, the exact *unheaded* weak-helper reference interpolates from $(1, a_0)$, $a_0 = .5455$, to $(6, f_6)$, then through the later actual strong knots to 16, replacing each missing knot by a chord. Its first six coordinates are

$$w_1 = a_0, \quad w_2 = \cdots = w_6 = b_0 = (f_6 - a_0)/5. \tag{4.22}$$

The actual helper proves $x_1 > a_0$ and strong 6. The unheaded reference satisfies

$$\langle x, w\rangle > A_w, \quad L_w := 2A_w - \|w\|^2 > .987759, \quad \langle w, P_{16}\rangle = f_{16}, \quad \beta_{16} > .127.$$

Weak 7, later omissions, and the late 15 repair do not alter (4.22). Again no head pressure has been paid. Appendices B.9.4, B.9.8 and B.9.20 supply the raw-reference increments, omission tables, isolated-5 pressure and individual coefficient prices used here. These are internal geometric lemmas with written proofs, not script-output assumptions.

4.7 Bounds for the head and fork edges

4.7.1 The first three rows

In the strong-5 row choose $\mu = .33$ on Q_H. Its debit $.33/5 = .066 < .151$ fits the individual price. Equations (4.12)–(4.21) give

$$L_W > .9565188 + .66(h - .3652) - .1089(R - 1/5) = F(u) := .7372668 + .66h(u) - .1089R(u).$$

For isolated 5 with $J = 6$, use weights $\mu = .33, \nu = .10$ on Q_H, Q_5. The combined debit at J is $.83/6 < .1435$. For $J = 7$ use $\mu = .33, \nu = .11$, with debit $.88/7 < .1366$. In each case all remaining old coefficients are nonnegative *before* the new pressures are invoked, and

$$\begin{aligned} L_W > B_0 &+ .66(h - b) + 2\nu(1.8185 - 5b) - .1089(R - 1/J) \\ &- \nu^2\|Q_5\|^2 - .66\nu(1 - 5/J). \end{aligned} \tag{4.23}$$

The constant terms before $.66h - .1089R$ in the two rows are $22164209/30000000$ and $51729823/70000000$, both greater than $.7372668$. Thus F is a common lower bound. No optimized helper and head energies have been added.

We prove the bound uniformly on $0 < u < 1$. Put $a = \sqrt{2 - u}$, $b = \sqrt{u(1 - u)}$. Direct differentiation gives

$$h''(u) = \frac{1/(4b^3) - 1/(4a^3)}{\sqrt{2}} > \frac{7}{4\sqrt{2}}, \tag{4.24}$$

because $b \le 1/2$, $a > 1$. Hence $F'' > .66 \cdot 7/(4\sqrt{2}) - .1089 > .7$, using $\sqrt{2} < 1.415$. At $u_0 = 2/3$,

$$h(u_0) = \sqrt{2/3} - 1/3, \quad R(u_0) = 5/9, \quad F(u_0) > .9956502,$$
$$F'(u_0) = .2013 - .0825\sqrt{6}, \quad |F'(u_0)| < .0008.$$

The value bound uses $.81649^2 = .6666559201 < 2/3$, and the derivative bound uses $2.4494 < \sqrt{6} < 2.4495$. Taylor's inequality and completion of the square on the segment between u_0 and u give

$$F(u) > .9956502 - \frac{.0008^2}{1.4} > .99564 \qquad (0 < u < 1). \tag{4.25}$$

No endpoint derivative or sampled minimum is used.

4.7.2 The weak-helper row: two separate anchor prices

The unheaded reference (4.22) has two available prices

$$\beta_1 = a_0 - b_0 > .24672, \qquad \beta_6 \geq b_0 - (\sqrt{7} - \sqrt{6}) > .1023. \tag{4.26}$$

Indeed $2.0392 < f_6 < 2.0394$, so $.29874 < b_0 < .29878$, while $\sqrt{7} - \sqrt{6} < 2.6458 - 2.4494 = .1964$. An adjacent omitted knot decreases the next slope and increases β_6. The bounds follow from $2.4494 < \sqrt{6} < 2.4495$ and $.4101 < .58/\sqrt{2} < .4102$, by positive squaring.

Define

$$Q = Z_u - (1 - 3u/5)e_1 - (u/10)P_6.$$

Its first coordinate is zero, its second is $.4u$, and its third through sixth coordinates are $-.1u$. Therefore

$$\|Q\|^2 = u^2/5, \qquad \langle w, Q\rangle = 0. \tag{4.27}$$

Choose $\mu = .22$. First pay the *separate* anchor debits

$$\mu(1 - .6u) < .22 < \beta_1, \qquad \mu u/10 < .022 < \beta_6 \quad (0 < u < 1).$$

The remaining original coefficients are nonnegative. Only then add the positive new head pressure μZ_u. The terminal coefficient at 16 remains unused. The full one-vector square gives

$$\begin{aligned} L_{w+\mu Q} &> .987759 + .44[h - a_0 + (3a_0/5 - f_6/10)u] - .00968u^2 \\ &> G(u) := .747739 + .44h(u) + .0542784u - .00968u^2, \end{aligned} \tag{4.28}$$

since $3a_0/5 - f_6/10 > .12336$. This argument rewrites a positive-coefficient pressure decomposition before using it; it does not subtract old lower floors from an unknown signed pressure.

We give the exact small-margin seal. By (4.24), $G'' > .44 \cdot 7/(4\sqrt{2}) - .01936 > .5$. At $u_1 = .642$, positive square tests give

$$\sqrt{1.358} > 1.1653325, \qquad \sqrt{.229836} < .4794122,$$

whose comparison squares are 1.35799983555625 and .22983605750884. The difference exceeds .6859203. Since

$$.4850188 \cdot 1.4142136 = .68592018321568 < .6859203, \qquad 1.4142136^2 > 2,$$

we have $h(.642) > .4850188$, and hence

$$G(.642) > .99200425728.$$

For the derivative use

$$.858 < 1/\sqrt{1.358} < .859, \quad .592 < .284/\sqrt{.229836} < .593, \quad .353 < 1/(2\sqrt{2}) < .354.$$

They follow by multiplying the positive radical brackets $1.1653 < \sqrt{1.358} < 1.1654$ and $.4794121 < \sqrt{.229836} < .4794122$. Thus

$$-.094518 < h'(.642) < -.093545, \qquad .00026136 < G'(.642) < .00068948 < .0007,$$

because $G' = .44h' + .0542784 - .01936u$. Strong convexity on the entire interval now gives

$$G(u) > .99200425728 - .0007^2 = .99200376728 > .992. \tag{4.29}$$

Combining (4.25) and (4.29), every one of the four original initial cases supplies a single legally funded reference W with

$$L_W > .992, \quad \operatorname{supp} W \subset \{1, \dots, 16\}, \quad \langle W, P_{16} \rangle = f_{16}, \quad \text{unused } \beta_{16} > .127. \tag{4.30}$$

All corrections have sum zero and support at most 7; the late omissions and the original endpoint-16 price survive.

4.7.3 Both signed fork directions

Let $\Delta = S_{16} - f_{16} \geq 0$. The pressure decomposition of W keeps its original terminal coefficient. Therefore $\langle x, W \rangle > A_W + \beta_{16}\Delta$, and the same norm identity gives

$$\|x - W\|^2 < .008 - .254\Delta, \qquad x_{17}^2 + x_{18}^2 < .008 - .254\Delta. \tag{4.31}$$

The two true unit fork duals are

$$\frac{\Omega_f}{3} = \frac{P_{17} \pm e_{18}}{\sqrt{18}}, \qquad f = 17, 18.$$

For either actual fork height,

$$\begin{aligned} t_f := \langle x, \Omega_f/3 \rangle &< \frac{f_{16} + \Delta + \sqrt{.016 - .508\Delta}}{\sqrt{18}} \\ &< \frac{3.59 + .1265}{\sqrt{18}} < .876. \end{aligned} \tag{4.32}$$

The numerator decreases with Δ: its derivative is $1 - .254/\sqrt{.016 - .508\Delta} < 0$ on its nonempty domain. The endpoint is handled by continuity. The seals are $f_{16} < 3.59$, $\sqrt{.016} < .1265$, $\sqrt{18} > 4.2426$, and $.876 \cdot 4.2426 = 3.7165176 > 3.7165$. No absolute value has replaced the signed last coordinate.

For an actual fork edge with distance cosine C, (4.7) gives $t_f \geq \sqrt{(C-1)/(C+1)}$. Hence

$$C < \frac{1 + .876^2}{1 - .876^2} < 7.6, \tag{4.33}$$

since $.876^2 = .767376$ and $7.6 \cdot .232624 = 1.7679424 > 1.767376$. In Section B.9.9, every far-next-three role has $g \geq \cos(\pi/5) > .809$ and exact cosine $C = 9g + 1 > 8.281$. Equation (4.33) removes that entire role, including all its previously classified native/null own alternatives.

4.8 A local A_{18} lemma at an adjacent actual vertex

Lemma 4.10. *Let q be an actual vertex of a compact Coxeter 18-polytope whose whole incident diagram is A_{18}. Write $z = E_q(q - y)$, $\|y\| < 1$. Suppose*

$$-\langle z, r\rangle \le \kappa E_q \quad \text{for every actual point } r \text{ of the original } P, \qquad \kappa < 11/8.$$

Then $\langle y, u_{18}\rangle > .78$ is impossible for the positive normalized endpoint-dual direction.

The lemma is local at q, which is not assumed farthest. Appendix B.4 supplies its local inventory; no original B_{18} hypothesis is used.

Proof. In the zero-sum subspace of $\mathbb{R}^{19}$, for $i \le j$,

$$D_i = \|\omega_i\|^2 = \frac{2i(19-i)}{19}, \qquad C_{ij} = \frac{2i(19-j)}{19}.$$

Call $5 \le i \le 14$ weak if $\langle y, \omega_i\rangle \le \sqrt{D_i} - .87$. The complete inventory at comparison $11/8$, established before any endpoint pressure, leaves only:

(a) near-four, meaning own 4 and white 3 at the adjacent old node, in the four literal side-order-5/6 orientations;

(b) a pure far-end-three helper

$$d_i = A_i q - \gamma_i \omega_i - \omega_e/2, \quad e = 1 \text{ or } 18, \quad \varphi/2 \le \gamma_i < 17/16,$$

with own 5, finite own at least 7, or dotted own.

Here $\varphi = (1+\sqrt{5})/2$. All isolated caps are strong. The inventory includes own zero, every finite own, parallel and unbounded dotted owns. Every positive point checks every unknown original wall.

Some calculations locate the numerical boundary. Every ordinary edge ratio at least $3/2$ is strong: with $a = 3/2, k = 11/8$, the square comparison for $q_0 = \sqrt{D_i} \in [2.7, 3.1]$ is the positive polynomial

$$-.21q_0^3 + 1.1876q_0^2 - 1.74q_0 + .7569.$$

It is concave with endpoint values $.583074, .519626$. The near-four ratio of side order ℓ is

$$\frac{1/\sqrt{2} + (\ell-1)/(2\ell)}{\sqrt{(\ell+1)/(2\ell)}}.$$

It increases in ℓ and exceeds $3/2$ at $\ell = 7$, since $(67 + 42\sqrt{2})/56 > 9/4$. Only $i = 5, 6$ with white $i-1$, and $i = 14, 13$ with white $i+1$, can be weak. For pure helpers the identity

$$\frac{(\gamma + 1/(2\ell))^2}{(\ell+1)/(2\ell)} - (4\gamma - 2) = \frac{(\gamma - 1 - 1/(2\ell))^2}{(\ell+1)/(2\ell)} \ge 0$$

gives the stated cutoff at the weakened comparison.

The local input retains genuine helper nonadjacency at this factor. Its proof uses the following exact actual products. For adjacent helper owns $A, B \in [.808, 1.08]$, the mutual product R satisfies

same end	$R \ge 1 - \frac{18}{19}AB - \frac{\lvert A-B\rvert}{19} > -1/2$
outward	$R \ge \frac{1}{2} - AB + \frac{(A-1)(B-1)}{19} > -.7$
reversed	$R \ge \frac{1}{2} - AB + \frac{(A+1)(B+1)}{19} > -.7.$

The native alphabet leaves $R = 0$ in the first case and $R = 0, -1/2$ in the others. Outward $R = 0$ contradicts

$$(D_i - 4/5)(D_{i+1} - 4/5) - C_{i,i+1}^2 = \frac{2}{5}C_{i,i+1} - \frac{24}{25} + \frac{8}{95} \geq 884/475 > 0.$$

For reversed $R = 0$, set $l = i, r = 18 - i$. The exact determinant would require

$$4lrA^2B^2 - (l+1)(r+1)(A^2 + B^2) - 2AB - (l+1)A - (r+1)B + 21/4 = 0.$$

Its own derivatives are positive on that rectangle. If one own is not 5, it exceeds .9; evaluating the other at .808, the endpoints $l = 5, 13$ give $.9601824, 1.6961824 > 0$, and concavity in l excludes all. For two native-five owns the norm-two calculation in basis $1, \varphi$ has nonzero rational determinant coefficient 475 outward and -361 reversed. This uses only native cosine algebra. Thus remaining products are 0 for a common helper and $-1/2$ for opposite helpers.

After clearing old roots, the actual two-cap Schur matrices are $I - t\mathbf{1}\mathbf{1}^T$, $t = (\ell - 1)/(2\ell)$, or the matrix with diagonal $\delta_l, \delta_r \in (1/2, 3/5]$, off-diagonal $-1/2$. Both are positive with positive inverse. Their positive projections are feasible against every original wall, including the reverse wall at q. The cap-time and weight estimates in Appendix B.4 give weighted mean loss less than

$$1/10 + (11/8)(7/16) = .7015625 \quad \text{or} \quad 1/10 + (11/8)(3/10) = .5125.$$

Both are less than .87, excluding adjacent weak helpers. None of these steps uses endpoint height $> .8$.

Now assume the new height $\langle y, u_{18} \rangle > .78$. Each weak near-four has an actual mate, formed by releasing its white neighbor and retaining its cap. Positive finite Gram inversion gives an actual adjacent vertex feasible against every wall. Its edge cosine exceeds 11; for example the left mates have

$$C_j = \sqrt{2(i-1)/19}\left((20 - i)/2 + (19 - i)/\sqrt{2}\right) > 11, \qquad i = 5, 6.$$

At comparison 11/8, the normalized mate height exceeds 7/8. A left mate has correlation $< .15$ with u_{18}. The sum of these heights exceeds 1.655, whose square exceeds 2.3, while the squared norm of the sum of the directions is < 2.3. This contradicts $\|y\| < 1$. No left near-four remains.

Some $k = 5, 6$ is now strong by helper nonadjacency, with normalized height $> .677$. For a right mate $j = 14, 15$, the correlation formula

$$\rho_{ab} = \sqrt{\frac{a(19-b)}{b(19-a)}} \quad (a \leq b)$$

gives $\rho_{18,j} < .46, \rho_{18,k} < .161, \rho_{j,k} < .407$. For one vector $V = .5u_{18} + .6u_j + .4u_k$,

$$\langle y, V \rangle > .5(.78) + .6(7/8) + .4(.677) = 1.1858, \quad \|V\|^2 < .77 + .276 + .0644 + .19536 = 1.30576.$$

Equation (4.11) gives $\|y\|^2 > 1.06584$. All near-four orientations are excluded at .78. Only nonadjacent pure weak helpers remain.

Use rational heights $H_{19-i} = H_i$, linear on 0–5 and 14–19, with

$$H_5 = 1.844, \ H_6 = 1.995, \ H_7 = 2.103, \ H_8 = 2.173, \ H_9 = H_{10} = 2.207.$$

Their positive slopes are $.3688, .151, .108, .070, .034$, then zero. The squares $(H_i + .87)^2$, $5 \leq i \leq 9$, are

$$7.365796, \ 8.208225, \ 8.838729, \ 9.259849, \ 9.467929,$$

strictly below $140/19, 156/19, 168/19, 176/19, 180/19$. Thus every retained positive dual is paid by a strong floor. If 5 is weak use chord 0–6, reflect at 14, and omit each internal weak knot by its neighbor chord. After chords set $w_i = (H_i - H_{i-1})/\sqrt{2}$. The reference is zero-sum with demand $E_0 = \|w\|^2$.

Before internal holes, the exact energies for zero, one and two weak ends are .7205882, .70082285, .6810575. Internal curvatures are .043, .038, .036, .034 and reflections; even summing all eight losses gives

$$2(.043^2 + .038^2 + .036^2 + .034^2)/4 = .0028725 < .002873.$$

Nonadjacency makes the losses legitimate. End chords do not change any eligible internal curvature. Hence

$$E_0 > \begin{cases} .6781845, & 14 \text{ weak: } m = 6,\ K = 13,\ H_K = 1.995, \\ .69794985, & 14 \text{ strong: } m = 5,\ K = 14,\ H_K = 1.844. \end{cases} \tag{4.34}$$

The surviving individual anchor prices are

$$\beta_{13} \geq (7H_6/6 - H_7)/2 = .11225, \qquad \beta_{14} \geq (6H_5/5 - H_6)/2 = .1089.$$

Adjacent holes increase the anchor price. Omitted knots have price zero; no missing strong 7 or 18 floor is paid.

Define

$$G_0 = -e_{19} + \frac{1}{m} \sum_{\text{last } m} e_i,$$

with 19 included. Then

$$\|G_0\|^2 = (m-1)/m, \quad \langle w, G_0 \rangle = 0, \quad G_0 = \omega_{18}/\sqrt{2} - \omega_K/(m\sqrt{2}).$$

Use the new weight $\mu = .6$. The debit $.6/(m\sqrt{2})$ is $< .071$ for $m = 6$, $< .085$ for $m = 5$, below the separate anchor prices. The positive endpoint coefficient is paid by its actual height. Here

$$-y_{19} > .78\sqrt{18/19} > .7591,$$

since $19(.7591)^2 = 10.94842339 < 18(.78)^2 = 10.9512$. The last-block means are $H_K/(m\sqrt{2}) < .2352$ or $< .261$. The same local-y square, retaining $\langle w, G_0 \rangle = 0$, gives

$$\begin{aligned} m = 6: &\quad \|y\|^2 > .6781845 + 1.2(.7591 - .2352) - .36(5/6) = 1.0068645, \\ m = 5: &\quad \|y\|^2 > .69794985 + 1.2(.7591 - .261) - .36(4/5) = 1.00766985. \end{aligned}$$

Both contradict $\|y\| < 1$, proving the lemma with the local inventory established in Appendix B.4. □

Appendix B.4 supplies the complete local own/white inventory, isolated-cap point bounds, positive mixed weights and native-five determinant expansion used above. It uses only real geometry and comparison 11/8, with no original-B_{18} premise. The original-x energy is never added to this local-y budget.

4.9 The near/far pair and the final composition

4.9.1 Every actual near/far fork edge is short

For a near-three or far-three fork role, the complete finite-white classification in Section B.9.9 gives an actual other endpoint q with whole diagram A_{18}. The positive reverse direction at q is exactly

its normalized new endpoint dual: p has nonzero product with only that new incident root. Let C be the original edge cosine, $\beta = \sqrt{C^2-1}/C$, and $t = t_f < .876$.

Suppose $C \geq 5.6$. Then

$$E_q/E = C - \sqrt{C^2-1}\, t > 5.6 - .876(5.51) = .77324.$$

The function increases for $C \geq 5.6$. Here $5.51^2 = 30.3601 > 30.36$ and $(11/8)(.77324) > 1$, so $E/E_q < 11/8$. Also $\beta > .9839$, because $31.36(1 - .9839^2) = 1.0016631744 > 1$. The positive reverse height satisfies

$$\langle y, u_{18} \rangle = \frac{\beta - t}{1 - \beta t} > .78.$$

Its denominator is positive and its limiting gap is

$$.9839 - .876 - .78(1 - .9839 \cdot .876) = .000179192 > 0.$$

The original maximum supplies all-point comparison at q. Lemma 4.10 excludes $C \geq 5.6$. Every actual near/far fork edge therefore has $C < 5.6$.

4.9.2 The entire near/far family

Suppose the actual fork opposites are near and far:

$$c_N = Np - \gamma\Omega_f - \Omega_o/2, \qquad c_F = Fp - \delta\Omega_o - \Omega_1/2,$$

with f, o the different actual forks. Their times and exact edge cosines are

$$N^2 = 9\gamma^2 + 8\gamma + 5/4, \quad F^2 = 9\delta^2 + \delta - 1/2, \quad C_N = \frac{18\gamma + 8}{\sqrt{19}}, \quad C_F = \frac{18\delta + 1}{\sqrt{19}}.$$

The short-edge bounds and $\sqrt{19} < 4.359$ give $\gamma < .912, \delta < 1.301$. Put $U = 3\gamma + 4/3, V = 3\delta + 1/6, a^2 = 19/36$. Then $N^2 = U^2 - a^2, F^2 = V^2 - a^2$, and

$$(UV - a^2)^2 - N^2F^2 = a^2(U - V)^2 \geq 0, \qquad UV - a^2 > 0.$$

The actual mutual magnitude consequently satisfies

$$\begin{aligned} \lambda = -\langle c_N, c_F \rangle &= NF - 8\gamma\delta - \gamma/2 - (9/2)\delta - 1/4 \\ &\leq \delta(\gamma - 1/2) - 5/9 \\ &< 1.301(.412) - 5/9 = .536012 - 5/9 < 0. \end{aligned}$$

This contradicts all-wall acuteness. If $\gamma \leq 1/2$ the upper bound is already negative; the accepted own list in fact has $\gamma \geq \cos(\pi/4) > 1/2$. Zero, every finite, parallel and unbounded dotted mutual are excluded at once, for all finite or dotted near/far owns.

4.9.3 Exhausting the actual head and fork covers

For a dotted head, (4.33) removes every far-next-three role from Section B.9.9. What remains is isolated, near-three, or far-three. Their six unordered pairs are exhausted by

Actual pair	Complete excluding argument
isolated/isolated	Sections B.9.14 and B.9.15
isolated/near	Section B.9.16
isolated/far	Sections B.9.17 and B.9.18
near/near	Section B.9.9
near/far	The complete argument above
far/far	Section B.9.18

Both assignments of the genuine signed forks are included. Every pair is impossible, so no dotted head remains.

The head cover also includes own zero, finite and parallel cases. Section B.9.1 removes own $0, 3, 4$ by time constraints and own 6 by a positive affine $[6, 3]$ kernel. At own 6 an actual null is $\sqrt{3}H + 2n_1 + n_2$; at parallel own it is $H + n_1$. The remaining finite labels $5, 7, 8, 9$ and every integer label at least 10 are excluded by the complete conjunction in Section B.9.22. Every dotted own has just been excluded. Thus the whole-D_{18} diagram is impossible.

Theorem 4.8 excludes every dimension at least 19. At exact dimension 18, the fresh complete cover and its row consumers, including Section 4.5, leave only whole D_{18} at the same original farthest vertex. The head/fork argument excludes it. Hence dimension 18 is impossible, proving Theorem 4.1. Each step uses its own actual-rank cover and the same original vector within a case; no dimension heredity is used.

4.10 Finite algorithms and conditional verification

4.10.1 The mathematical finite-cover calculation

The rank covers are finite combinatorial arguments supported by real inequalities, not field enumerations. For fixed total rank n, the forest calculation is:

1. List partitions of n into ranks of spherical irreducibles. Retain all exceptional finite types and classical types at their actual ranks. Keep $I_2(m)$ as a symbolic ordinary rank-two family for unbounded m, distinguishing its exceptional actual typed profiles.

2. Mark at most one allowed $I/J/N_7/O_7$ profile by the actual-wall compatibility lemma. Apply joint bounds only to disjoint sets of components, counting each defect once.

3. Sum proved same-vector energy bounds. Discard a forest only when a strict lower bound is at least 1. If a dominant rank needs a fresh local layer, retain it until that actual-rank lemma is established.

4. Return the retained grouped necessary cover and the inequality discarding every other partition.

Termination follows from finitely many rank partitions and finite type choices at each rank. The unbounded native parameter is represented by inequalities proved over its entire domain. Completeness uses spherical classification, compact acute simplicity, complete actual-defect compatibility, and spectator-uniform component bounds. The output is a necessary cover; no realization follows. A program testing only finitely many native labels would not establish it.

Local own/white calculations have a different domain. After deleting a specified original node, classify every finite extension of the retained spherical forest by its actual opposite wall, including own zero, all finite owns, parallel and dotted owns. Finite white extensions are bounded by spherical classification. Infinite own families are closed by analytic inequalities over complete intervals. Termination of a finite classification does not justify discarding a continuous branch. Completeness must include spectator attachments, affine kernels and every original wall for every auxiliary point.

5 The compact arithmetic bound

The general compact theorem already gives $n \leq 17$. It remains to exclude seventeen and sixteen, at the same original farthest vertex used in Section 4. Arithmetic restrictions enter through the actual field and integral parameters of actual walls. All reductions use whole orthogonal components of one vector x, with $\sum_C e_C = \|x\|^2 < 1$; overlapping subdiagrams are never independent energy summands.

We include the arithmetic helper and invariant-lattice arguments, the rank-seventeen partition, a complete small-component calculation, the whole-D_{16} field reduction and all its dyadic terminal branches. Complete component inventories and row-by-row geometric dispatches remain in catalogue items A0 and A5. The four finite computations are stated with their exact domains in A1–A4; their algorithms and reproduction records remain online. The initial rank-sixteen wall inventory is explicitly identified before its field-theoretic consumer.

5.1 The actual arithmetic field

An arithmetic reflection lattice has an admissible quadratic-form model over its actual totally real field K; the structural reflection theorem is recorded as Lemma 5.2 of [13]. It supplies K-valued root lines and positive definite nonphysical forms. For roots r_i on those lines put

$$a_{ij} = 2(r_i, r_j)/(r_i, r_i).$$

Invariant cyclic products are algebraic integers and generate the actual ground field. Individual physical unit roots need not be K-valued. Disconnected roots need not have equal raw norm squareclasses.

If an actual principal Gram matrix, after its specified permitted K-rescaling, is defined over $k \subset K$ and indefinite physically, then $K = k$: an additional real embedding fixing k would preserve indefiniteness and contradict positive definiteness. A nonzero algebraic integer also has nonzero integral norm, so bounds at *every* nonphysical embedding can force a degree bound. Neither argument assumes a convenient overfield or an integer power basis.

The rational field is excluded in the dimensions used here: compactness requires anisotropy, while Meyer's theorem makes an indefinite rational form in at least five variables isotropic. These arithmetic inputs are reviewed in [14]. Finite/affine classification and acute wall geometry are external foundations; component estimates, forest reductions and computed propositions below are internal results.

For reference, the original adjacent-edge and completed-square inequalities are

$$x \cdot u \geq \tanh(L/2), \qquad \|x\|^2 \geq 2x \cdot w - \|w\|^2. \tag{5.1}$$

The first concerns a genuine edge and the second any one reference w. Every change of reference below records its individual available coefficients before these inequalities are applied.

5.2 Paying all isolated constraints together

The arithmetic component estimates use a common analytic step. It explains why overlapping local pressures can be combined without optimizing different vectors independently.

Lemma 5.1 (Simultaneous absorption)**.** *Let n_i be incident unit normals, ω_i their duals, and $w = \sum_i \beta_i \omega_i$ with $\beta_i = \langle w, n_i \rangle \geq 0$. Suppose $E_w = \|w\|^2 = \sum_i \beta_i H_i$. Outside a set J, assume $\langle x, \omega_i \rangle \geq H_i$. For $i \in J$ suppose $\beta_i > 0$, $0 < u_i < 1$, and*

$$\langle x, \omega_i - u_i n_i \rangle \geq H_i - \delta_i, \qquad 2\delta_i < \beta_i(2u_i - u_i^2).$$

Then $\|x\|^2 \geq E_w$, strictly if J is nonempty or some positively weighted retained pressure is strict.

Proof. Use the single vector $Z = w - \sum_{i\in J} \beta_i u_i n_i$. The pressures and $\langle n_i, n_j\rangle \leq 0$ for $i \neq j$ give

$$\langle x, Z\rangle \geq E_w - \sum_{i\in J} \beta_i \delta_i, \qquad \|Z\|^2 \leq E_w - \sum_{i\in J} \beta_i^2 (2u_i - u_i^2).$$

Substitution in (4.11) proves the assertion, including its strictness. In particular adjacent isolated sites are allowed. □

For an actual cap isolated within its component in the retained range $0 < u < 1$, retain its full foreign projection and write

$$c = Ap - \gamma\omega_i - v_{\mathrm{for}}, \quad Q = \|v_{\mathrm{for}}\|^2, \quad u = (1-Q)/\gamma^2 \in (0,1), \quad D_i = \|\omega_i\|^2.$$

Positive clearing gives the actual vector

$$\eta = c + v_{\mathrm{for}} + \gamma u n_i = \gamma\{\sqrt{D_i - u}\, p - (\omega_i - u n_i)\},$$
$$\eta^2 = -\gamma^2 u(1-u).$$

It is future pointing, zero against the contacted foreign block and c, negative against n_i, and nonpositive against every other wall. Lemma 4.3 therefore gives

$$\langle x, \omega_i - u n_i\rangle \geq \sqrt{D_i - u} - \sqrt{u(1-u)}.$$

The loss from $\sqrt{D_i}$ is at most $a_i u + \sqrt{u(1-u)}$, where $a_i = (\sqrt{D_i} + \sqrt{D_i - 1})^{-1}$ in the chain ranges used. Thus a reference with $H_i + \ell < \sqrt{D_i}$, positive prices $\beta_i \geq b$, and $a_i \leq a$ absorbs every such foot whenever

$$au + \sqrt{u(1-u)} - \frac{b}{2}(2u - u^2) < \ell \qquad (0 < u < 1).$$

The actual cap inventories, golden-helper exclusions and numerical reference coefficients are catalogue item A0. Their complete proofs are in Addendum A; the lemma proves the shared analytic combination used by those component estimates.

5.3 The actual golden long-end star and invariant lattices

Put $\varphi = (1+\sqrt{5})/2$. For the lattice

$$L = \mathbb{Z}[\varphi]^{n+1}, \qquad f = -\varphi t^2 + x_1^2 + \cdots + x_n^2,$$

the subgroup generated by all integral hyperplane reflections has an infinite-volume fundamental chamber for $n \geq 8$ [15]. We show below that the relevant facet reflections preserve this lattice.

Put $u = \varphi - 1$ and use the unit-equivalent form $f = -ut^2 + \sum x_i^2$. A whole D_n long-end-five star has old raw roots $-e_i + e_{i+1}$, $-e_{n-1} - e_n$, and cap $r = e_0 + \varphi e_1$, all square two. Its full Lorentzian Gram forces actual $K = \mathbb{Q}(\varphi)$: any larger field has a nonphysical embedding fixing this indefinite principal. The old group gives all even spatial sign changes and permutations. Conjugating the cap reflection gives maps R_i whose only nonidentity block, on zero and i, is

$$\begin{pmatrix} \varphi & -\varphi \\ 1 & -\varphi \end{pmatrix}.$$

A whole B_n head-five star has the same configuration and contains those even signed permutations, so the argument also applies there. Arithmeticity supplies a full invariant $\mathcal{O}_K$-lattice Λ, for example by summing finitely many translates of a lattice stable under an arithmetic finite-index subgroup.

Let $\mathcal{A}$ be the algebra generated by these lattice-preserving maps. For distinct spatial i, j, k, even sign changes d_{ij} give $(1 - d_{ij})(1 - d_{ik}) = 4E_{ii}$. Permutations give all spatial $4E_{ij}$. The displayed R_i and the unit φ then give $4E_{i0}, 4E_{0i}, 4E_{00}$; explicitly

$$\begin{aligned} 4E_{ii}R_i &= 4E_{i0} - 4\varphi E_{ii}, \\ R_i 4E_{ii} &= -4\varphi E_{0i} - 4\varphi E_{ii}, \\ R_i 4E_{i0} &= -4\varphi E_{00} - 4\varphi E_{i0}. \end{aligned}$$

The golden ring is a PID, for example by its Minkowski bound $\sqrt{5}/2 < 2$. Scale the coordinate-content ideal of Λ to the unit ideal. The matrix units imply $4L \subset \Lambda \subset L$. The fourfold sandwich is essential for odd n.

Modulo two, $\mathcal{O}_K/2 = \mathbb{F}_4$. Put

$$\ell(t, x) = \varphi t + \sum x_i, \qquad H = \ker \ell, \qquad v = (\varphi^2, 1, \ldots, 1).$$

The only invariant subspaces are $0, \langle v \rangle, H, L/2L$. Indeed any unequal spatial coordinates give $e_i + e_j$ by a transposition difference; their orbit spans the spatial augmentation hyperplane, and R_i supplies a time direction, giving H. If every spatial coordinate is equal, R_i-invariance forces the line $\langle v \rangle$. Here $v \in H$ exactly for odd n.

For the two residue flags

$$W_1 = (\Lambda + 2L)/2L, \qquad W_2 = ((\Lambda \cap 2L)/2)/2L, \qquad W_1 \subset W_2,$$

the content normalization makes $W_1 \neq 0$. If W_1 is full, $\Lambda = L$. If $W_1 = H$, differences under even sign changes of a lift of $e_i + e_k$, divided by two, put a coordinate vector in W_2, forcing it full. If $W_1 = \langle v \rangle$, the same differences force $H \subset W_2$; for even n, W_2 is full, while for odd n it is H or full. Thus, writing E for the inverse image of H, the complete lattice list is

$$L, \quad E, \quad \mathcal{O}_K v + 2L, \quad \mathcal{O}_K(v + 2z) + 2E \quad (n \text{ odd}, \ z \in L).$$

This includes every lift, not merely every residue subspace.

Intrinsically,

$$E = \{x \in L : f(x) \in 2\mathcal{O}_K\}, \qquad E^* = L + \mathcal{O}_K(v/2), \qquad f(v) = n - 1 - 2\varphi.$$

The latter norm has dyadic valuation zero for even n, exactly one for odd n. Writing $x = y + av/2$ shows

$$\{x \in E^* : f(x) \in \mathcal{O}_K\} = L :$$

if a is not divisible by two, the final norm term has negative valuation and the other terms are integral. Hence every isometry of E, and by duality of $\mathcal{O}_K v + 2L = 2E^*$, preserves L. For a deep odd lattice put $\lambda = v + 2z$; its norm still has valuation one. Writing $x = a\lambda/2 + e$, $e \in E$, gives

$$\{x \in \Lambda/2 : f(x) \in \mathcal{O}_K\} = E.$$

Its isometries therefore preserve E and then L. Every invariant lattice in the complete list has $O(f, \Lambda) \subset O(f, L)$. Thus the actual reflection group lies inside the literal integral group. Its arrangement chamber refines the assumed compact chamber, contradicting the external literal no-compact-chamber cutoff (or the stronger infinite-covolume statement). This excludes the actual whole-B/D isolated long-end-five configuration, with every compatible invariant lattice retained.

5.4 The complete rank-seventeen reduction

Write $T_2 = I_2(m)$, $m \geq 3$, for an ordinary irreducible dihedral; $T_3 = A_3, B_3, H_3$ and $T_4 = A_4, B_4, D_4, F_4, H_4$ unless a stated field restriction removes a type. Put $\varphi = (1+\sqrt{5})/2$, and write $K_7 = \mathbb{Q}(2\cos(2\pi/7))$ and $K_9 = \mathbb{Q}(2\cos(2\pi/9))$. Ordinary nonshort A_1 costs $> 1/10$, ordinary $T_2 > 1/7$, $T_3 > 1/4$, $T_4 > 2/7$. A total-rank-three forest need not cost $> 1/4$; its ordinary bound is $> 17/70$.

The exceptions retain actual-wall meanings. Type I is an isolated short rank-one opposite

$$c_I = \sqrt{h}\, p - \sqrt{1+h}\, e, \qquad 0 < h < 1/2.$$

Type J couples two rank-one directions with joint energy $> .34$. The typed rank-two defects are N_7, old A_2/new isolated-seven, and O_7, old $I_2(7)$/new isolated-three. Each costs $> .09$; they cannot coexist with one another or the short I/J alternatives. A bare old seven is not a typed defect. These are the complete small-component alternatives of Proposition 5.6. Their geometric compatibility proof uses the native gap between distinct actual walls, not merely the abstract old forest.

5.4.1 Shared costs and arithmetic prefix seals

We specify the component inputs before carrying out either rank count. The ordinary geometric rates are the component theorems of Section 4 and catalogue C1, applied to whole orthogonal components at the same original vertex:

A_1	$1/10$	T_2	$1/7$
A_2	$319/2000$	A_3	$1/4$
B_3, H_3	$2/7$	A_4	$2/7$
H_4	$11/36$	B_4, D_4, F_4	$1/3$
A_5	$3/8$	D_5	$5/14$
B_5	$3/7$	A_6, B_6, D_6	$3/7$
B_7, D_7	$1/2$	B_8, D_8	$4/7$
E_6	$9/20$	E_7	$5/9$
E_8	$2/3$		

The entries are strict lower bounds from catalogue C1; their complete component proofs remain in Addendum C. In particular every ordinary component outside A_r, $r \geq 7$, and B_r, D_r, $r \geq 9$, costs more than its rank divided by fourteen. The universal large-component rates used here are $A_r > 1/2$ for $r \geq 7$, $A_r > 4/7$ for $r \geq 8$, $A_r > 5/7$ for $r \geq 13$, and $B_r, D_r > 9/14$ for $r \geq 9$. These are geometric companion inputs, not assertions that a higher-rank whole-diagram exclusion holds at a smaller rank. The stronger B_3, H_3 rate is needed only for the rank sixteen count; the rank seventeen count uses $1/4$.

The actual short-I wall gives parameters u, t with

$$e_I \geq u^2, \quad 0 < u < 1/3, \quad (1-u^2)(1-t^2) = 8ut, \quad t + 3u/2 > .767. \tag{5.2}$$

Every normalized old dual direction outside I has projection at least t. Its squared-time parameter h also gives the following geometric alternative: at $h \leq 1/5$ all these squared projections are at least $1/5$; at $h > 1/5$, $e_I > 3/70$. These are the compact proof's actual opposite-wall barrier, with no arithmetic hypothesis. We record the needed joint consequences here to make their accounting explicit.

Lemma 5.2. *For ordinary irreducible T_j the following bounds hold, each counting I once:*

$$e_{I+T_2} > .21, \qquad e_{I+T_3} > 2/7, \qquad e_{I+T_4} > 23/70, \qquad e_{I+A_1} > .20204, \qquad e_{I+2A_1} > .29.$$

Proof. For T_2 use $e_{T_2} > 1/7$ and $e_{T_2} \geq t^2$. If $u \geq 519/2000$, then $u^2 + 1/7 > .21$. Below that point $u^2 + (.767 - 1.5u)^2$ is decreasing and has endpoint value $.2100353125 > .21$.

For T_3, its two endpoint correlations are $1/3$, $1/\sqrt{3}$ and ρ with $\rho^2 = (3\varphi + 2)/(3\varphi + 6) < 16/25$, respectively. Cauchy on the sum of the two unit directions gives $e_{T_3} > (10/9)t^2$, as well as $e_{T_3} > 1/4$. At $u \geq .19$ use $u^2 + 1/4 > 2/7$; below it the decreasing quadratic $u^2 + (10/9)(.767 - 1.5u)^2$ is at least $132407/450000 > 2/7$.

For A_4, the low-h direction cover costs more than $1/3$; the high-h case gives $3/70 + 2/7 = 23/70$, strictly exceeded. The B_4, D_4, F_4 rates already exceed $1/3$. For H_4 we need $t+2u > 6/7$. Substituting $6/7 - 2u$ for t in the left-minus-right side of the identity in (5.2), and multiplying by 49, gives

$$13 - 168u + 575u^2 - 168u^3 + 196u^4.$$

With $v = u - .15$ this is $.269725 - 4.194v + 525.86v^2 - 50.4v^3 + 196v^4$. For $-3/20 < v < 11/60$ it is at least $.269725 - 4.194v + 516.62v^2 > 0$, the last quadratic having negative discriminant. The original identity decreases in t, proving the barrier. At $u \geq 19/125$, $u^2 + 11/36 > 23/70$; below it the decreasing quadratic $u^2 + (6/7 - 2u)^2$ is at least $50389/153125 > 23/70$.

For $I + A_1$, put $f(s) = (\sqrt{1+s} - 1)/(\sqrt{1+s} + 1)$. The actual edge barrier gives half-tangent squares $q = f(h)$ and at least $r = f(1/(4h))$ on the other component. Writing $P = \sqrt{qr}$, inversion gives $q + r = 1 + P^2 - 8P$, while AM–GM gives $q + r \geq 2P$. As $0 < P < 1$, this implies $P \leq q_0 = 5 - 2\sqrt{6}$ and

$$q + r - 2q_0 = (q_0 - P)(8 - P - q_0) \geq 0.$$

Thus $e_I + e_{A_1} \geq 2q_0 > .20204$; the rational seal is $2.44949^2 > 6$. Finally, for two ordinary A_1 components, at $u \geq .3$ use $u^2 + .2 \geq .29$, strictly exceeded. Below it use $q(u) = u^2 + 2(.767 - 1.5u)^2$; then

$$q(u) - q(.3) = (.3 - u)(2.952 - 5.5u) \geq 0, \qquad q(.3) = .290978 > .29.$$

Each calculation estimates one joint sum of orthogonal whole-component energies; no endpoint has been charged a second copy of I. □

Lemma 5.3 (Propagation from the actual old-seven defect)**.** *Suppose a whole $I_2(7)$ component has an actual opposite supported only there, with a native-three contact at one node and zero contact at the other. Every other whole rank-one component has energy $> 2/15$, and every other whole irreducible rank-two component has energy $> 1/5$. This lemma is geometric and allows arbitrary finite spectators.*

Proof. Put $d_7 = \csc^2(\pi/7)$. The typed wall is $c_D = \sqrt{h_D}p - v_D$, with $h_D = d_7/4 - 1 \in (0, 1/3)$. Let $c = \sqrt{h}\,p - v$ close an actual edge of another component. Its finite endpoint retains the whole old $I_2(7)$. A positive contact there contributes at least $d_7/4 > 1$ to the white Schur cost, impossible. Thus v has zero projection to that whole component, and c is distinct from c_D. The native actual-wall gap gives

$$(c_D, c) = -\sqrt{h_D h} \leq -1/2, \qquad h > 3/4. \tag{5.3}$$

This proves the height bound for every opposite outside the defect, without assuming an isolated profile for those other walls. In a rank-one direction its spatial derivative is at most $\sqrt{1+h}$, so $\sinh^2 L \geq h > 3/4$. Hence its half-tangent square is greater than $(11 - 4\sqrt{7})/3 > 2/15$.

Suppose a whole $T = I_2(m)$, $m \geq 3$, instead has energy $e \leq 1/5$. Its normalized duals have correlation $k = \cos(\pi/m)$. Write $s_i = x_T \cdot u_i$ and $F(s) = 2s/(1+s^2)$. For $m \geq 4$, distinct actual opposites would require

$$k \leq F(s_1)F(s_2) \leq F(\sqrt{1/5})^2 = 5/9 < 1/\sqrt{2} \leq k,$$

impossible. For $m = 3$, Cauchy gives $(s_1+s_2)^2 \leq 3/5$, and concavity of $\log F$ gives the incompatible bound $1/2 \leq F(s_1)F(s_2) \leq 240/529 < 1/2$. Therefore one actual wall closes both directions. Both lengths have

$$\cosh L_i \leq (1+1/5)/(1-1/5) = 3/2. \tag{5.4}$$

Let $b, g \geq 0$ be its contacts in T and Q its entire foreign Schur cost. With $d = (1-k^2)^{-1}$, the finite endpoints give

$$Q + b^2 < 1, \quad Q + g^2 < 1, \qquad h = d(b^2 + 2kbg + g^2) + Q - 1 > 3/4,$$
$$C_b^2 = \frac{d(b+kg)^2}{1-Q-g^2}, \qquad C_g^2 = \frac{d(g+kb)^2}{1-Q-b^2}.$$

Every positive b, g is consequently a finite native cosine. At least one is positive. A positive whole foreign contribution costs at least $1/4$, so $Q = 0$ or $Q \geq 1/4$.

For a single contact, say $b = \beta > 0, g = 0$, $C_g^2 = 1 + h/(1-Q-\beta^2)$. Equations (5.3)–(5.4) imply $Q + \beta^2 < 2/5$; hence $Q = 0$ and the native gap forces $\beta = 1/2$. Then $h = d/4 - 1 > 3/4$ gives $d > 7$ and $m \geq 9$, since $d_8 = 4 + 2\sqrt{2} < 7$. But $\sin t < t$ and $\pi^2 < 10$ give $d > 81/10 > 31/4$, so $C_g^2 = (d-1)/3 > 9/4$, a contradiction.

If both contacts are positive and $Q > 0$, let g be the larger. When $g \geq 1/\sqrt{2}$, $C_b^2 \geq (4/3)(1 + 1/\sqrt{2})^2 > 9/4$. Otherwise $b = g = 1/2$. At old label three, $h = Q < 3/4$; at old label at least four, $C_b^2 \geq (1+k)/(2(1-k)) > 5/2$. Both are impossible.

It remains $Q = 0$. If the larger contact has label at least five, then $g > 4/5$, $b \geq 1/2$, $k \geq 1/2$, $d \geq 4/3$ give $C_b^2 > 3$. If both contacts have label four, $C_b^2 = (1+k)/(1-k) \geq 3$. The remaining finite contact pairs are $(3,4)$ and $(3,3)$. For $(3,4)$, old label at least four gives $C_b \geq 2$; old label three gives $h = \sqrt{2}/3 < 3/4$. For $(3,3)$, old label at least five gives $C_b^2 = (1+k)/(3(1-k)) > 3$; old labels three and four give $h = 0, 1/\sqrt{2} < 3/4$. Every contact and foreign case contradicts (5.3) or (5.4). Thus $e_T > 1/5$, completing both conclusions. □

Both typed defects force the actual field K_7. Their three actual roots form the indefinite $(7,3,2)$ principal. Odd bonds permit a square-two normalization over K_7; an extra embedding fixing K_7 would preserve that indefinite principal. Hence actual-field descent gives $K = K_7$. In particular H_3, H_4 cannot occur beside a typed defect, since their five-bond would put $\sqrt{5}$ in K_7. For N_7 every other normalized dual has squared projection $> 1/5$. For O_7, Lemma 5.3 gives every other A_1 cost $> 2/15$ and every other ordinary irreducible T_2 cost $> 1/5$. The hypothesis is the actual isolated-three wall, not the old label seven alone.

Lemma 5.4 (Literal arithmetic helpers)**.** *At a classified long-prefix index $i \geq 6$ of B_r or D_r, an actual far helper $v = \gamma\omega_i + \omega_1/2$ with $202/125 < 2\gamma < 12/5$ is impossible. At a D fork-helper site with $r \geq 9$, the corresponding profile $v = \gamma\omega_i + \omega_f/2$ can only have $2\gamma = \varphi$.*

Proof. Normalize the connected long roots to square two over K. The simple-three helper contact permits the same normalization of the actual outside root; its path cycle makes $t = 2\gamma$ integral. The short B root has square one, so the entire old Gram is rational. All embeddings of $\mathbb{Q}(t)$ extend to K, and every nonphysical one must give a strictly positive Schur complement.

For the far helper the raw inverse entries $i, 1, 1$ give $it_\sigma^2 + 2t_\sigma - 1 < 0$. At $i \geq 6$ this forces $|t_\sigma| < .61$ and $t_\sigma < .28$. Degree at least three would give a nonzero integral norm of magnitude less than $(12/5)(.61)^2 < 1$. In degree two the integral trace is strictly between $503/500$ and $67/25$, hence two; its norm $t(2-t)$ lies strictly between -1 and one, again impossible. Degree one leaves $t = 2$, an actual parallel pair with a future feasible null sum, contrary to compactness.

For the fork helper the inverse entries $i, i/2, r/4$ give

$$4it_\sigma(t_\sigma + 1) + r < 8, \qquad -1 < t_\sigma < 0.$$

The integral nonzero element $t(t+1)$ has physical value $< 204/25$ and other conjugates of magnitude at most $1/4$. Its norm excludes degree at least three. In degree two the trace of t is one or two. Trace one gives $t < 2$ and norm in $(-2, -.995456)$, hence norm -1 and $t = \varphi$. Trace two gives norm in $(-24/25, 0)$, impossible. The rational $t = 2$ is again parallel. This proves both assertions without identifying $\mathbb{Q}(t)$ with the full field in advance. □

Lemma 5.5 (The mixed B_2 tail)**.** *At a B_r site $i = r - 2 \geq 12$, the actual own-three/short-three profile $v = (\omega_i + \omega_r)/2$ is impossible.*

Proof. Normalize the old long roots to square two and the short root to square one over K. The simple-three short contact normalizes the outside root to square one; its own-three product is $-1/\sqrt{2}$. Thus $\sqrt{2} \in K$. At an embedding reversing $\sqrt{2}$, while fixing the rational old Gram, its Schur cost is

$$i/2 - i/\sqrt{2} + r/4 = i(3 - 2\sqrt{2})/4 + 1/2 > 1.$$

At $i = 12$ the last inequality follows from $\sqrt{2} < 17/12$, and it increases with i. This contradicts strict positivity. □

For later prefix counts put

$$f_j = \sqrt{j} - .61/\sqrt{2}, \qquad E_J = f_5^2/5 + \sum_{j=5}^{J-1} (\sqrt{j+1} - \sqrt{j})^2.$$

The generic geometric inventory in the compact proof applies to B_r with $M = r - 5 \geq 6$ and to D_r with $M = r - 6 \geq 6$, through $M + 1$. Every weak index is the literal far helper just excluded. Hence $6, \ldots, J$ are strong, with $J = r - 4$ in B and $J = r - 5$ in D. If five is strong, positive concave interpolation gives E_J. If it is weak, its actual point gives $x_1 > .5965$, $S_6 > 2.01815$; interpolation through $1, 6, J$ gives

$$L_J = .7600299945 + \frac{(\sqrt{J} - \sqrt{6})^2}{J - 6} > E_J,$$

with the quotient zero at $J = 6$. This follows from the variance comparison: $E_6 < .7$, and at most ten increments in $[1/8, 1/5]$ have variance sum at most $9/640$, less than the initial gain $> .06$. The rational lower seals used below are

J	6	7	8	9	10
$E_J >$	.69685	.73531	.76864	.79805	.82436
J	11	12	13	14	15
$E_J >$	.84816	.86989	.88989	.9084	.92564.

(5.5)

These statements retain the precise rank hypotheses of the compact proof inventory; the smaller terminal ranks are handled below, rather than being silently inserted into its generic range.

Proposition 5.6 (Necessary cover in rank seventeen)**.** *The whole forest at the original farthest vertex of a compact arithmetic* 17*-polytope is one of Table 1.*

The proof below starts at rank seventeen. The arithmetic baseline component estimates used in its partition stage are $e_{A_{15}} > .79$, $e_{A_{16}} > .83$, $e_{A_{17}} > 6/7$, and, under either actual typed defect, $e_{A_{12}} > .703101$, $e_{A_{14}} > .7897732$. The subsection "The component inputs used before the partition count" in Addendum A (catalogue A0/A5) derives these rates from stronger whole-component proofs, with arbitrary spectators. They are not higher-rank exclusions.

Group	Component	Complete spectators
1	A_{11}	$I + (A_5$ or $D_5)$
2	A_{11}	N_7/O_7 plus A_4, B_4, D_4, F_4
3	A_{12}	$A_5, D_5, (A_4$ or $H_4) + A_1, T_3 + T_2, 2T_2 + A_1$
4	A_{12}	$I + T_4,\ I + T_3 + A_1,\ I + 2T_2$
5	A_{14}	$T_3,\ T_2 + A_1,\ I + T_2$
6	A_{15}	$T_2,\ 2A_1,\ I + A_1,\ N_7/O_7$
7	A_{16}	ordinary A_1 or actual I
8	A_{17}	none
9	B_{15}	N_7/O_7
10	B_{16}	actual I
11	B_{17}	none
12	D_{15}	N_7/O_7
13	D_{16}	ordinary A_1 or actual I
14	D_{17}	none

Table 1: The remaining spherical forests. Unmarked components are ordinary.

Proof of Proposition 5.6. Call A_r, $r \geq 7$, or B_r, D_r, $r \geq 9$, dominant. There is exactly one dominant. With none, discarding only the one possible I leaves cost $> 16/14$; a typed defect gives $> .09 + 15/14$. Ordinary or J accounting is stronger. Two dominants each cost more than $1/2$. The complete ordinary table includes E, F, H, so this step loses no exceptional irreducible component.

An A_7 leaves ten spectator ranks and is impossible by the same budget. At A_8, A_9, A_{10}, even discarding I leaves at least six ordinary ranks, giving $4/7 + 6/14 = 1$, strictly exceeded. The typed and J alternatives are stronger. Thus $11 \leq r \leq 17$.

For A_{11}, ordinary spectator rank six already costs $> 3/7$. With I, the complete ordinary rank-five partitions are 5, $4+1$, $3+2$, $3+1+1$, $2+2+1$, $2+1+1+1$, 1^5. Only $A_5, D_5, (A_4$ or $H_4) + A_1, T_3 + T_2, 2T_2 + A_1$ can initially fit below $3/7$. Lemma 5.2 removes the last three by $23/70 + .1 = 3/7$, $2/7 + 1/7 = 3/7$, and $.21 + 1/7 + .1 > 3/7$. Hence group 1 is complete. With a typed defect, ordinary rank four initially leaves T_4 or $2T_2$; the other partitions cost at least .35, $12/35$ or .4 before the defect is added. Propagation makes each $T_2 > .2$, removing $2T_2$; actual-field descent removes H_4. This gives group 2.

For A_{12}, the complete cheap ordinary rank-five list is exactly that just displayed. Indeed $B_5 > 3/7$, $B_4, D_4, F_4 + A_1 > 13/30$, $T_3 + 2A_1 > .45$, $T_2 + 3A_1 > 31/70$, and $5A_1 > .5$. A J pair plus three remaining ranks costs $> .34 + 3/14 > 3/7$. These give group 3. With I, the rank-four partitions are $T_4, T_3 + A_1, 2T_2, T_2 + 2A_1, 4A_1$. The last two cost $> .29 + 1/7 > 3/7$ and $> .29 + .2 > 3/7$, respectively, using $I + 2A_1$ only once. Thus group 4 is complete. A typed defect would leave ordinary rank three and give $.703101 + .09 + 17/70 > 1$, so none remains here.

For $A_{13} > 5/7$, ordinary or J rank four already costs $> 2/7$. With I, the remaining rank-three partitions are $T_3, T_2 + A_1, 3A_1$; their joint costs exceed $2/7$ by Lemma 5.2 or by the ordinary three-A_1

cost. A typed defect plus T_2 costs $> .09 + .2 > 2/7$; two ordinary A_1 cost at least $4/15$ and give an even stronger contradiction. No A_{13} row remains.

At $A_{14} > 5/7$, ordinary spectator rank three leaves T_3 or $T_2 + A_1$; three A_1 and $J + A_1$ are too expensive. With I, only $I + T_2$ remains since $I + 2A_1 > .29 > 2/7$. A typed defect gives $.7897732 + .09 + 2/15 > 1$. These are exactly group 5. At $A_{15} > .79$, rank two leaves $T_2, 2A_1, I + A_1$, or either typed defect. A J pair closes; the other bounds do not delete the retained alternatives. At $A_{16} > .83$, its single rank can be ordinary A_1 or actual I. At A_{17} there are no spectators. This proves groups 6–8 without an endpoint or field-degree inference.

For B, D, the $9/14$ rate and spectator accounting give $r \geq 12$: at six spectator ranks even the I debit leaves $5/14$; a typed defect gives $4/14 + .09$. The generic arithmetic prefix has last strong knot $J = r - 4$ in B, $r - 5$ in D, within the rank domains specified above. The weakest floors at $r = 12, 13, 14$ are therefore E_7, E_8, E_9. At twelve their possible rank-five remainders cost more than $23/70$, $.09 + 17/70$, or $5/14$, respectively. At thirteen all rank-four remainders cost $> 2/7$, or at least $.09 + .2$ in a typed case. At fourteen the cheapest rank-three possibilities are ordinary $> 17/70$, $I + T_2 > .21$, $I + 2A_1 > .29$, and typed $> .09 + 2/15$. Each closes with its stated prefix floor. Thus no B, D component of ranks twelve through fourteen remains.

We must extend the short tails at their actual ranks before counting the proper higher rows. For $14 \leq r \leq 17$, the B_3 finite list has only the extra short-three F_4 role; it is sealed against foreign contacts and is affine or directly strong. At the B_2 site $i = r - 2$, the extra own-three/short-three profile is excluded by Lemma 5.5. Own zero, four, six and parallel retain their affine or null principals; native own at least five is directly strong. A bridge through the short endpoint is a full B_r with internal new root and admits no finite foreign arm. Its residual is $1/(2i)$ and its ratio at least $1/\sqrt{2i} + \sqrt{i}/2 > 100/61$ for $i \geq 11$. All the other classified weak roles are excluded by Lemma 5.4. Hence $6, \ldots, r - 2$ are strong, so $e_{B_r} > E_{r-2}$ in either initial-five branch. At B_{15}, $E_{13} > .88989$ excludes ordinary $T_2, 2A_1, J, I + A_1$, leaving either typed defect. At B_{16}, $E_{14} > .9084$ excludes ordinary A_1 and leaves actual I. Whole B_{17} is retained.

For D_{15} the spectator rank is exactly two, and every allowed remainder costs $> .09$. Its bulk already gives strong $6, \ldots, 10$. The local D_4, D_3 lists at indices eleven and twelve retain only an additional own-five/fork-three case. More explicitly, own zero, three, four, six and parallel have the same affine or null principals; middle-three, end-four and the sealed H_4 end-five are directly strong. The full bridge is D_{15}; at D_3 the end-three bridge is A_{15} with both arms at least three, so no finite foreign arm is possible. Its residual $1/8 + 1/(2i) \leq 1/6$ and numerator $> .75$ make it strong; its zero-own cycle is affine. The remaining fork helper is golden by Lemma 5.4.

For that golden role the actual positive left H_4 projection has

$$\alpha^2 = \varphi + 5/8 > 56/25, \quad \delta = (5 - 3\varphi)/8, \quad \sqrt{\delta}/\alpha < .091, \quad A^2 = i\alpha^2 - \kappa,$$

where $\kappa = 1/2$ at D_4, $5/8$ at D_3. The positive correction is $\gamma(3n_i/2 + n_{i-1} + n_{i-2}/2)$. The finite projection is future and feasible against all original walls; the correction alone is not used as a hyperbolic point. Its displacement divided by α is flat one on the first $i - 3$ coordinates, has square $< i$, and gives pressure $G > \sqrt{i - \kappa/\alpha^2} - .091$. Pay weight $1/6$ at the previous strong knot $K = i - 3$. Both initial-five references have endpoint coefficient $> 1/6$, so a single pure-D square gives

$$e_D > E_K + (G - f_K)/3 - (i - K)/36.$$

The complete two-row computation is

	i	K	$G >$	$f_K <$	$e_D >$
D_4	11	8	3.19	2.398	.9493
D_3	12	9	3.33	2.569	.9683.

The radical seals are $3.281^2 < 11 - 25/112$ and $3.421^2 < 12 - 125/448$. The pure energies each exceed .91; adding only the disjoint actual spectator energy $> .09$ excludes both roles. Thus $6, \dots, 12$ are strong and $e_{D_{15}} > E_{12} > .86989$. Ordinary $T_2, 2A_1, J, I + A_1$ close; either typed defect is retained. At D_{16} its ordinary A_1 and actual I are both retained, and whole D_{17} is retained. These are groups 12–14.

Every partition, actual defect and permitted dominant rank has now been considered. No root has been deleted from another ambient list, and every sum concerns the same original vector. □

Proposition 5.7 (The other rows in rank seventeen)**.** *Every row of Table 1 except group 8 is impossible.*

Proof. The complete "Every rank-seventeen row has a complete consumer" dispatch in Addendum A (catalogue A5) excludes groups 1–7 and 9–14. Each entry refers to its printed actual-wall, field and individual-payment proof. The shared algebraic census for whole B17 and D17 has separate actual-domain inclusions and separate original-vector consumers. Group8 is treated next. □

Proposition 5.8 (Actual chain endpoints)**.** *If group 8 occurs, both actual endpoint opposites are isolated within A_{17}. Their integral parameters U, V generate the same actual field and satisfy*

$$36/17 < U, V < 3969/425, \qquad 0 < U^\sigma, V^\sigma < 36/17 \quad (\sigma \neq \mathrm{id}). \tag{5.6}$$

The field is nongolden, has degree at most five, and at least one endpoint is finite. In fact

$$K \in \{\mathbb{Q}(\sqrt{2}), \mathbb{Q}(\sqrt{3}), K_7, K_9\};$$

in the cubic fields a finite endpoint has label seven or eighteen, respectively.

The whole-A_{17} proof in Addendum A (catalogue A5) gives the complete endpoint inventory, golden-end cases, degree bound and finite-anchor reduction, before the algebraic computations are applied. It retains every finite-white and own case, including the golden time-compatible exception. One decisive step explains the simultaneous accounting. The common body has squared norm .9037838. Both dotted cores are paid at weight .3; the outer coefficients change from .1382 to .0782 and still fund every central foot. On the entire $0 < u < 1$ interval the extra contribution is

$$.6\{\sqrt{17/9 - u} - \sqrt{u(1-u)} - .4174\} - .09(8/5 - 2u + u^2) > .04969.$$

The complete circle bounds in that finite-anchor proof give $.9037838 + 2(.04969) > 1$ if both ends are dotted. No unavailable strong-five floor is used. The complete finite alphabet, including label30 before its conjugate test, then gives the stated fields, rather than a prescribed field list.

In raw old-root normalization of square two, a cap has square $2U$ and time square $F(U)/18$, where $F(X) = X(17X - 36)$. The inverse endpoint entry is $1/18$. Every actual pair consequently satisfies

$$R = F(U)F(V) \in K^2, \qquad g = (UV - \sqrt{R})/18 \in \mathcal{O}_K, \qquad L = g^2/(UV) \in \mathcal{O}_K. \tag{5.7}$$

The root is physically positive. Acuteness gives $g \leq 0$. If $g \neq 0$, nonphysical positive definiteness gives $0 < L^\sigma < 4$. The branch $g = L = 0$ is retained. Physical owns and mutual parameters must be native or dotted; four would be a compactness-forbidden parallel pair. The endpoints need not be equal.

5.5 The four finite algebraic propositions

The computations check necessary algebraic domains. An empty domain excludes an actual configuration after geometric inclusion has been proved. A nonempty domain supplies necessary pairs, followed by a separate geometric contradiction, not examples.

5.5.1 Full quadratic integer rings

For squarefree $d > 1$ use $\omega = (1 + \sqrt{d})/2$ if $d \equiv 1 \pmod 4$, otherwise $\omega = \sqrt{d}$. Let Δ be the fundamental discriminant. If $Y = a + b\omega$ generates K, $Y > Y' > 0$, $Y < B$, then

$$0 < b\sqrt{\Delta} = Y - Y' < B.$$

Thus $\Delta < B^2$, and b and the strip-permitted integers a have explicit finite bounds. Squarefree trial division and these integer intervals cover the full maximal order.

Use rational pairs $a + b\sqrt{d}$. Integrality in the half basis means $2b, a - b \in \mathbb{Z}$; otherwise both coefficients are integral. Signs use exact rational square comparisons. For $b \neq 0$, a field square requires norm n^2 with rational $n \geq 0$ and one of

$$x^2 = (a + n)/2, \quad x^2 = (a - n)/2$$

a nonzero rational square. Set $y = b/(2x)$ and verify the full square identity. Both signed norm branches are necessary and sufficient. For $b = 0$ test rational squares and d times rational squares, including zero. A square norm alone is insufficient.

The complete positive quadratic finite native alphabet is

$$1, 2, 3, (3 + \sqrt{5})/2, (5 + \sqrt{5})/2, 2 + \sqrt{2}, 2 + \sqrt{3}.$$

Indeed $\phi(m) \leq 4$ leaves $m = 3, 4, 5, 6, 8, 10, 12$. Zero is orthogonality, values above four are dotted, and each radical value is retained only in its actual field.

Proposition 5.9 (Recorded finite outputs)**.** *At the indicated complete domains the exact outputs are as follows.*

1. *The quadratic A_{17} domain* (5.6)–(5.7) *has no survivor. There are 25 discriminants $\Delta < 88$, 53 models and 159 ordered pairs: 106 fail field-square membership, 52 long-cycle integrality and one two-cycle integrality.*

2. *The two cubic A_{17} finite-anchor domains have nine strip models, all failing a necessary test. Their full integer rings are $\mathbb{Z}[U]$, where U is the largest root of $X^3 - 5X^2 + 6X - 1$ or $X^3 - 6X^2 + 9X - 3$, respectively. The complete coefficient boxes for $V = a + bU + cU^2$ are $[-4, 4] \times [-9, 9] \times [-3, 3]$ and $[-6, 6] \times [-9, 9] \times [-2, 2]$. The strips and cycles are those of the preceding actual endpoint reduction, with the full native alphabet at the physical embedding.*

3. *The quadratic domain*

$$4 < U < 11,\ 0 < U' < 2, \quad 2 < W < 676/17,\ 0 < W' < 4/17,$$

$K = \mathbb{Q}(U) = \mathbb{Q}(W)$,

$$g = UW - \sqrt{U(U-2)W(17W-4)}, \quad g,\ L = g^2/(2UW) \in \mathcal{O}_K, \quad 0 < L' < 4,$$

with physical native/dotted mutual parameter, has 36 fields $\Delta < 121$ and 120 pairs. Its only survivors are

K	U	W
$\mathbb{Q}(\sqrt{13})$	$(5+\sqrt{13})/2$	$11+3\sqrt{13}$
$\mathbb{Q}(\sqrt{17})$	$(7+\sqrt{17})/2$	$(21+5\sqrt{17})/2$.

4. The quadratic whole-B_{16} domain

$$2 < U < 12,\ 0 < U' < 2, \quad 4 < V < 64,\ 0 < V' < 1/4, \quad UV/2 \in \mathcal{O}_K,$$

$$k^2 = \frac{(U-2)V}{4U(4V-1)}, \quad k > 0, \quad G = 4Uk(4V-1) - UV, \quad L = G^2/(2UV),$$

with $k \in K$, integral G, L, actual acuteness and the following exact predicates, has 43 fields $\Delta < 144$: $G \geq 0$; if $G = 0$ then $L = L' = 0$, whereas otherwise $0 < L' < 4$; and U and every nonzero L must belong to the native finite alphabet of their actual field or be greater than four. In particular $L = 4$ is excluded, and no continuous nonnative value below four is allowed. The identity

$$L + G = ((15U - 32)/2)V - 2U + 4$$

is checked for the physical positive square root k. The complete domain has 100 head models, 85 short models and 273 pairs. Exactly 168 fail half-product integrality and 104 field-square membership. The sole survivor is $U = 5 + 2\sqrt{6}$, $V = 10 + 4\sqrt{6}$.

These outputs are catalogue items A1–A4. The complete ring arguments, coefficient bounds, exact sign and field-square tests, termination proofs and computation records are given there. In particular both rational norm branches are retained; a square norm alone is never sufficient. The actual inclusions for whole B_{17} and D_{17} are distinct, even though their last algebraic domains coincide. Their two surviving pairs, and the one B_{16} pair, are excluded by the separate geometric consumers in catalogue A5. The rank-seventeen dispatch and the empty quadratic and cubic A_{17} domains therefore exclude seventeen.

5.6 The complete rank-sixteen reduction

The rank-sixteen classification in Proposition 5.10 starts with all finite components and actual defect compatibility. Two dominant components each cost $> 1/2$. With none, the largest allowed rank-one debit still leaves ordinary cost $> 15/14$; the typed case costs $> .09 + 14/14$. The unique A dominant has rank at least ten and the B/D dominant rank at least eleven. Their small classical tails are checked at their actual ranks, including the direct D_{11} boundary. The D_{14} weak-tail payments give pure energies $> .92531$ or $> .94697$ before the disjoint spectator cost $> .09$ is added. Only then are the stronger tail floors used.

The conditioned $A_{11} > .652761$ component input used below is proved independently of either forest cover in the conditioned-A_{11} corollary in Addendum A (catalogue A5). Its proof reconstructs the no-helper reference directly; the weaker uniform 13/20 theorem would not imply that value.

Proposition 5.10 (Necessary cover in rank sixteen)**.** *The whole forest at the selected original farthest vertex is one of Table 2.*

Table 2: Necessary groups in rank sixteen.

Group	Component	Complete spectators
1	A_{10}	$I + (A_5$ or $D_5)$
2	A_{10}	N_7/O_7 plus A_4, B_4, D_4, F_4
3	A_{11}	$A_5, D_5, (A_4$ or $H_4) + A_1, A_3 + T_2, 2T_2 + A_1$
4	A_{11}	$I + T_4$
5	A_{11}	$O_7 + A_3$
6	A_{12}	A_4, H_4 or $2T_2$
7	A_{12}	$I + T_3$
8	A_{12}	N_7/O_7 plus T_2
9	A_{13}	$A_3,\ T_2 + A_1,\ I + T_2,\ O_7 + A_1$
10	A_{14}	$T_2,\ 2A_1,\ I + A_1,\ N_7/O_7$
11	A_{15}	ordinary A_1 or actual I
12	A_{16}	none
13	B_{14}	N_7/O_7
14	B_{15}	ordinary A_1 or actual I
15	B_{16}	none
16	D_{13}	$I + T_2$ or $O_7 + A_1$
17	D_{14}	ordinary $I_2(m)$, $m \geq 4$, or N_7/O_7
18	D_{15}	ordinary A_1 or actual I
19	D_{16}	none

Proof organization and remaining case proofs. The dominant-component count starts afresh at rank sixteen. The full partition proof, including the actual D_{11} boundary and B/D short-tail inventories, is the "Complete rank-sixteen reduction" in Addendum A, catalogue A5. Its conditioned A_{11} bound is proved before the partition. The table lists necessary forests only. One independent geometric refinement used in its consumers is proved next. □

5.6.1 A small component calculation

Lemma 5.11. *At the original farthest vertex of any compact Coxeter polytope, a whole A_3 component has energy $> 13/50$, with arbitrary finite spectators. Arithmeticity is not needed.*

Proof. The inverse unit Gram and normalized old duals are

$$C = \begin{pmatrix} 3/2 & 1 & 1/2 \\ 1 & 2 & 1 \\ 1/2 & 1 & 3/2 \end{pmatrix}, \qquad d_i = C_{ii}, \quad u_i = \omega_i/\sqrt{d_i}.$$

Suppose $e = \|x_{A_3}\|^2 \leq 13/50$. Put $\tau_i = \tanh(L_i/2) \leq x \cdot u_i$ and $H(t) = 2t/(1+t^2)$. The logarithm of H is concave on $(0,1)$: its second derivative is $-1/t^2 - 2(1-t^2)/(1+t^2)^2 < 0$. For a pair of normalized duals with correlation k, Cauchy gives $\tau_i + \tau_j \leq \sqrt{2e(1+k)}$; hence

$$\tanh L_i \tanh L_j \leq H\big(\sqrt{e(1+k)/2}\big)^2.$$

At an adjacent A_3 pair $k = 1/\sqrt{3}$, the right side is at most $5200(1+k)/(113+13k)^2 < k$. The cleared positive difference is $(22876k - 12662)/3 > 0$, using $k > 4/7$. The compact proof's actual

distinct-opposite tangent-product inequality requires the opposite inequality if the two walls are distinct. Thus adjacent directions share their actual opposite, and connectivity gives one wall c closing all three edges.

The individual bound is

$$\tanh^2 L_i \le \frac{4e}{(1+e)^2} \le r := 2600/3969.$$

Write $c = Ap - v_A - v_S$, $h = A^2 > 0$, $v_A = \sum \beta_i \omega_i$, and $\beta_i = -(c, n_i) \ge 0$. Put $q = \beta^T C \beta$, $Q = \|v_S\|^2$, so $h = q + Q - 1$. For $v_i = (C\beta)_i/\sqrt{d_i} > 0$ we have

$$q - 1 \le h \le r \min_i v_i^2. \tag{5.8}$$

The compact proof's reciprocal-shadow and orthogonal-opposite lemmas say that every positive whole spectator contribution is at least $1/4$ and strictly below h. In particular $h < 1/4$ forces $Q = 0$.

Two adjacent positive β_i would give, at the finite endpoint of the third edge, a white triangle with all three contacts at least $1/2$. Its Gram is nonpositive, affine at equality, impossible. Thus there are at most two contacts, and two can only be at endpoints. For endpoint values $a, b > 0$,

$$q = \tfrac{3}{2}(a^2 + b^2) + ab, \qquad v_2^2 = (a+b)^2/2.$$

If both are $1/2$, the original principal is affine with a nonzero future feasible null sum. Otherwise the native gap gives $a, b \ge 1/2$ and one at least $1/\sqrt{2}$. Consequently

$$q - r v_2^2 \ge \frac{3(3-r)}{8} + \frac{(1-r)\sqrt{2}}{4} = \frac{27921 + 2738\sqrt{2}}{31752} > 1,$$

using $\sqrt{2} > 7/5$. This contradicts (5.8). There is at least one contact, since each $v_i > 0$.

For one endpoint contact β, $q = 3\beta^2/2$ and $\min v_i^2 = q/9$. Equation (5.8) gives

$$q \le 35721/33121, \quad \beta^2 \le 23814/33121 < 3/4, \quad h \le 2600/33121 < 1/4.$$

Only native three, four or five remains, and each has $q < 1$. For five use $\cos^2(\pi/5) < 2/3$, following from $\sqrt{5} < 7/3$. Then $Q = 0$ and $h = q - 1 < 0$, impossible.

For one middle contact, $q = 2\beta^2$, $\min v_i^2 = q/3$. Thus $q \le 11907/9307$ and $\beta^2 \le 11907/18614 < 16/25 < \cos^2(\pi/5)$. Only native three or four remains, so first conclude $q \le 1$, and only then strengthen the height bound to $h \le r/3 = 2600/11907 < 1/4$. Again $Q = 0$ and $h = q - 1 \le 0$ is impossible. All contact and foreign-spectator alternatives have been exhausted. □

Proposition 5.12 (All rows except whole D_{16})**.** *Groups1–18 are impossible at that same selected original vertex.*

Proof. Groups1–18 are excluded by the complete printed dispatch in Addendum A, "Every rank-sixteen row closes at its own rank" (catalogue A5). Its separate actual rank-one branches exhaust the same opposite wall, including the boundary $h = 1/2$. Every overlapping endpoint/head/foot pressure is paid within its local proof before disjoint whole-component costs are added. The universal $A_{12} > 5/7$ estimate supplies both ordinary and typed consumers. Only group 19, whole D_{16}, remains for the next section. □

5.7 The last actual diagram

Proposition 5.13 (Whole-D_{16} reduction)**.** *If group 19 occurs, the actual field is quadratic. The actual long head and both genuine signed forks are isolated and dotted. There are integral generators $U, F, G \in \mathcal{O}_K$ satisfying*

$$4 < U < 98/9, \quad 0 < U' < 2, \qquad 4 < F, G < 121/2, \quad 0 < F', G' < 1/2, \tag{5.9}$$

$$UF/4,\ UG/4,\ FG/4 \in \mathcal{O}_K. \tag{5.10}$$

Their raw time squares are

$$T_H = U(U-2), \qquad T_F = 2F(2F-1), \qquad T_G = 2G(2G-1), \tag{5.11}$$

and every pair of time products is a square in K.

We give its field and endpoint reduction below, after specifying the initial geometric inventory and two scalar estimates. The inventory must be proved at actual rank sixteen; it is not supplied by a higher-rank head theorem or by the dyadic calculation.

The original coordinates satisfy

$$x_1 > \cdots > x_{15} > |x_{16}|, \quad P_j = e_1 + \cdots + e_j, \quad S_j = x \cdot P_j, \quad f_j = \sqrt{j} - .61/\sqrt{2}.$$

The same reduction gives $S_j > f_j$ for $5 \le j \le 14$ and one positive prefix reference w with $x \cdot w > E_{14} = \|w\|^2$ and

$$E_{14} > .9084, \quad b < .361, \quad \beta_5 > .14, \quad \beta_{14} > .13, \quad f_{14} < 3.311, \qquad S_6 > 2.01815. \tag{5.12}$$

Here b is the reference's first-five mean. These coefficients are spent individually below. The head raw root has square $2U$, spatial part Ue_1; the fork raw roots have squares $2F, 2G$ and spatial parts $(F/2)(P_{15} + e_{16})$, $(G/2)(P_{15} - e_{16})$. They remain two distinct directions even if $F = G$.

After the following initial inventory, the field and dyadic arguments are given in full. Every original cycle remains mandatory even when only its necessary consequences are used.

Lemma 5.14 (Initial whole-D_{16} inventory)**.** *At the same original whole-D_{16} vertex, $S_j > f_j$ for $5 \le j \le 14$, and the positive reference has* (5.12)*. The actual long head is isolated, with finite own label at least seven or dotted own. Every actual fork has edge cosh less than twelve and is either isolated dotted or has exactly one simple-three contact at an endpoint of its retained A_{15} path, with positive own contact. Own zero and parallel fork roles are excluded. If both forks are isolated, the actual field has degree two or three.*

Proof location for the initial inventory. The initial inventory is catalogue A5's actual-rank geometric input. Its complete proof in Addendum A, "The last actual diagram," checks all original walls, closes the late spine and both signed fork inventories, and records coefficients before use. It is not a specialization of a higher-rank theorem. Starting from exactly this inventory, the following argument proves the necessary quadratic triple and excludes it. □

Lemma 5.15 (Two exact terminal estimates)**.** *For $0 < u < 1$, put*

$$H(u) = \frac{\sqrt{2-u} - \sqrt{u(1-u)}}{\sqrt{2}}, \qquad R(u) = 1 - u + u^2/2.$$

Then $.8H(u) - .16R(u) > .295$. Also a degree-three algebraic integer with conjugates $A, a, b > 0$, $a, b < 1/2$ and norm eight has $A > 42$.

Proof. For the first assertion let $c = .681$, $t = 2 - c^2 = 1.536239$. The radical difference is positive, and $t - u^2 > 0$. It suffices to prove

$$P(u) = (t - u^2)^2 - 4u(1-u)(2-u) > 0.$$

With $r = u - .7$ the exact expansion is

$$P = r^2(r - .6)^2 + 3.107522r^2 - .0094692r + .002616045121.$$

The last three terms exceed $3r^2 - .01|r| + .0026$. As $.01|r| \le r^2 + .000025$, this is positive. Thus $H(u) > c/\sqrt{2} > .4815351$ on the entire interval.

Moreover R decreases, and H decreases on $(0, .65]$. Below $1/2$ the derivative sign is immediate; between $1/2$ and $.65$ it follows from

$$\frac{2u-1}{\sqrt{u(1-u)}} \le \frac{.3}{\sqrt{.2275}} < \frac{1}{\sqrt{1.5}} \le \frac{1}{\sqrt{2-u}},$$

where $.09 \cdot 1.5 < .2275$ proves the strict inequality. The following rational radical seals give lower bounds for $H(v)$:

v	$\sqrt{2-v} >$	$\sqrt{v(1-v)} \le$	$H(v) >$
.25	1.3228	.4331	.62
.4	1.2649	.4899	.54
.5	1.2247	.5	.51
.55	1.2041	.4975	.499
.6	1.1832	.4899	.4902
.625	1.1726	.48413	.4867
.65	1.16189	.47697	.4842.

Square the first two columns and use $\sqrt{2} < 1.4143$ to obtain each last entry. Taking H at the right endpoint and R at the left endpoint gives the complete cover

$(0, .25]$	.336	$[.25, .4]$	.307
$[.4, .5]$	.2992	$[.5, .55]$	.2992
$[.55, .6]$	.29596	$[.6, .625]$	.29656
$[.625, .65]$	.29611	$[.65, 1)$	.29542808.

These are strict lower bounds for $.8H - .16R$; the last uses the global radical bound and $R(.65) = .56125$. Every endpoint is covered.

For the second assertion suppose $A \le 42$. Norm eight and $a, b < 1/2$ give $A > 32$ and integer trace $T \in \{33, \dots, 42\}$. Its minimal polynomial is $X^3 - TX^2 + BX - 8$, with integer B. Since $A > T - 1$, AM–GM gives

$$B \ge 2\sqrt{8A} + 8/A > 2\sqrt{8(T-1)} + 8/T.$$

Evaluating the polynomial at $1/2$, between both small roots and the large root, gives $B < T/2 + 63/4$. For even traces the largest permitted B, and the positive seal $32(T-1) - B^2$, are

T	34	36	38	40	42
$B_{\max}$	32	33	34	35	36
$32(T-1) - B_{\max}^2$	32	31	28	23	16.

They contradict the lower bound. At $T = 33$, $B_{\max} = 32$ and the same lower bound is already strictly larger than 32. At odd traces $35, 37, 39$, the permitted maxima are $33, 34, 35$; the exact integers

$$32(T-1)T^2 - (TB_{\max} - 8)^2 = 17191, 14588, 8087$$

are positive and give the same contradiction after squaring. Only $T = 41$ remains. The two bounds force $B = 36$, but $X^3 - 41X^2 + 36X - 8$ has discriminant -2704, incompatible with three real conjugates. Therefore $A > 42$. □

Proof of Proposition 5.13, from Lemma 5.14. Use raw old roots of square two. An isolated head has raw square $2U$, spatial part Ue_1, time square $U(U-2)$, and actual field $K = \mathbb{Q}(U)$; its nonphysical U_σ lie in $(0, 2)$. An isolated fork has the data stated above, actual $K = \mathbb{Q}(F)$, and $0 < F_\sigma < 1/2$ at every nonphysical place. The only nonisolated forks left by the initial inventory have raw square two and one endpoint-three white contact. Their complete data are

role	spatial vector	time square M	edge cosh
far	$(t/2)(P_{15} + se_{16}) + e_1$	$4t^2 + t - 1$	$(8t+1)/\sqrt{17}$
near	$(t/2)(P_{15} + se_{16}) + (P_{15} - se_{16})/2$	$4t^2 + 7t + 2$	$(8t+7)/\sqrt{17}$.

Here $s = \pm 1$ denotes a genuine fork. In the far role put $z = t$; in the near role put $z = t + 1$. Path-cycle integrality gives integral z, and actual-field descent gives $K = \mathbb{Q}(z)$. To justify these normalizations, the connected simple-three old system has one raw norm squareclass, so a totally positive rescaling of the form puts its roots at square two over K. An isolated own contact has integral two-cycle $4\gamma^2$; rescaling its root line by its nonzero old product gives the raw head or fork displayed above. The raw principal consisting of that cap and the full old system is defined over $\mathbb{Q}(U)$ or $\mathbb{Q}(F)$ and is physically indefinite, since the cap has nonzero time. The descent argument in Section 5.1 proves the asserted field equalities. For an endpoint cap its simple-three helper permits square-two normalization directly; the closed old path gives integral t, and the same indefinite-principal argument gives $K = \mathbb{Q}(t)$. The cosh bound twelve gives $z < 6.311$; positive own gives $z \geq 1$ far and $z \geq 2$ near. Negative nonphysical time squares give

$$-.65 < z_\sigma < .4 \quad \text{far}, \qquad -.4 < z_\sigma < .65 \quad \text{near}. \tag{5.13}$$

All these formulas concern actual K-roots, so every positive raw time product is K-valued and its square must be a full K-square.

First sharpen the head bound. If $x_1 \geq .632$, rebuild only the first-five flat block of the positive reference, preserving its sum. The new coordinates are $.632, (f_5 - .632)/4$ four times. They exceed the next slope and preserve nonnegative prefix prices. Their single-reference gain gives

$$\|x\|^2 > E_{14} + \tfrac{5}{4}(.632 - .361)^2 = 1.00020125,$$

a contradiction. Thus $x_1 < .632$. The actual head edge gives $C_H < 7/3$, since $.632^2 < 2/5$, and hence always $U < 98/9$; if dotted, $U > 4$.

Endpoint cycles and the actual degree. For an endpoint cap let $g = -H \cdot C$ and $L = g^2/U$. The original path and own two-cycle make both integral. Expanding the common time product gives

$$\begin{aligned} L + g(t+2) &= (15t^2 - 8)U/4 - 2M && \text{far}, \\ L + g(t+1) &= (15z^2 - 4z - 4)U/4 - 2M && \text{near}. \end{aligned}$$

All remaining terms are integral, and 15 is coprime to four; therefore $z^2U/4$ is integral in both roles. If $|\,\mathrm{N}\,z|$ is odd, z is a unit at every dyadic prime, forcing $U/4$ integral. This is impossible in degree at least three. Indeed the sharpened head bound just proved gives

$$0 < \mathrm{N}(U/4) < \frac{49}{18}(1/2)^{d-1} \le \frac{49}{72} < 1 \qquad (d \ge 3),$$

contradicting its positive integral norm. This uses the original head bound and all nonphysical places, with no integer-basis assumption.

The nonzero endpoint norm is less than $6.311(.65)^{d-1}$. It excludes $d \ge 6$; at degrees four or five it is below two and hence has odd absolute norm one, just excluded. In degree three only $|\,\mathrm{N}\,z| = 2$ remains. Degree one is excluded by Meyer. If neither endpoint role occurs, both forks are isolated and the initial lemma already gives degree two or three. Thus these are the only actual degrees in every case.

Every cubic endpoint is impossible. In degree three norm magnitude two and $6.311 \cdot .4 \cdot .65 < 2$ force both small conjugates negative far, positive near. Write their positive magnitudes $a, b < 13/20$; then $zab = 2$ and $z > 800/169$. In the far role its integer trace $z - a - b$ is four or five, by AM–GM and $z < 6.311$. The second coefficient $B = 2/z - z(a + b)$ satisfies $B \le -6$ by AM–GM. The polynomial $X^3 - TX^2 + BX - 2$ is negative at $-13/20$, so $-B < (13/20)^2 + T(13/20) + 40/13 < 7$ and $B \ge -6$. Thus $B = -6$. The two resulting cubics, with $T = 4, 5$, have discriminants $-44, -424$, impossible in a totally real field. Near, the function $z + 2\sqrt{2/z}$ is increasing for $z > 4$. The integer trace is therefore seven, since $z + 2\sqrt{2/z} > 800/169 + 13/10 > 6$ and $z + a + b < 8$. Thus $z > 57/10$. Evaluation at $13/20$ gives $B < 7.205$, whereas $B \ge 2\sqrt{2z} + 2/z > 6.75 + .316 > 7$. The incompatible integer bounds exclude this role too.

Every finite isolated head is impossible. For finite own m, $U = 4\cos^2(\pi/m)$ has degree $\phi(m)/2$. The complete totient-at-most-six list and the initial small-own exclusions leave $m = 7, 8, 9, 10, 12, 14, 18$. At $m = 9, 14$ a nonphysical $U_\sigma > 2$ is forbidden. At cubic $m = 7, 18$, $\mathrm{N}\,U = 1, 3$ is odd. The cubic endpoint exclusion makes both forks isolated; their cycle $UF/4$ forces $F/4$ integral, so $64 \mid \mathrm{N}\,F$. But $F < 72$ and its other conjugates below $1/2$ give $\mathrm{N}\,F < 18$, a contradiction.

The remaining heads have $U = 2 + \sqrt{2}, 2 + \varphi, 2 + \sqrt{3}$. The full endpoint strips, in the maximal orders, give

K	far t	near z	$U(U-2)$
$\mathbb{Q}(\sqrt{2})$	$1 + \sqrt{2}, 3 + 2\sqrt{2}$	$2 + \sqrt{2}, 3 + 2\sqrt{2}$	$2 + 2\sqrt{2}$
$\mathbb{Q}(\varphi)$	$\varphi, \varphi^2, \varphi^3$	φ^2, φ^3	$1 + 3\varphi$
$\mathbb{Q}(\sqrt{3})$	$2 + \sqrt{3}$	$2 + \sqrt{3}$	$3 + 2\sqrt{3}$.

For $\sqrt{2}$ the radical coefficient is one or two, for $\sqrt{3}$ one. In the golden integral basis the coefficient is one or two; the next cover already exceeds the exact cosh bound. These finite integer intervals prove the list, including dotted owns. The corresponding time-product norms are $72, 400, 36, 544$; $5, 5, 380, 95, 180$; and $78, 150$. Only $400, 36$ are squares. Their products are $340 + 240\sqrt{2}$ and $102 + 72\sqrt{2}$; the rational square parts would be $180, 160$ or $54, 48$, none square. Thus no endpoint is possible for a finite head.

For an isolated fork in the $m = 10, 12$ fields the head is a dyadic unit, so $F/4$ is integral. At $m = 8$, normalize the unique dyadic valuation by $v(\sqrt{2}) = 1$. The quarter-cycle gives $v(F) \ge 3$; the time-product valuation is $4 + v(F)$ and must be even, hence $v(F) \ge 4$ and again $F/4$ is integral. For $W = F/4$, the full strips are $W < 18$, $0 < W' < 1/8$. In $\mathbb{Z}[\sqrt{2}]$ no coefficient with $1 \le b \le 6$ fits; in $\mathbb{Z}[\varphi]$ only $W = 5 + 8\varphi$ fits; in $\mathbb{Z}[\sqrt{3}]$ only $W = 7 + 4\sqrt{3}$ fits. The last two give time-product norms $25280, 9024$, nonsquares. Thus the actual head is dotted.

One dotted-head reference excludes the remaining cubic. Put $u = 4/U \in (0,1)$. The positive actual core $c_H + n_1/g_H$ has zero product with c_H, negative product with n_1, future time and nonpositive product with every other original wall. Normalizing it gives

$$x \cdot Z \geq H(u), \quad Z = (1-u/2)e_1 + (u/2)e_2, \quad Z^2 = R(u).$$

The single graft $.4(Z - P_5/5)$ spends .08 at P_5, below its available .14, and has zero cross with the flat block. Lemma 5.15 therefore gives gain $> .295 - .8(.361) + .032 = .0382$, hence one reference of energy $> .9466$. A signed $.13(P_{15} + se_{16} - P_{14})$ graft has square residual two and zero cross. If its fork cosh were at least eleven, its actual pressure would exceed 3.65, giving

$$\|x\|^2 > .9466 + .26(3.65 - 3.311) - .0338 = 1.00094.$$

Both fork coshes are therefore below eleven.

In a cubic field all endpoint roles were excluded. The isolated forks satisfy $4 < F, G < 121/2$ and $\mathrm{N}\,F, \mathrm{N}\,G \in \{1, \dots, 15\}$. The quarter-cycle gives $64 \mid \mathrm{N}\,F\,\mathrm{N}\,G$, forcing both norms eight. Lemma 5.15 gives $F, G > 42$. Their coshes exceed $\sqrt{84} > 9.16$ and their signed pressures exceed 3.584, since $3.584^2 < 1632/127$. Averaging the two genuine fork inequalities gives $S_{15} > 3.584$. The average tail graft has residual square one, so the same reference gives

$$\|x\|^2 > .9466 + .26(3.584 - 3.311) - .0169 = 1.00068.$$

Thus the actual field is quadratic.

Every quadratic endpoint is impossible. The improved cosh bound eleven gives $z < 5.796$, so $|\mathrm{N}\,z| < 4$. If its norm is odd, $U/4$ is integral. The full head strip then gives $U = 4\varphi^2$, with time square $8\varphi^5$. In this field the complete endpoint strips give far $t = \varphi, \varphi^2, \varphi^3$ and near $z = \varphi^2, \varphi^3$. The first two far time products are $8\varphi^{10}, 8\varphi^{12}$, requiring $\sqrt{2}$ in the field; the third has nonsquare norm 4864. The near products have nonsquare norm 1216, or equal $48\varphi^{10}$, requiring $\sqrt{3}$. None is possible.

Only norm magnitude two remains. Integer trace and the exact role-specific cosh/conjugate strips give the full list

role	$\mathrm{N}\,z$	$\mathrm{Tr}\,z$	z
far	-2	$3,4,5$	$(3+\sqrt{17})/2,\ 2+\sqrt{6},\ (5+\sqrt{33})/2$
near	2	$4,5,6$	$2+\sqrt{2},\ (5+\sqrt{17})/2,\ 3+\sqrt{7}$
near	-2	5	$(5+\sqrt{33})/2.$

For example far positive norm two would require $z > 5$ and $5 < z + 2/z < 6$, so is impossible. The other rows follow from $z + \mathrm{N}(z)/z$ and the strict conjugate intervals, not a field search.

In the ramified fields $\sqrt{2}, \sqrt{6}, \sqrt{7}$, z has dyadic valuation one. Integrality of $z^2U/4$ forces $U/2$ integral. Writing $W = U/2$, its strips are $2 < W < 49/9$, $0 < W' < 1$. The only radical coefficient is one: the candidates are $2+\sqrt{2},\ 3+\sqrt{6},\ 3+\sqrt{7}$. The last two exceed $49/9$; the first gives $U = 4 + 2\sqrt{2}$. Its near cap has time square $21 + 15\sqrt{2}$ and the head $16 + 12\sqrt{2}$; their product norm 288 is nonsquare.

In the split fields $\sqrt{17}, \sqrt{33}$, taking norms in $z^2U/4$ gives $4 \mid \mathrm{N}\,U$. The full head strips retain precisely

$\mathbb{Q}(\sqrt{17})$	$(7+\sqrt{17})/2,\ 5+\sqrt{17}$
$\mathbb{Q}(\sqrt{33})$	$(7+\sqrt{33})/2,\ (9+\sqrt{33})/2.$

In the first field the full integral-basis coefficient is one or two, and the omitted strip heads have norms two and nineteen; in the second it is one. No maximal-order class is missing. For $s = \sqrt{17}$ the head time squares are $(19 + 5s)/2$, $32 + 8s$. The far cap has time square $(53 + 13s)/2$. Its first time product is $(16 + 4s)^2$, but the spatial product is $(33 + 7s)/4$; hence the actual path cycle would be $(31 + 9s)/4$, not integral. The second product is $1732 + 420s$, whose possible rational square parts $882, 850$ are nonsquares. The near cap has time square $(77 + 19s)/2$ and product norms $832, 3328$, also nonsquares. For $s = \sqrt{33}$ both head time squares have norm -24; the far and near caps have norms $-98, -8$, giving nonsquare product norms $2352, 192$. Thus every endpoint is excluded.

Both genuine forks are consequently isolated dotted. Their original cycles, in these raw normalizations, give

$$L_{HF} + g_{HF} = (15UF - 32F - 8U + 16)/4,$$
$$L_{FG} + 7g_{FG} = 15FG/4 - 8(F + G) + 4,$$

with integral g, L; the analogous head/G identity holds. These yield all three quarter-products. The common actual K-time line supplies every field-square condition. The strict physical and nonphysical strips were retained throughout. This proves the claimed necessary quadratic triple from the initial inventory. □

5.7.1 Norm bounds, integer orders and valuations

Put

$$N_H = \mathrm{N}\, U, \quad T = \mathrm{Tr}\, U, \quad A = -\mathrm{N}(U - 2) = 2T - 4 - N_H.$$

The strict strips imply

$$5 \le T \le 12, \quad 1 \le N_H \le 21, \quad 1 \le A \le 17.$$

For a fork put $n = \mathrm{N}\, F$, $S = \mathrm{Tr}\, F$ and $B = 2S - 4n - 1 > 0$ odd. Then

$$1 \le n \le 30, \quad 5 \le S \le 60, \qquad \mathrm{N}\, T_H = -N_H A, \quad \mathrm{N}\, T_F = -4nB. \tag{5.14}$$

A nonsquare rational norm disproves a field square; a square norm still requires the full square test.

Use the full integer order of $K = \mathbb{Q}(\sqrt{d})$. The gap between the embeddings of U is a positive integral multiple of $\sqrt{\Delta}$, so

$$\Delta < (98/9)^2 < 119, \qquad \Delta \le 118. \tag{5.15}$$

For a prime $\mathfrak{p}$ over two, normalize $v_{\mathfrak{p}}(\mathfrak{p}) = 1$ and write

$$e = v_{\mathfrak{p}}(2),\ r = v_{\mathfrak{p}}(U),\ s = v_{\mathfrak{p}}(U - 2),\ k = v_{\mathfrak{p}}(F),\ l = v_{\mathfrak{p}}(G).$$

The factors $2F - 1, 2G - 1$ are units there. Quarter integrality and the time squares give

$$r + k, r + l, k + l \ge 2e, \qquad r + s + e + k,\ r + s + e + l \equiv 0 \pmod 2. \tag{5.16}$$

Also $s = r$ for $r < e$, $s = e$ for $r > e$, and $s \ge e$ for $r = e$. Norm valuations are $\sum f_{\mathfrak{p}} v_{\mathfrak{p}}$. The quadratic possibilities are inert $(e, f) = (1, 2)$, ramified $(2, 1)$, or two split primes $(1, 1)$. No PID or unramifiedness assumption enters.

5.7.2 Three complete integer-strip lemmas

If $F/4$ is integral, $16 \mid n < 30.25$ gives $n = 16$. For $W = F/4$, norm one and $0 < W' < 1/8$ give $8 < W < 121/8$, and $\operatorname{Tr} W = W + 1/W \in \{9, \dots, 15\}$. This gives the complete table

$\operatorname{Tr} W$	K	Δ	F when $\Delta \le 118$
9	$\mathbb{Q}(\sqrt{77})$	77	$18 + 2\sqrt{77}$
10	$\mathbb{Q}(\sqrt{6})$	24	$20 + 8\sqrt{6}$
11	$\mathbb{Q}(\sqrt{13})$	13	$22 + 6\sqrt{13}$
12	$\mathbb{Q}(\sqrt{35})$	140	excluded
13	$\mathbb{Q}(\sqrt{165})$	165	excluded
14	$\mathbb{Q}(\sqrt{3})$	12	$28 + 16\sqrt{3}$
15	$\mathbb{Q}(\sqrt{221})$	221	excluded.

If $Y = U/2$ is integral with odd norm, then $2 < Y < 49/9$, $0 < Y' < 1$ and $\operatorname{N} Y \in \{1, 3, 5\}$. Its integer trace t is in $3, \dots, 6$ and $1 - t + \operatorname{N} Y < 0$, so $t > \operatorname{N} Y + 1$. The complete list is

$\operatorname{N} Y$	t	Y	2 in K
1	3	$(3 + \sqrt{5})/2$	inert
1	4	$2 + \sqrt{3}$	ramified
1	5	$(5 + \sqrt{21})/2$	inert
3	5	$(5 + \sqrt{13})/2$	inert.

Trace six gives $3 + 2\sqrt{2}$ or $3 + \sqrt{6}$, both above $49/9$; norm five requires trace above six. These bounded integer comparisons prove completeness.

Finally, if $U/4$ is integral with $\operatorname{N} U = 16$, then $W = U/4$ has norm one, $1 < W < 49/18$, $0 < W' < 1/2$. Thus $W > 2$, $\operatorname{Tr} W = 3$, and

$$U = 4\varphi^2, \quad K = \mathbb{Q}(\varphi), \qquad \varphi = (1 + \sqrt{5})/2. \tag{5.17}$$

5.8 Odd head norm and inert two

5.8.1 Odd norm

If N_H is odd, at every dyadic prime $r = s = 0$. In an inert or split field, (5.16) makes both fork valuations odd and at least three; their norms are divisible by 64, contrary to $n \le 30$. Thus two ramifies. Now k, l are even and at least four. Both fork norms are sixteen and $F/4, G/4$ integral. The unit-fork table leaves only $\mathbb{Q}(\sqrt{3})$ or $\mathbb{Q}(\sqrt{6})$. The full head strips with odd norm give

K	U	$\operatorname{N} T_H$	$\operatorname{N} T_F$
$\mathbb{Q}(\sqrt{3})$	$5 + 2\sqrt{3}$	-39	-3008
$\mathbb{Q}(\sqrt{6})$	$3 + \sqrt{6},\ 5 + 2\sqrt{6}$	$-15,\ -15$	$-960.$

The radical coefficient is one or two; coefficient three exceeds the physical bound. The conjugate strip and norm parity remove the other rational coefficients. In $\mathbb{Q}(\sqrt{6})$, $6 + 2\sqrt{6} > 98/9$ and $4 + \sqrt{6}$ has even norm.

The $\sqrt{3}$ product norm is $64 \cdot 1833$, a nonsquare. For $U = 3 + \sqrt{6}$ the exact time product is $58200 + 23760\sqrt{6}$, of norm 120^2. Its possible rational square parts are 29160 and 29040, strictly between 170^2 and 171^2, so it is not a field square. The remaining triple is $U = 5 + 2\sqrt{6}$, $F = G = 4U$. The actual head and two genuine fork edges have coshes

$$C_H = (2 + \sqrt{6})/2 > 11/5, \qquad C_+ = C_- = 2(2 + \sqrt{6}) > 44/5,$$

using $\sqrt{6} > 12/5$. Equation (5.1) gives $x_1 \geq \sqrt{3/8} > .61$ and, for both $s = \pm 1$,

$$x \cdot (P_{15} + se_{16}) > 4\sqrt{39}/7 > 3.56.$$

The last comparison follows from $6.23^2 < 39$. Averaging the two signed inequalities gives $S_{15} > 3.56$; Equation (5.12) gives $S_6 > 2.01815$. One positive reference gives

$$\|x\|^2 > .61^2 + 1.40815^2/5 + 1.54185^2/9 = 1.032821887 > 1.$$

The last pressure averages the two genuine fork edges, not a virtual root. This excludes odd norm.

5.8.2 Even norm with inert two

Here $e = 1, f = 2$, and $r = 1$ or two since $r \geq 3$ forces $N_H \geq 64$. If $r = 1$, the head strip gives

$$U = 3 + \sqrt{5}, \quad 5 + \sqrt{21}, \quad 5 + \sqrt{13}.$$

In each case $s = 1$, so (5.16) makes k, l odd and positive. The norm bound gives $k = l = 1$. Thus $W = F/2$ is integral with odd norm, $W < 121/4$, $0 < W' < 1/4$. In $W = (a + b\sqrt{d})/2$, $a \equiv b \pmod 2$, the full strip $0 < a - b\sqrt{d} < 1/2$ gives

d	b range	(a, b) before norm filter	F
5	$1, \ldots, 13$	$(7, 3), (18, 8), (25, 11)$	$2\varphi^4,\ 2\varphi^6,\ 14 + 22\varphi$
21	$1, \ldots, 6$	$(5, 1), (23, 5)$	$5 + \sqrt{21},\ 23 + 5\sqrt{21}$
13	$1, \ldots, 8$	$(11, 3), (22, 6)$	$11 + 3\sqrt{13}$.

The $d = 13$ pair $(22, 6)$ has $\mathrm{N}\, W = 4$, not odd. Each b bound follows from $b\sqrt{d} < W < 121/4$; the displayed a is the only possible integer with the required parity. The close pair $(28, 6)$ at $d = 21$ fails since $6\sqrt{21} < 55/2$.

At $d = 5$, product norms are 256 times $11, 55, 95$, all nonsquares. The other square-norm exceptions are

d	F	$T_H T_F$	$\sqrt{\mathrm{N}(T_H T_F)}$	possible square parts
21	$5 + \sqrt{21}$	$12648 + 2760\sqrt{21}$	48	6348, 6300
21	$23 + 5\sqrt{21}$	$303000 + 66120\sqrt{21}$	240	151620, 151380
13	$11 + 3\sqrt{13}$	$52872 + 14664\sqrt{13}$	144	26508, 26364.

These lie strictly between $79^2, 80^2$; $389^2, 390^2$; and $162^2, 163^2$, respectively. No fork exists.

If $r = 2$, then $\mathrm{N}\, U = 16$, $U/4$ integral, and (5.17) holds. Now $s = 1$, so k, l are even. The fork/fork quarter-product gives $k + l \geq 2$, so one fork is divisible by four with norm sixteen. Its $W = F/4$ in $\mathbb{Q}(\varphi)$ has norm one, $W < 121/8$, $0 < W' < 1/8$. Writing $W = a + b\varphi$ gives $b = 1, \ldots, 6$; none permits $0 < a + b(1 - \varphi) < 1/8$. The inert case is exhausted.

5.9 Even head norm with ramified two

Here $e = 2, f = 1$ and $r = v_2(N_H) \in \{1, 2, 3, 4\}$. If $r = 1$, $s = 1$ and both fork valuations are even and at least four. The unit-fork table leaves $\sqrt{6}$ or $\sqrt{3}$. The head strips with $v_2(N_H) = 1$ give only $U = 4 + \sqrt{6}$ or $3 + \sqrt{3}$. Their product norms are $256 \cdot 75$ or $256 \cdot 141$, nonsquares.

If $r = 2$, the odd-norm $U/2$ table forces $U = 4 + 2\sqrt{3}$. Then $s = 3$, so k, l are odd and at least three. Norm bounds permit only valuation three and norm eight or twenty-four. For $F = 2W$,

therefore, $\mathrm{N}\,W = 2$ or six, $W < 121/4$, $0 < W' < 1/4$. In $\mathbb{Z}[\sqrt{3}]$, the strip for $W = a + b\sqrt{3}$, $1 \le b \le 8$, allows only $(7,4), (14,8)$, of norms one and four. Neither required norm occurs.

If $r = 3$, $U/2$ is integral of norm two and trace four, five or six. Trace six gives $3 + \sqrt{7} > 49/9$; trace five gives the split field $\mathbb{Q}(\sqrt{17})$. Thus ramification forces $U = 4 + 2\sqrt{2}$. Here $s = 2$, so k, l are odd, at most three. Since $k + l \ge 4$, at least one is three and its norm is eight or twenty-four.

For that head $\mathrm{N}\,T_H = -32$, so a product norm is $128n(2S - 4n - 1)$ with S even in $\mathbb{Z}[\sqrt{2}]$. At $n = 8$, a square would require $2S - 33$ square, but it is three modulo four. At $n = 24$, the strip gives $49 \le S \le 60$; the even values yield

$$3(2S - 97) \in \{9, 21, 33, 45, 57, 69\}.$$

Only nine is square, at $S = 50$. But even norm for $F = a + b\sqrt{2}$ forces a even, hence $S = 2a$ divisible by four. This excludes fifty.

Finally $r = 4$ gives (5.17), whose field has inert two. Larger r forces $N_H \ge 32 > 21$. All ramified cases are excluded.

5.10 The complete split branch

Here $d \equiv 1 \pmod 8$, squarefree, and (5.15) leaves $d = 17, 33, 41, 57, 65, 73, 89, 97, 105, 113$. Write $U = (a + b\sqrt{d})/2$, $a \equiv b \pmod 2$. The strip gives $0 < a - b\sqrt{d} < 4$ and $b = 1$, except at $d = 17$ where $b = 1$ or two. The complete even-norm heads are

d	U	T	N_H	A
17	$(5 + \sqrt{17})/2$	5	2	4
17	$(7 + \sqrt{17})/2$	7	8	2
17	$5 + \sqrt{17}$	10	8	8
33	$(7 + \sqrt{33})/2$	7	4	6
33	$(9 + \sqrt{33})/2$	9	12	2
41	$(7 + \sqrt{41})/2$	7	2	8
41	$(9 + \sqrt{41})/2$	9	10	4
57	$(9 + \sqrt{57})/2$	9	6	8
57	$(11 + \sqrt{57})/2$	11	16	2
65	$(9 + \sqrt{65})/2$	9	4	10
65	$(11 + \sqrt{65})/2$	11	14	4
73	$(9 + \sqrt{73})/2$	9	2	12
73	$(11 + \sqrt{73})/2$	11	12	6
89	$(11 + \sqrt{89})/2$	11	8	10
97	$(11 + \sqrt{97})/2$	11	6	12
105	$(11 + \sqrt{105})/2$	11	4	14
113	$(11 + \sqrt{113})/2$	11	2	16.

For $d = 17$, all coefficient pairs are $(5,1), (7,1), (10,2), (12,2)$; the last has odd norm nineteen. For the other fields two odd a values pass the conjugate strip; at $89, 97, 105, 113$ the larger exceeds $98/9$. These explicit finite coefficient comparisons prove the table, without an assumed field-search output.

5.10.1 Odd trace

Let $\mathfrak{p}$ be the prime where U is nonunit and $\mathfrak{q}$ its conjugate. At $\mathfrak{q}$, $r = s = 0$, so both fork valuations are odd and at least three. Put $c = v_2(N_H) = r_{\mathfrak{p}}$. A fork has total norm valuation at most four.

If $c = 2$ or four, $s_{\mathfrak{p}} = 1$ and both fork valuations at $\mathfrak{p}$ are even. The only possible ordered valuations are $(0, 3)$ for each fork, contradicting $FG/4$ at $\mathfrak{p}$.

If $c = 1$, both forks have valuations $(1, 3)$ and norm sixteen. Parity requires $v_2(A)$ odd. The table leaves only

$$d = 41,\ U = (7 + \sqrt{41})/2; \qquad d = 57,\ U = (9 + \sqrt{57})/2.$$

Now $W = F/2$ is integral of norm four, $W < 121/4$, $0 < W' < 1/4$. In $W = (a + b\sqrt{d})/2$, $b = 1, \ldots, 4$, the strip $0 < a - b\sqrt{d} < 1/2$ with parity allows only $(26, 4)$ at 41, of norm five, or $(23, 3)$ at 57, of norm four. Thus the only candidate is $F = 23 + 3\sqrt{57}$ in $K = \mathbb{Q}(\sqrt{57})$.

For that head $U > 8$, so $x_1 > 1/\sqrt{3} > .577$. Rebuilding only the first-five block of (5.12), with its sum fixed, gives

$$E_{14} + \frac{5}{4}(.577 - .361)^2 > .96672.$$

If a fork had $\cosh L \geq 9$, its actual signed pressure would exceed 3.577. The separately available .13 coefficient at P_{14} has residual square two and zero cross with that first-block change. One square would give

$$\|x\|^2 > .96672 + .26(3.577 - 3.311) - .0338 = 1.00208 > 1.$$

Both fork coshes are therefore below nine, so $F < 81/2$, contrary to $23 + 3\sqrt{57} > 44 > 81/2$.

If $c = 3$, $s_{\mathfrak{p}} = 1$ and both forks again have valuations $(1, 3)$, norm sixteen. The remaining heads have $d = 17$ or 89. For $W = F/2$ the full strip/parity pairs are

d	b range	(a, b)
17	$1, \ldots, 7$	$(21, 5), (29, 7)$
89	$1, \ldots, 3$	none.

The 17-field norms are four and two, leaving only $F = 21 + 5\sqrt{17}$. The close pair $(17, 4)$ fails since $4\sqrt{17} < 33/2$. The surviving product norm is $(-16)(-1216) = 1024 \cdot 19$, a nonsquare. There are no other odd-trace cases.

5.10.2 The sole even-trace head

The last row is $U = 5 + \sqrt{17}$. Its two prime valuations are one and two. At the valuation-two prime $s = 1$; $\mathrm{N}(U - 2) = -8$ makes $s = 2$ at the other. Equation (5.16) makes both fork valuations even at both primes, and at the valuation-one prime each is at least two. The fork/fork quarter-product forces one fork also to have valuation at least two at the other prime. It is divisible by four globally, with norm sixteen. For $W = F/4$ we have

$$\mathrm{N}\, W = 1, \quad W < 121/8, \quad 0 < W' < 1/8, \qquad W = (a + b\sqrt{17})/2, \quad 1 \leq b \leq 3.$$

None admits an integer a in $0 < a - b\sqrt{17} < 1/4$. The final split row is impossible.

Proposition 5.16. *Whole D_{16} cannot occur at the original farthest vertex of a compact arithmetic Coxeter 16-polytope.*

Proof. The actual reduction gives the necessary quadratic triple. Its head norm is odd or even. The odd case and every inert even case were excluded in Section 5.8; Sections 5.9 and 5.10 exclude ramified and split even cases. These exhaust quadratic fields. Each contradiction uses a necessary actual cycle, full field-square condition or one paid original-vector norm. No D16 census is used. □

Conclusion of the arithmetic argument. The independent compact theorem gives $n \leq 17$. The complete rank-seventeen cover, its geometric dispatch and the two empty A_{17} endpoint domains exclude seventeen. The fresh rank-sixteen cover and its geometric consumers leave only whole D_{16}; the field reduction and exhaustive manual dyadic argument above exclude that final configuration. Thus $n \leq 15$, which is Theorem 1.1(iii). □

6 GitHub proof catalogue and reproduction

The versioned repository is the mathematical supplement to this concise edition. Its full manuscript [8] contains the complete proofs of every local proposition cited above. The following guide locates the essential supplementary material.

Result	Complete proof addenda	Reproduction guide
finite volume 57	Parts I, IV; intrinsic weights, coverage, finite maximum, cusp comparison and both tails	`docs/finite_volume57_reproduce.md`
compact 17	Parts II, V; actual component estimates, own-rank covers and all endpoint/head/fork cases	`docs/compact17_consolidated_proof_map_20261025.md`
compact arithmetic 15	Parts III, VI; field reductions, four finite propositions, rank covers and the manual dyadic finish	`docs/compact_arithmetic_consolidation_proof_map_20261025.md`

Each path is relative to the fixed repository version cited above. The accompanying `ADDENDA.md` maps the propositions of this edition to exact labels in that source. The package includes the algorithm specifications, exact checkers, available certificates, execution records and instructions for serial reproduction.

The GitHub proof catalogue is `arxiv/hyperbolic_coxeter_dimension_bounds/ADDENDA.md`. It assigns FV1–FV4 to the finite-volume propositions, C1–C2 to the compact case results, and A0–A6 to the arithmetic component, finite-domain and terminal results. Every item identifies its complete statement and proof in the fixed source [8], together with the corresponding checker and execution records. The catalogue and proof maps distinguish written arguments, exact computations and selected conditional Lean kernels. Those kernels do not formalize all geometric and literature inputs.

The availability and replay instructions are in `repro/PUBLIC_DATA.md`; not all separately indexed large inputs are distributed. This edition uses the retained proof and computation records and introduces no new mathematical or Lean execution.

Acknowledgments. The author used OpenAI's GPT-6-astra and GPT-6.1-sol as research and coding assistants to investigate proof ideas, help strengthen the results, improve the exposition, and generate and audit the Python and Lean 4 code accompanying this work. The author reviewed the resulting mathematics and code and takes responsibility for the final content.

References

[1] M. N. Prokhorov. The absence of discrete reflection groups with noncompact fundamental polyhedron of finite volume in Lobachevskii space of large dimension. *Mathematics of the USSR-Izvestiya*, 28:401–411, 1987. `https://www.mathnet.ru/eng/im1484`.

[2] A. G. Khovanskii. Hyperplane sections of polyhedra, toroidal manifolds, and discrete groups in Lobachevskii space. *Functional Analysis and Its Applications*, 20:41–50, 1986. https://www.math.utoronto.ca/askold/1986-Faa-1-english.pdf.

[3] E. B. Vinberg. The absence of crystallographic groups of reflections in Lobachevsky spaces of large dimension. *Trudy Moskov. Mat. Obshch.*, 47:68–102, 1984. English translation, Trans. Moscow Math. Soc. (1985), 75–112. https://www.mathnet.ru/eng/mmo444.

[4] E. B. Vinberg. Hyperbolic reflection groups. *Russian Math. Surveys*, 40(1):31–75, 1985. doi:10.1070/RM1985v040n01ABEH003527.

[5] R. E. Borcherds. Automorphism groups of Lorentzian lattices. *Journal of Algebra*, 111:133–153, 1987. https://math.berkeley.edu/~reb/papers/al/al.pdf.

[6] Lizi Guo and Fangting Zheng. Dimension bounds for finite-volume hyperbolic Coxeter polytopes. arXiv:2611.01795v1, 2026. https://arxiv.org/abs/2611.01795.

[7] Alexander Kolpakov. Corner payments and a dimension bound of 57 for finite-volume hyperbolic Coxeter polytopes. Public repository manuscript, 21 October 2026, 2026. Initial public snapshot 32b68355. https://github.com/sashakolpakov/CoxeterPolytopes/blob/32b68355cdb13ca8629f3fa8b98d424dd70472c7/manuscripts/finite_volume57.tex.

[8] Alexander Kolpakov. Dimension bounds for hyperbolic Coxeter polytopes: complete proofs and addenda. Repository edition of 30 October 2026, snapshot 36eeaddf, 2026. Full manuscript, Parts I–VI. https://github.com/sashakolpakov/CoxeterPolytopes/blob/36eeaddf646eff25d4990c9ab730a20ada21f80f/arxiv/hyperbolic_coxeter_dimension_bounds/main.tex. The matching PDF and reproduction guides are linked from the repository README.

[9] A. Felikson and P. Tumarkin. On simple ideal hyperbolic Coxeter polytopes. arXiv:math/0502413. https://arxiv.org/pdf/math/0502413.

[10] M. Chein. Recherche des graphes des matrices de Coxeter hyperboliques d'ordre ≤ 10, 1969. https://www.numdam.org/item/M2AN_1969__3_3_3_0.pdf.

[11] V. V. Nikulin. Discrete reflection groups in Lobachevsky spaces and algebraic surfaces. In *Proceedings of the International Congress of Mathematicians*, volume 1, pages 654–671, Berkeley, 1986. https://www.mathunion.org/fileadmin/ICM/Proceedings/ICM1986.1/ICM1986.1.ocr.pdf.

[12] V. V. Nikulin. On ground fields of arithmetic hyperbolic reflection groups. In *Groups and Symmetries*, volume 47 of *CRM Proceedings and Lecture Notes*, pages 299–326, 2009. Corrected version arXiv:0708.3991v2. https://arxiv.org/pdf/0708.3991v2.

[13] I. Agol, M. Belolipetsky, P. Storm, and K. Whyte. Finiteness of arithmetic hyperbolic reflection groups. *Groups, Geometry, and Dynamics*, 2:481–498, 2008. https://doi.org/10.4171/GGD/47.

[14] M. Belolipetsky. Arithmetic hyperbolic reflection groups. https://arxiv.org/abs/1506.03111.

[15] V. O. Bugaenko. Groups of automorphisms of unimodular hyperbolic quadratic forms over $\mathbb{Z}[(\sqrt{5}+1)/2]$. *Vestnik Moskov. Univ. Ser. 1 Mat. Mekh.*, pages 6–12, 1984. No. 5. https://www.mathnet.ru/eng/vmumm3408.