Dimension bounds for hyperbolic Coxeter polytopes

Alexander Kolpakov

4 November 2026

Abstract

We prove that a finite-volume hyperbolic Coxeter polytope has dimension at most 57, that a compact one has dimension at most 17, and that a compact arithmetic one has dimension at most 15. The finite-volume argument compares corner weights with the geometry of vertex links. The compact arguments use projections of one vector associated with an original farthest vertex; arithmeticity further restricts the ground field and integral wall parameters. We explain the three proof strategies, prove their central geometric and arithmetic steps, and give precise technical propositions from which the global bounds follow. Complete technical proofs, finite inventories and computational certificates are provided in a versioned GitHub supplement, with a reproduction guide and selected conditional Lean kernels.

1 Introduction

All polytopes in this article are convex, finite-sided and full-dimensional. A hyperbolic Coxeter polytope has dihedral angles that are integer submultiples of π; reflection in its facets generates a discrete group with the polytope as fundamental domain. Hyperbolic space has curvature -1.

Theorem 1.1. *Let $P \subset \mathbb{H}^n$ be a finite-volume hyperbolic Coxeter polytope. Then:*

(i) $n \leq 57$*;*
(ii) *if P is compact, then $n \leq 17$;*
(iii) *if P is compact and its reflection group is arithmetic, then $n \leq 15$.*

The classical finite-volume ceiling is 995, due to Prokhorov [1], using the combinatorial estimates of Khovanskii [2]. Vinberg obtained the compact ceiling 29 [3, 4]. The largest known finite-volume examples have dimension 21 [5]; the bounds here are upper bounds and make no new existence assertion. Compactness is essential in Theorem 1.1(iii).

Guo and Zheng recently obtained the general finite-volume bound 81 [6]. An earlier version of our bound 57, including its corner-weight method and higher-dimensional argument, was publicly posted on 21 October 2026 [7]. The present article condenses the three proof families assembled in the repository edition of 30 October [8].

The methods separate after the common local geometry. In the finite-volume case, triangles and quadrilaterals require corner payments that high-dimensional vertex diagrams cannot accommodate. Nonsimple ideal vertices are treated through the product structure of their links. In the compact case, local walls force quadratic lower bounds for components of a single vector of norm less than one. Arithmeticity enters only in the third argument, through integral cycles and positivity at the other real embeddings.

Detailed proofs and version. The full proof addenda [8] are fixed at repository commit `36eeaddf646eff25d4990c9ab730a20ada21f80f`. Parts I and IV contain the finite-volume argument, Parts II and V the compact argument, and Parts III and VI the arithmetic argument. We refer to these as Addenda FV, C and A, respectively. This article explains the mechanisms of the proofs: how local corner charges become a global curvature contradiction, how actual walls force energy into one fixed vector, and how arithmeticity converts continuous wall parameters into integral data. We give the central arguments and representative calculations, state the technical propositions with their hypotheses, and prove the deductions of the three dimension bounds. Detailed geometric case analyses, finite inventories and computational proofs remain in the supplement. They are essential parts of the proofs, cited at a fixed version independent of subsequent research. Section 6 identifies the current proof catalogue and reproduction/Lean companion.

Contents

1 Introduction **1**

2 Actual walls and vertex diagrams **3**

3 The finite-volume bound **4**
3.1 Actual cores, extensions and whole components 5
3.2 Why the final unit matters 6
3.3 The dimension-specific final local rule 7
3.4 The finite maximum after the transfer 8
3.5 Ideal vertices and positive surplus 9
3.6 Shape curvature and the contradiction in dimension 58 10
3.7 Independent rules in dimensions 59–163 11
3.8 The unbounded tail: actual framing walls and two budgets 12
3.8.1 Outgoing edges give actual framing walls 12
3.8.2 The common-height Lorentz identity 13
3.8.3 Local distances and the original endpoint budgets 14
3.8.4 A narrow facet without a cusp 16

4 The compact bound **17**
4.1 Farthest vertices and projections 17
4.1.1 Positive inverse matrices and feasible projections 18
4.1.2 Opposite walls and common components 19
4.2 Quadratic estimates from reference vectors 20
4.3 Component estimates 21
4.3.1 Ordinary components and exceptional actual defects 21
4.3.2 Common-wall forcing and the short defects 21
4.3.3 The short directions and their exact debits 22
4.3.4 The A-chain branch 22
4.3.5 The common B/D prefix branch 23
4.4 The remaining spherical diagrams 24
4.4.1 A fresh cover in exact dimension 18 24
4.5 The final whole-D18 geometry 25
4.5.1 A future point valid for every dotted head 25
4.5.2 One reference shared by the head and both forks 26

4.5.3 Both signed fork directions . . . 27
4.6 An adjacent A18 vertex . . . 27
4.7 The near/far pair and the final composition . . . 28
4.7.1 Every actual near/far fork edge is short . . . 28
4.7.2 The entire near/far family . . . 28
4.7.3 Exhausting the actual head and fork covers . . . 29

5 The compact arithmetic bound **29**
5.1 The actual arithmetic field . . . 30
5.2 Paying all isolated constraints together . . . 30
5.3 Why the golden field requires a lattice argument . . . 31
5.4 From component bounds to actual endpoint parameters . . . 32
5.5 The four finite algebraic propositions . . . 33
5.5.1 Full quadratic integer rings . . . 34
5.6 The final whole-D16 configuration . . . 35
5.6.1 Using a prefix reference without spending it twice . . . 36
5.6.2 Norm bounds, integer orders and valuations . . . 37
5.7 A worked dyadic branch and the final exclusion . . . 38
5.7.1 Odd norm . . . 38
5.7.2 The remaining dyadic cases . . . 39

6 Proof catalogue and reproduction **40**

2 Actual walls and vertex diagrams

We use a Lorentz space of signature $(-,+,\dots,+)$ and the future unit hyperboloid model of $\mathbb{H}^n$. Outward wall normals are spacelike; the chamber inequalities are $\langle y, e_i\rangle \le 0$. Distinct wall normals have nonpositive products. Finite vertex diagrams are spherical. A horospherical link at an ideal vertex is a product

$$L_v = \prod_i \Delta^{r_i}, \qquad \sum_i r_i = n-1, \tag{2.1}$$

whose factors correspond to irreducible affine diagrams of orders $r_i + 1$. These standard face and link facts are used in their actual-wall form [4, 9].

Two elementary observations account for much of the local geometry.

Lemma 2.1. *A positive-definite Gram matrix with nonpositive off-diagonal entries has entrywise nonnegative inverse. A positive-definite principal set of actual chamber walls has a feasible finite common point.*

Proof. If B is the matrix, choose s larger than its largest eigenvalue. Then $M = sI - B$ is entrywise nonnegative and has spectral radius less than s, so $B^{-1} = s^{-1}\sum_{j\ge 0}(M/s)^j$.

Take a strict interior future timelike point o and put $b_i = -\langle o, e_i\rangle > 0$ for the selected walls. The vector

$$q = o + \sum_i (B^{-1}b)_i e_i$$

is orthogonal to the selected normals and has square $q^2 = o^2 - b^T B^{-1} b < 0$. Its product with o is negative, so it is future pointing. Every other chamber inequality is strict, because the coefficients are nonnegative and all other wall products are nonpositive. Normalize q to the hyperboloid. □

An irreducible affine principal set has a positive kernel vector (u_i). The root combination $\nu = \sum_i u_i e_i$ is null, nonzero since $\langle o, \nu \rangle < 0$, and feasible against every wall. It represents an actual ideal point. Irreducibility and the positive definiteness of proper deletions identify its whole affine factor. In contrast, a kernel vector in an indefinite core represents the zero vector: that vector is null and orthogonal to a timelike vector in the core span, whose perpendicular space is positive definite. Both interpretations are needed when controlling extensions.

At an actual two-face, the supporting walls form a positive-definite diagram of rank $n-2$. Adding one side gives an edge diagram, and adding two adjacent sides gives a positive-semidefinite corner diagram. For opposite sides of a quadrilateral, the corresponding principal diagram is indefinite. A positive or affine alternative would give a feasible common finite or ideal point of those sides. Minimal indefinite subdiagrams, inverse Gram matrices and the classification of finite and affine diagrams consequently reduce the local payment questions to explicit marked configurations.

3 The finite-volume bound

In dimension 58 the proof compares corner payments with the curvature of vertex links. Its improvement comes from transferring one unit away from eight extremal finite forests while preserving the required face payments. Nonsimple cusps contribute a strictly positive surplus. Separate dimension-specific rules, followed by an all-rank narrow-facet argument, exclude the higher dimensions; we do not assume that arbitrary faces inherit Coxeter angles.

We develop these mechanisms and their final deductions below. Catalogue FV1–FV4 in the pinned Addendum FV supplies the exhaustive local classifications, numerical tables and computational proofs at the interfaces where they are used.

For each two-face F, let $s(F)$ be its number of sides, and put

$$\mathcal{D}_5(P) = \sum_{\dim F = 2} (5 - s(F)).$$

A nonnegative corner rule assigns $w(v, F)$ to an actual incidence (v, F); its charge at v is $C(v) = \sum_{F \ni v} w(v, F)$.

Lemma 3.1 (Payment comparison)**.** *Suppose every triangular face receives total payment at least 2 and every quadrilateral receives at least 1. Then*

$$\mathcal{D}_5(P) \le \sum_v C(v).$$

More generally, suppose an incidence argument gives

$$\mathcal{D}_5(P) \ge B v_f + \sum_{v \text{ ideal}} \kappa(v) + \Delta, \qquad \Delta > 0,$$

while a simultaneous payment rule has $C(v) \le B$ at every finite vertex and $C(v) \le \kappa(v) - \varepsilon$ at every ideal vertex, where $\varepsilon \ge 0$. These hypotheses are inconsistent.

Proof. A face with s sides needs payment at least $5 - s$. This is 2 for a triangle and 1 for a quadrilateral. For $s \ge 5$ it follows from nonnegativity. Sum over faces and then over their corners. Combining the resulting upper bound with the proposed lower bound gives $0 \ge \Delta + \varepsilon v_i > 0$. □

3.1 Actual cores, extensions and whole components

We use outward norm-two roots, with diagonal Gram entries 2 and nonpositive off-diagonal entries. At an actual two-face in dimension n, the supporting white roots form a positive-definite rank $n-2$ diagram. White roots together with one black side give a strictly positive-definite edge diagram; adjacent corners are positive semidefinite. A finite corner has an elliptic forest of rank n.

Lemma 2.1 makes these diagrams geometric: a positive principal set has an actual common finite point, strict against all other walls. An irreducible affine set has a positive relation giving a nonzero feasible null vector, hence an actual ideal point. It is a whole cusp factor, since each proper principal subdiagram is positive definite; see also [9, Section 1].

Lemma 3.2 (Opposite sides and minimal cores)**.** *For either opposite pair of sides of an actual quadrilateral, the supporting whites and that pair form an indefinite principal set. An inclusion-minimal indefinite subset contains both marked sides. Its order is at most ten; if all proper principal subsets are elliptic, its order is at most five.*

Proof. A positive set would have an actual common finite point by the projection argument. A positive-semidefinite singular set would have a common ideal point by its affine kernel. Opposite closed sides of a nondegenerate quadrilateral are disjoint, so neither is possible. Deleting either marked side leaves a subset of an actual finite edge diagram. Hence both marks lie in every minimal indefinite subset. The published Lannér/quasi-Lannér order bounds now apply; the classification through order ten is [10]. □

Two opposite cores cannot be orthogonal: their negative vectors would give two negative directions. Their overlap or a joining path therefore supplies an adjacent-side route. The local theorem must restore every other supporting white. All presence conditions refer to connected components of the *whole actual corner diagram*, not merely to the displayed core.

Lemma 3.3 (Two ways to seal a component)**.** *Let G be the Gram matrix of an actual indefinite core.*

1. *A Gram-kernel vector z of this core gives the actual zero root relation $\sum_i z_i e_i = 0$. Every further wall therefore satisfies $\sum_i z_i(e, e_i) = 0$.*

2. *Suppose G is nonsingular, an added forest contacts it only at a position p, and $(G^{-1})_{pp} = r > 1$. If every connected component of the added forest contacts p, its total rank is at most $1/(r-1)$.*

Proof. The span of an indefinite core contains a timelike vector. The root combination associated with a Gram-kernel vector is null and orthogonal to that timelike vector. The perpendicular space is positive definite, so the combination is zero. This differs from the nonzero feasible null vector of an affine block.

For the second statement let B be the positive Gram matrix of all added supporting whites, including indirect tails, and let u be their contact column at p. Schur elimination and the one-negative- direction condition give $B - ruu^T \succeq 0$, whence $u^T B^{-1} u \leq 1/r$.

For a connected square-two finite Coxeter Gram of rank k, every inverse diagonal is at least $k/(k+1)$. This follows by induction: delete the chosen node, let the remaining component ranks be k_j, and use a contact of magnitude at least one into each component. Inverse positivity gives Schur cost at least $\sum_j k_j/(k_j+1) \geq (k-1)/k$. The resulting inverse diagonal is at least $1/(2-(k-1)/k) = k/(k+1)$; the rank-one base is $1/2$. For an added forest of total rank M contacted in each component, the same bound gives $u^T B^{-1} u \geq M/(M+1)$. Thus $M/(M+1) \leq 1/r$, proving $M \leq 1/(r-1)$. □

The Schur argument includes every indirect tail. A radical seal instead pairs its zero relation with every further root: the contact signs force either an already paid shortcut or orthogonality. An indirect extension has a first contacting root, so these tests really control whole components.

At an ideal vertex the link is $\prod_i \Delta^{r_i}$, with $\sum r_i = n-1$. A marked pair in factor i occurs in $V/(r_i+1)$ actual corner fibers, where $V = \prod_i (r_i+1)$: an edge varies in that factor and chooses an arbitrary vertex in every other factor. Thus an affine factor envelope $(r+1)c(r)$ gives the actual charge bound

$$C_{\mathrm{act}} \le C_{\mathrm{up}} = V \sum_i c(r_i). \tag{3.1}$$

The marked pair restores the varying affine factor; other factors appear with one node deleted. Omitting these fiber multiplicities would not count a nonsimple cusp correctly.

The intrinsic tables, including role priorities, are fixed in Addendum FV, "The intrinsic corner weights" (FV1). They specify within-component, cross-component and contextual finite weights and the varying affine-factor weights. The preliminary rule below restores the B_{13} endpoint unit and omits the three rank-15 changes. All later modifications are to this one simultaneous rule.

3.2 Why the final unit matters

The finite calculation ranges over spherical forests of total rank 58. Its alphabet consists of the classical components through that rank, $E_6, E_7, E_8, F_4, H_3, H_4$, and a single cost symbol I_2^* for the remaining dihedral components. The last symbol is justified by the all-label local theorem; it imposes no cutoff on the dihedral angle. Every actual finite vertex lies in this domain. Its formal members need not all be realized by polytopes.

After multiplying charges by 64, the complete cost has the form

$$\begin{aligned} C_{64}(m) = &\sum_T b(T) m_T + \sum_T j(T,T) \binom{m_T}{2} \\ &+ \sum_{T<U} j(T,U) m_T m_U + \Psi(m). \end{aligned} \tag{3.2}$$

The tables b, j give component and cross-component costs; Ψ contains the contextual additions. FV2 fixes context-sensitive counts before an additive recurrence treats the residual rank. Every last-component transition, not just a maximizing path, is checked. The exact tables, complete domains and computational proof are in Addendum FV, "Contexts in the finite-forest calculation." Their geometric identification with actual charges is the separate local theorem FV1.

Proposition 3.4 (Computed finite interface)**.** *The preceding simultaneous rule in ambient 58 pays every triangle at least 2 and every quadrilateral at least 1. Every actual rank 58 finite forest has charge at most 377 except precisely the following eight formal forests: one B_{13} or D_{13} together with any multiset of three A_{15} or D_{15} components. Each of those eight has charge* 378*. All ideal weights admit the envelope of the ideal-vertex discussion below.*

Computed interface FV2. The complete context cover and residual recurrence in FV2 dominate every actual finite vertex. Reconstructing each full contextual cost, including interactions, leaves only the stated eight formal forests above 377; each has cost $78 + 3 \cdot 100 = 378$. The local geometric coverage and triangle proof are supplied by FV1. No realization of these extremal formal forests is asserted. □

Number A_r by its path $0, \dots, r-1$ and B_r by the same path with its last bond doubled. Number D_r by the spine $0, \dots, r-3$ with leaves $r-2, r-1$ at $r-3$; its long endpoint is 0. These

conventions describe actual whole components. The intrinsic weight tables are invariant under their automorphisms.

The separate B_{13} endpoint theorem removes pair $(0, 12)$. Two older affine-B_{13} singleton routes genuinely used that unit; their whole cores instead give distinct already-unit A_1/D_{12} short-leaf contributions. Pair $(1, 12)$, needed by the D_{12} theorem, is retained. This repair, proved in FV1, lowers the four B_{13} extremal forests to 377.

3.3 The dimension-specific final local rule

Theorem 3.5 (Three-rank 15 transfer)**.** *In ambient dimension 58, make these changes to the simultaneous rule:*

1. *In a whole finite D_{13}, give pairs $(0, 11), (0, 12)$ weight $1/2$ instead of 1 only if its actual corner has three distinct whole unmarked A_{15} or D_{15} components.*

2. *Add $1/2$ to whole D_{28} pair $(0, 12)$ only if its corner has two such whole unmarked rank 15 components.*

3. *Add $1/2$ between an endpoint of whole A_{12} and the long endpoint of whole D_{16} under the same two-component presence guard.*

All ideal weights and other finite weights remain. Every triangle still pays at least 2 and every quadrilateral at least 1.

Witnesses are not counted with multiplicity. The negative guard uses $13 + 3 \cdot 15 = 58$ ranks, so the corner has exactly those four components. The two positive guards similarly force exactly $D_{28} + 2 \cdot 15$ or $A_{12} + D_{16} + 2 \cdot 15$. The theorem is not asserted in larger ambient dimensions.

Why the guard survives additional walls. FV1 classifies every affected core and route after restoring all supporting whites. A new white either supplies an already paid short path, violates a projection bound, or is restricted by a zero root relation as in Lemma 3.3. Pairing such a relation with every further wall either supplies a shortcut or seals a small factor incompatible with three rank-15 witnesses.

The remaining growing-factor cases illustrate why a bound on immediate neighbors is insufficient. Their old small factor K has rank $s = 1, 2, 3$, and all additional contacts, after excluding paid shortcuts, meet one old position with inverse square $r = 4/3, 3/2, 2$, respectively. Include *all* added whites in a forest Y, of total rank M, with contact column u. The Lorentz condition gives

$$G_Y - ruu^t \succeq 0, \qquad \frac{M}{M+1} \leq u^t G_Y^{-1} u \leq \frac{1}{r}.$$

Thus $M \leq 3, 2, 1$ and the whole component has rank at most four. This proves the guard impossible in each such case, however long an indirect extension was initially allowed to be. FV1 supplies the finite classification placing every affected route in one of these cases, including the proper completions and all contact signs.

Why an affine route needs a new payment. An older affine-D_{12} route really uses the lowered D_{13} unit, so it requires a repair rather than a retention argument. This case explains the two new half-payments. Number the affine factor by spine $0, \ldots, 8$, leaves $9, 10$ at 0 and $11, 12$ at 8. Take opposite-end marks $a = 9, b = 11$, with the other two black roots c, d. Let Z be the orthogonal elliptic spectator forest, and x, y their nonnegative contact rows on Z.

An adjacent-black bond or shared spectator neighbor is an independently paid short path. Otherwise x, y have disjoint supports. The all-label end-profile reduction in FV1 gives costs $g_c, g_d \le 8$, with a nonextreme end costing at most four, and the exact Schur identity

$$\frac{\det S}{h^2k^2} = 1 - \frac{g_c + g_d}{12} + \frac{1}{12}(x/h - y/k)^t G_Z^{-1}(x/h - y/k),$$

where $h, k > 0$ are its height scales. The actual four-black star has determinant zero. If an end is nonextreme, equality forces costs eight and four and $x = y = 0$. The complete equality cases give a retained short-leaf unit, an independent affine-B_{13} priority, or a second guarded D_{13} half at another corner. Those cases, including the earlier B_{13} endpoint repair, are proved separately in FV1; the latter does not depend on the newly financed D_{13} route.

When both ends are extreme, $h = k = 1/2$ and the identity becomes

$$(x - y)^t G_Z^{-1}(x - y) = 1. \tag{3.3}$$

The affected output lies in the one-sided branch, say $y = 0$; the two-sided branch keeps its short paths or sealed alternatives. If the quiet corner has the new guard, Z consists of exactly three whole A_{15}/D_{15} factors. The following local consequence makes the compensation transparent.

Lemma 3.6. *Under the strict edge conditions and disjoint-support hypothesis, energy one on three whole A_{15}/D_{15} factors forces a single unit contact at the long endpoint of one D_{15}, by one black root. Both black roots are orthogonal to the other two factors.*

Reduction to the profiles in FV1. The inverse diagonals of A_{15} below two are $15/16, 7/4, 7/4, 15/16$ at nodes $0, 1, 13, 14$. In D_{15} only the long endpoint has inverse diagonal below two, namely one. Strict edge positivity and the native contact gap leave the simple contacts at those A_{15} nodes, doubled contacts at its endpoints, and the single unit D_{15} contact. The complete inverse-profile check in FV1 also treats disjoint two-sided differences: every nonzero component energy is at least $15/16$, and no A_{15} profile has energy one. Two active components would cost at least $15/8 > 1$. Equation (3.3) therefore leaves just the stated contact. □

The two inactive factors remain whole: they are orthogonal to the entire star and its active factor. Attaching c to the selected leaf yields, at the opposite corner, whole A_{12} at an endpoint and whole D_{16} at its long endpoint. Attaching it to the mate yields whole D_{28} at $(0, 12)$. The new half in either case is guarded by the two inactive rank-15 factors and lies at a different corner from the quiet D_{13} half. The required unit is therefore restored.

Retention is part of the simultaneous rule. FV1 proves that every older selected output is retained or receives one of these replacements. In particular the changed distance is eleven, while the triangle proof uses distinct short paths of length at most five, paths of length at most seven, or protected E_8 outputs. Those protections and the affine triangle exclusions are part of the complete actual-face proof. Pairwise tests in an abstract corner diagram would not suffice. This finite retention classification, the whole-component seals, and the repaired affine-D_{12} route together prove Theorem 3.5.

3.4 The finite maximum after the transfer

Proposition 3.7. *Every actual finite rank 58 forest under the final simultaneous rule has charge at most 377. The exact formal maximum is 377.*

Proof. The B_{13} theorem lowers four forests in Proposition 3.4 from 378 to 377. The new D_{13} guard lowers the other four by 1. Every other forest was already at most 377, except that the two new positive coupon families must be checked.

Rank capacity makes those families exact. They have respectively

$$170 + 2(100) + \tfrac{1}{2} = 370.5, \qquad 66 + 108.5 + 2(100) + 1 = 375.5.$$

The second total counts both A_{12} endpoint coupons. The previous A_{12}/D_{16} cross cost is zero. Each family has three possible multisets of two A_{15}/D_{15} factors, so all six actual forests are covered. No additional context witness can fit. Outside them the change is pointwise nonincreasing. Thus the bound 377 holds everywhere, and the eight changed forests attain it formally. A geometric realization of an attainer is neither required nor asserted. □

3.5 Ideal vertices and positive surplus

For affine ranks r_i summing to 57 set

$$H(t) = \prod_i (1 + t + \cdots + t^{r_i}) = \sum_{j=0}^{57} h_j t^j, \qquad V = \prod_i (r_i + 1),$$
$$G = \sum_j h_j \min(j, 57 - j), \qquad \kappa = \frac{5G - 57V}{2}.$$

The geometric all-family envelope is unchanged by the finite transfer. Its function c is fixed in FV3 (Addendum FV, "Positive ideal surplus," `fv:sec:ideal`); in particular

$$c(r) = r/2 \quad (1 \le r \le 8), \qquad c(r) = \frac{5r + 56}{r + 1} \quad (24 \le r \le 57).$$

The remaining fifteen rational entries and their factor-charge proofs are supplied there. We need the linear part for a concentration argument, and the full table for one finite calculation. The geometric premise is $C_{\mathrm{act}} \le V \sum_i c(r_i)$ from (3.1); evaluating a polynomial alone would not establish it.

Let $\mathcal{P}(n, 8)$ denote the nondecreasing partitions of n with parts at least eight, including the empty partition when $n = 0$. Define the terminal profiles

$$\mathcal{T}_{57} = \bigcup_{\ell=0}^{7} \{(\ell, p) : p \in \mathcal{P}(57 - \ell, 8)\}, \tag{3.4}$$

omitting a zero part. For any profile set $m = \kappa/V - \sum_i c(r_i)$.

Proposition 3.8 (Complete terminal calculation)**.** *The complete domain* (3.4) *has 2552 profiles. Exact polynomial convolution and rational evaluation give*

$$\min_{\mathcal{T}_{57}} m = \frac{87}{10240} > 0, \qquad \min_{\mathcal{T}_{57}} Vm = 36.$$

Computational proof in FV3. Sorted-partition recursion lowers the remaining rank at each step; the unique first part and its sorted tail establish completeness and absence of duplicates. Repeated convolution with $1 + t + \cdots + t^r$ gives every coefficient of H, and hence V, G and each rational margin. The certificate and its checker cover every profile and coefficient list before taking the two minima. The full algorithms, correctness proofs and recorded outputs are in Addendum FV, `fv:prop:terminal`, identified by FV3. The normalized and absolute minima are distinct assertions; they need not have the same attainer. □

The reason this finite domain controls arbitrarily many small simplex factors is convex order. Moving one rank from a smaller factor to a larger one spreads the link's index distribution, which reduces the curvature available to pay for corners.

Lemma 3.9 (Concentrating two simplex ranks)**.** *Let $1 \le a \le b$ be integers, and let U_r be uniform on $\{0, \dots, r\}$. With independent summands, $U_{a-1} + U_{b+1}$ dominates $U_a + U_b$ in convex order. Adding any common independent summand preserves this order.*

Proof. Both sums have mean $(a+b)/2$ and are symmetric about that mean. For an integer threshold t, let Δ_t be the difference of their expectations of $(X - t)_+$. Equal means allow us to use $(t - X)_+$ instead, and symmetry gives $\Delta_t = \Delta_{a+b-t}$. It suffices to take $0 \le t \le (a+b)/2$. Put $D = (a+1)(b+1)$ and $D' = a(b+2)$. For $0 \le t \le a$, direct summation of the increasing side of the two convolution triangles gives

$$\Delta_t = \binom{t+2}{3}\left(\frac{1}{D'} - \frac{1}{D}\right) \ge 0, \qquad D - D' = b - a + 1 > 0.$$

For $a \le t \le (a+b)/2 \le b$, summing over the first uniform instead gives

$$\Delta_t = \frac{t(a+b-t) - (a-1)(a+2b+2)/3}{2(b+1)(b+2)}.$$

The numerator is increasing up to $(a+b)/2$. At $t = a$ it is $(a+2)(b-a+1)/3 > 0$. Outside the common support the difference is zero; between integer thresholds it is linear. Thus all stop-loss differences are nonnegative. On the integer support the values of a convex function are an affine function plus a nonnegative sum of these stop-loss functions, proving convex order. Conditioning on a common independent summand proves the last assertion. □

Lemma 3.10 (Concentration below rank eight)**.** *Every actual ideal vertex has $\kappa - C_{\mathrm{act}} \ge 36$.*

Proof. For $1 \le a \le b \le 7$, replacing (a, b) by $(a-1, b+1)$ preserves normalized envelope cost, since all these ranks stay at most eight. Lemma 3.9, including its independent-summand assertion, gives the required convex order. Since

$$\frac{G}{V} = \frac{57 - \mathbb{E}|2S - 57|}{2},$$

the transfer does not increase normalized curvature or normalized margin. Its product decreases from $(a+1)(b+1)$ to $a(b+2)$. Iteration terminates at ranks at least 8 and at most one positive rank below 8. Indeed the integer potential $\sum_i r_i^2$ increases by $2(b-a+1) \ge 2$ at every step and is bounded above by 57^2. Its initial value is at least 57, so at most 1596 steps occur. Rank zero is omitted. No transfer passes through the changed rank 9 envelope. The stopping domain is exactly (3.4).

Proposition 3.8 gives positive normalized terminal margin and absolute envelope surplus at least 36. If the original and terminal quantities are (V, m) and (V', m'), then $V \ge V'$ and $m \ge m' > 0$. Thus $Vm \ge V'm' \ge 36$. This multiplication uses positivity. Finally (3.1) transfers the same lower bound to actual cusp charge. □

3.6 Shape curvature and the contradiction in dimension 58

Lemma 3.11. *For a proposed ambient 58 polytope,*

$$\mathcal{D}_5 \ge 377 v_f + \sum_{v \text{ ideal}} \kappa(v) + 4205.$$

Proof. Choose a generic linear function on the bounded Klein closure of the polytope. Let E_v be the number of edges of its simple vertex figure and E_v^0 the number not cut by the induced level section. If I counts all polygon vertices with multiplicity, the published incidence identities give $I = \sum_v E_v$ and $2f_2 = \sum_v E_v^0$ [2, Theorem 9]. Hence $\mathcal{D}_5 = \sum_v (5E_v^0/2 - E_v)$.

Orient each vertex figure by its induced generic linear function. Assign an uncut edge below the section to its upper endpoint and one above the section to its lower endpoint. A vertex of index j receives j or $57 - j$ contributions, at least their minimum. The index distribution is its h-vector [2, Theorems 2 and 6], so $E_v^0 \geq G_v$. At the distinct global minimum and maximum the whole vertex figure is uncut, and symmetry gives

$$E_v - G_v = \sum_{j>57/2} h_j(2j - 57) \geq \sum_{j=29}^{57} (2j - 57) = 841.$$

Every coefficient of a simplex or product-of-simplices h-polynomial is at least 1. Adding the two extrema and multiplying by $5/2$ gives 4205, counted once. For a finite vertex $h_j = 1$, so

$$G_0 = \sum_{j=0}^{57} \min(j, 57 - j) = 812, \qquad \frac{5G_0 - 57 \cdot 58}{2} = 377.$$

The remaining ideal terms are precisely κ. □

For faces with at least five sides,$5 - s(F) \leq 0$. The nonnegative local rule and Proposition 3.7 therefore give

$$\mathcal{D}_5 \leq 377 v_f + \sum C_{\text{act}}.$$

Subtract Lemma 3.11 and use Lemma 3.10:

$$0 \geq 4205 + \sum (\kappa - C_{\text{act}}) \geq 4205 + 36 v_i > 0.$$

This also excludes the all-finite case. Thus dimension 58 is impossible.

It remains to exclude dimensions at least 59. We now give the separate arguments used for that purpose.

3.7 Independent rules in dimensions 59–163

The same comparison applies at other ranks, but the rule and its finite maximum must be established in that actual dimension. Put $d = n - 1$ and

$$\beta_n = \begin{cases} n(n-6)/8 & n \text{ even}, \\ (n-1)(n-5)/8 & n \text{ odd}. \end{cases}$$

The incidence proof above gives

$$\mathcal{D}_5 \geq \beta_n v_f + \sum_{v \text{ ideal}} \kappa(v) + \Delta_n, \qquad \Delta_n \geq 5(n-1) > 0.$$

For a cusp profile $\boldsymbol{r}$ of total rank d, define H, V, G as before with 57 replaced by d. An affine envelope c gives the normalized margin

$$m_c(\boldsymbol{r}) = \frac{5G - dV}{2V} - \sum_i c(r_i). \tag{3.5}$$

A simultaneous rule with finite maximum $U_n \leq \beta_n$ and $m_c > 0$ at every cusp therefore excludes that dimension.

Proposition 3.12 (The bounded-range interfaces; FV4)**.** *For each* $59 \le n \le 163$*, the dimension-specific rules in Addendum FV pay two on every triangle and one on every quadrilateral, with whole-component guards valid against all additional supporting walls. Their finite forest maxima satisfy* $U_n \le \beta_n$*, and their affine-factor envelopes have positive margin* (3.5) *on every terminal profile. The families cover* 59–65*,* 66–97 *and* 98–163*. Their stated concentration moves carry the positive margin to every actual product-of-simplices cusp.*

The refined rules for the first two ranges, their exact forest domains and complete terminal profiles are supplied under "The complete ambient 59–65 layer table" and "Labelled rules for dimensions 66–97" in Addendum FV. This is a finite conjunction of results at the specified dimensions, not an extrapolation from dimension 58 or from a range endpoint.

The higher ranges illustrate the same mechanism more simply:

n	U_n	$c(r)$
98 : 108	$9n$	$\min(8, r/2)$
109 : 116	$9n$	$\min(9, r/2)$
117 : 126	$10n$	$\min(10, r/2)$
127 : 136	$11n$	$\min(11, r/2)$
137 : 146	$12n$	$\min(12, r/2)$
147 : 156	$13n$	$\min(13, r/2)$
157 : 163	$14n$	$\min(14, r/2)$.

For $k = 9, \dots, 13$ these rules pay a unit at distance at most k or in a whole component of order at most 17; at $k = 14$ distance alone suffices. The first range instead uses finite distance nine and affine distance eight, retaining certain distance-nine pairs near branches or nonsimple bonds. FV4 proves local routing and the affine pair counts for these choices; the latter acquire the product-link multiplicities above.

For $c(r) = \min(r/2, L/2)$, concentration leaves arbitrary copies of L, at most one part below L, and at most one part above L. Indeed each permitted transfer raises the integer sum of squared ranks, decreases the product of the link vertex counts, and does not increase (3.5). The process terminates because the total rank is fixed. The full terminal calculations in FV4 give strictly positive minima for every row; moreover $\beta_n > U_n$ in these seven ranges. Together with the two refined families this proves the exclusion of every $59 \le n \le 163$.

3.8 The unbounded tail: actual framing walls and two budgets

The range $n \ge 164$ requires a proof at all ranks, not a finite list of partitions. We use Nikulin's narrow-facet theorem [11, pp. 654–655]; its unrestricted statement is also given in [12, Section 4, p. 10]. A finite-volume chamber has a facet F whose actual framing hyperplanes have pairwise distance strictly less than $\operatorname{arccosh} 7$. With outward normals of square 2 this implies

$$-(x, y) < 14 \tag{3.6}$$

for any two framing walls. This is the external narrowness input. We prove the incidence, endpoint and scale statements needed to apply it; they are not additional hypotheses on the cusp.

3.8.1 Outgoing edges give actual framing walls

Suppose first that F contains a cusp v. Its horospherical link is a product of affine Coxeter simplices. Write their ranks r_a, with $\sum_a r_a = N = n - 1$, and normalize every old root to square 2. In each

factor choose two omitted nodes in advance. In the factor containing the normal α_s of F, both choices avoid s. The two choices need not be distinct. If that factor has affine rank one, use its other node twice; its modified target $2(r_a - 1)/9$ is zero. Deleting one node per factor gives a positive-definite support S of N roots and a vertex of the horospherical link. It therefore defines an actual outgoing edge in F. Finite volume gives a finite or ideal second endpoint.

At a finite endpoint choose any new wall x terminating that edge. It and α_s belong to a spherical diagram, so their rank-two Gram is positive definite. At an ideal endpoint let T_b be its affine factors. The positive-definite support S contains at most $\operatorname{rank} T_b$ nodes from each. The total rank N forces equality in every factor: exactly one node of each T_b is missing. If the factor containing α_s has rank at least 2, choose its missing node as x; every pair in such a factor has finite angle. If it is affine A_1, choose the missing node of another factor, which is orthogonal to α_s. Another factor exists in the range at issue. Thus again the pair is positive definite.

To see that this is an actual ridge, start with a strict interior timelike point z. Project it off the span of x, α_s. The projection correction is a nonnegative combination of those normals, because their Gram is a positive-definite matrix with nonpositive off-diagonal entries and nonnegative inverse. Every other wall inequality stays strict, and the projected square is even more negative. It gives a point of precisely their common face. Hence x is in the framing set controlled by (3.6). Apply the same construction to the second omission tuple and call its terminating wall y.

Neither x nor y is an old deleted cusp wall: that wall is strict along the outgoing edge away from v. Their old contact rows $p_j = -(x, \alpha_j)$ and $q_j = -(y, \alpha_j)$ are nonnegative, and every nonzero entry is at least 1. For each affine factor let $c_a = \sum_j \delta_{aj}\alpha_j$, with $\delta_{aj} > 0$ normalized by $\min_j \delta_{aj} = 1$. These are nonzero future null vectors, since their pairing with an interior point is negative. They are all positive multiples $k_a c$ of one cusp vector c. Therefore

$$\sigma_a = \delta_a \cdot p = -(x, c_a), \quad \tau_a = \delta_a \cdot q = -(y, c_a), \quad q_0 = \sigma_a/\tau_a$$

have a common positive ratio. Indeed, if a row height vanished, all its nonnegative contacts with that factor would vanish; proportionality of the c_a would force every factor height and every old contact to vanish. A square-two normal would then lie in the null orthogonal complement of the full cusp span, impossible.

For each endpoint the principal Gram $S+x$ or $S+y$ is positive semidefinite. Its Schur complement gives the separate budgets

$$\sum_a C_{0a} \le 2, \qquad \sum_a C_{1a} \le 2, \qquad C_{\ell a} = p^t_{\ell,S_a} G(S_a)^{-1} p_{\ell,S_a}. \tag{3.7}$$

The inverse is entrywise nonnegative. No bound on a deleted-node contact or on its Coxeter label has been used.

3.8.2 The common-height Lorentz identity

Choose a null partner d_0 with $(c, d_0) = -1$, and write $x = ac + bd_0 + u$, $y = a'c + b'd_0 + u'$ in the orthogonal Euclidean quotient. Then $b = -(x, c) > 0$, $b' = -(y, c) > 0$. Since $(x/b - y/b') \perp c$ and $x^2 = y^2 = 2$,

$$\|u/b - u'/b'\|^2 = 2/b^2 + 2/(b')^2 - 2(x, y)/(bb').$$

The quotient is the orthogonal sum of the affine factor spaces. In factor a put $E_a = \|u_a/\sigma_a - u'_a/\tau_a\|^2$. Since $\sigma_a = k_a b$ and $\tau_a = k_a b'$, the k_a^2 cancels and the preceding identity becomes

$$-(x, y) = \frac{1}{2}\sum_a \sigma_a \tau_a E_a - q_0 - q_0^{-1}. \tag{3.8}$$

This proves the scale cancellation for a product cusp; no commensurability or chosen horosphere is required. Let $M = \max(q_0, q_0^{-1})$ and $W_a = \min(\sigma_a, \tau_a)^2 E_a$. Then $\sigma_a \tau_a E_a = M W_a$ and $q_0 + q_0^{-1} \le 2M$.

The prescribed omission choices have targets

$$T_a = 2r_a/9 \quad \text{ordinarily}, \qquad T_a = 2(r_a - 1)/9 \quad \text{in the sole avoid-node factor.}$$

Thus $\sum T_a = (2N-2)/9$. We next prove

$$\sum_a W_a \ge \begin{cases} (2N-2)/9 - 4 & 1 \le M \le 21/20, \\ (2N-2)/9 - 16/3 & M \ge 21/20. \end{cases} \tag{3.9}$$

It follows from(3.8) that $-(x,y) \ge M(N-37)/9$ in the first range and $-(x,y) \ge M(N-43)/9$ in the second. At $N = 163 + t$ these are at least $14 + t/9$ and $14 + 7t/60$, respectively. They contradict (3.6) for every $t \ge 0$.

3.8.3 Local distances and the original endpoint budgets

Each affine diagram has Euclidean point potentials v_j: their squared distances $R(i,j)$ are the corresponding grounded inverse-Gram quantities. Normalizing a contact row by its affine height gives nonnegative weights of sum one on these potentials. Thus the local energy E is the squared distance between two convex means. An omitted-node contact is included in the mean; there is no bound on its magnitude.

For illustration, an A cycle of order m has

$$\delta_j = 1, \qquad R(i,j) = h(m-h)/m, \qquad h = (i-j) \bmod m.$$

The ordinary omissions are $0, \lfloor m/2 \rfloor$. Rotation avoids a prescribed node without weakening the estimate; at $m = 2$ the repeated allowed omission has modified target zero. For B, C, D, partial sums of orthonormal coordinates give the potentials and all-rank inequalities. Their complete formulas, the exceptional grounded matrices, and the prescribed ordinary and avoid-node choices are fixed in Addendum FV, "Point potentials and the complete local domain." Those specific choices are used throughout the following interfaces.

Lemma 3.13 (Full-budget classical unit charge)**.** *For the prescribed ordinary or avoid-node omissions of a classical affine factor, let $C_0, C_1 \le 2$ be its actual retained costs and $W = \min(\sigma, \tau)^2 E$. If the common height ratio is at most $21/20$, then*

$$W + C_0 + C_1 \ge T, \qquad T = 2r/9 \text{ or } 2(r-1)/9,$$

with the second target in the avoid-node factor. This includes arbitrary real contacts whose nonzero magnitudes are at least one, and both possibilities for each omitted-node contact.

Geometric reduction and the local certificates in FV4. For omissions i, j, write the two point means as $v_i + a, v_j + b$, with contact weights w_k, z_l. Polarization gives

$$\begin{aligned} E = R(i,j) &- \sum_k w_k \{R(i,j) + R(i,k) - R(k,j)\} \\ &- \sum_l z_l \{R(i,j) + R(j,l) - R(l,i)\} + \|a - b\|^2. \end{aligned} \tag{3.10}$$

The omitted-node displacement is zero. Every full row height is at least one, so $W \geq E$. If $c_{ik} = \delta_k^2 R(i,k)$, inverse positivity and the contact gap give

$$C_0 \geq \sum_{k \text{ retained, contacted}} c_{ik}, \qquad C_1 \geq \sum_{l \text{ retained, contacted}} c_{jl}.$$

These unit-contact costs therefore control the linear losses without a bound on the full contact magnitudes. For B, C, D the coordinate proof partitions contacts into those whose cost already exceeds T, and those whose linear loss is at most their unit cost. The complete partition, including the modified B_2 case, is in the full-budget lemma of FV4.

Here is the all-rank part for the A cycles. If $m \geq 18$, a retained contact at cyclic distance at least three costs more than two, since $3(m-3)/m > 2$ and resistance increases to the antipode. The remaining distances one and two have linear-loss to cost ratio at most $(m+\varepsilon)/(m-2)$, where $\varepsilon = m \bmod 2$. Using the two original budgets in the polarization identity leaves

$$W + C_0 + C_1 - T \geq R(0, \lfloor m/2 \rfloor) - T - \frac{4(2+\varepsilon)}{m-2}.$$

For $m = 18 + 2u$ its right side is $(128 + 84u + 4u^2)/(36(m-2))$; for $m = 19 + 2u$ it is $(360 + 1708u + 252u^2 + 8u^3)/(36m(m-2))$. Both are positive for every $u \geq 0$. The remaining orders $2 \leq m \leq 17$ are the finite convex-quadratic domain of FV4. Its certificates cover every retained support, both omitted-node states and the common-height ratio constraint. In particular this interface holds at the original budgets two, before selecting any failing factor. □

Why two ordered budgets must remain separate. A factor can fail the coefficient-one estimate only in a small exceptional domain. Removing it changes the two endpoint budgets by different amounts. Replacing them by their sum would lose the restriction that makes the remaining loss affordable. The exact local support classification and residual calculation used here are the following two finite interfaces from FV4.

Proposition 3.14 (Exceptional losses at the original budgets)**.** *For the prescribed omissions and $1 \leq M \leq 21/20$, a factor with $T - W > C_0 + C_1$ is either an ordinary E_8, with cost floors $(8/9, 2/3)$ and target $16/9$, or an ordinary E_7/modified E_8, with floors $(1/2, 1)$ or $(1, 1/2)$ and target $14/9$. After such a factor is removed, the maximum residual loss is at most $20/9$ under ordered budgets $(10/9, 4/3)$, and at most $22/9$ under $(1, 3/2)$ or $(3/2, 1)$. The residual bounds permit at most one modified factor.*

The classification is applied at the original budgets two: Lemma 3.13 first excludes classical failures, and the exceptional proof checks every cheap grounded support, with all inverse cross terms and both omitted-node states. In particular its ordinary E_7 two-contact case requires four distinct ordered full-support keys. Only after this classification may the smaller residual budgets be used.

The residual proof is an ordered packing calculation. Every local domain is dominated by one of 28 records with no larger ordered cost floors, no smaller loss bound and the same modified flag. A recurrence maximizes the sum of recorded losses subject to each remaining endpoint budget and at most one modified record. Each record consumes at least $1/2$ of total budget, so this recursion has finite depth. FV4 supplies the domination proof, continuous support certificates and every recurrence inequality giving the stated maxima. The records are necessary envelopes, not asserted simultaneous geometric realizations.

If no factor fails, summing its coefficient-one inequalities bounds the total loss by four. Otherwise an E_8 failure leaves $(10/9, 4/3)$, so its own loss at most $16/9$ and residual loss at most $20/9$ again

sum to four. The other failures give $14/9 + 22/9 = 4$. All further failures belong to the residual calculation. Even if the removed factor was modified, permitting one modified residual factor only increases the bound. This proves the first line of (3.9) without assuming a residual budget before the first failure.

The other height range. FV4 also proves, for the same omissions and the full actual contact domain,

$$W + \frac{4}{3}(C_0 + C_1) \geq T.$$

For B, C, D the coordinate coefficient-one proof already uses only that both heights are at least one. For A, write $C = C_0 + C_1$. If $C \geq 3T/4$, nonnegativity of W suffices. Below that threshold the contact gap and original budgets restrict the relevant cyclic distances to one and two; in the remaining range polarization leaves the nonnegative margin $d(m - 4d - 3\varepsilon)/(3m)$. The small orders and exceptional factors are the separately specified finite convex-quadratic domains in FV4. Their certificates include the omitted-node possibilities and every cheap support. Thus the inequality covers all ranks, with no cutoff on a contact magnitude. Summing it and using (3.7) proves the second line of (3.9).

3.8.4 A narrow facet without a cusp

The alternative is a compact facet. We use its actual ambient faces, without assuming its induced angles are Coxeter. A compact two-face has positive-definite corner diagrams. For a triangle, a minimal indefinite star contains all three black nodes and at most seven whites. A spanning tree joining the marks supplies two distinct corner paths of length at most eight. For a quadrilateral, the whites with either opposite pair are indefinite: a positive or affine block would give an actual finite or ideal intersection of the opposite sides. Minimal opposite cores each have at most eight whites and cannot be orthogonal. A shortest path in their overlapping or adjacent union therefore joins consecutive black sides through at most 17 edges. The complete core argument is also recorded in FV4.

Consequently unit weight at ambient spherical distance at most 17 pays two on a compact triangle and one on a compact quadrilateral. Its cost at an ambient finite vertex is at most $17n$: components of rank at most 35 have at most $\binom{r}{2} \leq 17r$ pairs; larger spherical components are paths or D forks, with at most r pairs at each fixed distance.

A compact facet of dimension $d = n - 1$ is simple. Its two-face corners form a subset of these ambient vertex pairs, so their total charge is at most $17(d+1)f_0$. Its symmetric nonnegative h-vector has $h_0 = h_d = 1$. Pairing the coefficients $\binom{k}{2}$ and $\binom{d-k}{2}$ gives

$$\frac{f_2}{f_0} > \begin{cases} d(d-2)/8 & d \text{ even}, \\ (d-1)^2/8 & d \text{ odd}. \end{cases}$$

The endpoint coefficients make the inequality strict. Its average required payment $5f_2/f_0 - \binom{d}{2}$ therefore exceeds $d(d-6)/8$ or $(d-1)(d-5)/8$. Subtracting the available $17(d+1)$ gives, at $d = 143 + 2t$ and $144 + 2t$, respectively,

$$t^2/2 + 36t + 3/2, \qquad t^2/2 + 73t/2 + 19,$$

both positive for $t \geq 0$. Thus a compact facet forces $n \leq 143$. The cusp and compact alternatives exhaust the finite-volume facet supplied by the narrow-facet theorem.

Proposition 3.15 (Independent higher-dimensional exclusion)**.** *There is no finite-volume hyperbolic Coxeter polytope in any dimension $n \geq 59$.*

Proof. The dimension-specific families exclude $59 \leq n \leq 163$. For $n \geq 164$, the framing-wall and two-budget argument gives $-(x, y) \geq 14$, contradicting narrowness, if the narrow facet has a cusp. If it has none, the compact-facet comparison already excludes $n \geq 144$. These alternatives exhaust the facet supplied by the narrow-facet theorem. □

Together with the contradiction in dimension 58, this proves Theorem 1.1(i).

4 The compact bound

The compact argument turns the geometry at a farthest vertex into a Euclidean budget. Each whole irreducible component consumes part of one vector of squared norm less than one. The main difficulty is obtaining simultaneous lower bounds: pressures from several actual walls may share coordinates, and their best separate estimates cannot simply be added. We first prove the projection and reference-vector principles that handle this dependence, then explain the component reduction and the final whole-D_{18} mechanism.

Catalogue C1 supplies the complete component inventories and proper-row proofs, and C2 supplies the terminal whole-component and higher-rank cases. References with a prefix B below denote exact numbered proofs in Part V of the pinned complete edition, called Addendum C. We retain these references at the points where their full cases enter the argument.

Theorem 4.1 (Compact dimension bound)**.** *Every compact hyperbolic Coxeter polytope has dimension at most* 17*.*

4.1 Farthest vertices and projections

Use the hyperboloid model with Lorentz signature $(- + \cdots +)$. Every wall has an outward unit spacelike normal n, and the polytope is given by $\langle r, n\rangle \leq 0$. Distinct actual wall normals have nonpositive products. Fix a strict interior point z, choose an actual vertex p farthest from z, and write

$$z = E(p - x), \qquad E = -\langle z, p\rangle > 0, \qquad x \in p^{\perp}, \qquad \|x\|^2 = 1 - E^{-2} < 1. \tag{4.1}$$

The space $p^{\perp}$ is Euclidean. The incident normals at p form a positive definite spherical diagram; compact acute simplicity implies that there are exactly n of them in dimension n. The whole irreducible components T have orthogonal projections with

$$\sum_T e_T = \|x\|^2 < 1, \qquad e_T = \|x_T\|^2. \tag{4.2}$$

We retain these original normals and this original choice of p, z, x, E throughout an energy argument.

The function $r \mapsto -\langle z, r\rangle$ takes its maximum on a compact hyperbolic polytope at a vertex. Its restriction to a unit speed geodesic has positive second derivative equal to the function itself, so a maximum on a segment occurs at an endpoint; successively passing to proper faces proves the assertion. Consequently

$$-\langle z, r\rangle \leq E \quad \text{for every actual point } r \in P. \tag{4.3}$$

At an adjacent vertex q, write $z = E_q(q - y)$. Equation (4.3) gives a comparison factor E/E_q, not a claim that q is farthest.

4.1.1 Positive inverse matrices and feasible projections

Inverse positivity is Lemma 2.1. We record its explicit projection formula in the coordinates used below.

Lemma 4.2 (An actual positive projection)**.** *Let n_j, $j \in J$, be actual wall normals with positive definite Gram matrix B_J, and let z be strict interior. Put*

$$q_0 = z - \sum_{i,j\in J} (B_J^{-1})_{ij}\langle z, n_j\rangle n_i.$$

Then q_0 is future timelike, orthogonal to all selected walls, and has negative product with every unselected actual wall. Its normalization is a point of the original polytope.

Proof. The added coefficients are nonnegative by Lemma 2.1. For an unselected wall every added product is nonpositive and the original product is negative; the selected products vanish. Moreover

$$q_0^2 = z^2 - (\langle z, n_i\rangle)_{i\in J}^T B_J^{-1}(\langle z, n_i\rangle)_{i\in J} < 0.$$

Since $\langle z, q_0\rangle = q_0^2 < 0$, it has future orientation. All original wall inequalities survive normalization. If fewer than n walls are selected this is a point on a face, not an asserted full-dimensional vertex. □

The same sign check applies to positive combinations of actual normals below: selected products are calculated explicitly; every unknown original wall has nonpositive product with every summand. A nonzero feasible future null vector would give a point of the ideal boundary of P, contradicting compactness. A positive affine Coxeter kernel on actual walls is an obstruction after its future sign and all remaining wall products have been checked.

Lemma 4.3 (Future-point pressure)**.** *Suppose $U \in p^\perp$, $\xi = Ap - U$ is future timelike with $\xi^2 = -k < 0$, and its normalization is feasible against every actual wall. Then the same original vector satisfies*

$$\langle x, U\rangle \geq A - \sqrt{k}.$$

At a different actual vertex q, use the decomposition $\xi = Aq - U$ with $U \in q^\perp$. With all-point comparison factor κ, the corresponding inequality is $\langle y, U\rangle \geq A - \kappa\sqrt{k}$.

Proof. The spatial hypothesis gives $A = -\langle p, \xi\rangle > 0$. Apply (4.3) to $\xi/\sqrt{k}$: $-\langle z, \xi/\sqrt{k}\rangle = E(A - \langle x, U\rangle)/\sqrt{k} \leq E$. The modified comparison is the identical calculation. □

For any Euclidean subspace $S \subset p^\perp$ containing U, let x_S be the orthogonal projection of x onto S. If $U \neq 0$, then $\|U\|^2 = A^2 - k$; Cauchy therefore gives

$$\|x_S\|^2 \geq \frac{A - \sqrt{k}}{A + \sqrt{k}}. \tag{4.4}$$

The case $U = 0$ gives the same nonnegative bound directly. Normalize a future vector, not a spacelike correction. For a cap $c = Ap - v$, clear a positive definite old block J:

$$w = \sum_{i,j\in J} (B_J^{-1})_{ij}\beta_j n_i, \quad \beta_j = -\langle c, n_j\rangle \geq 0, \quad \eta = c + w, \quad \delta = \eta^2 = 1 - \beta^T B_J^{-1}\beta.$$

If $\delta < 0$, η is the future vector to normalize. If $\delta > 0$, the actual finite projection instead uses

$$p + \frac{A}{\delta}\eta, \qquad \left(p + \frac{A}{\delta}\eta\right)^2 = -1 - \frac{A^2}{\delta}.$$

When its selected products are nonpositive, its positive coefficients check all unselected walls. The pressures are

$$\langle x, v - w\rangle \geq A - \sqrt{-\delta}, \qquad \langle x, v - w\rangle \geq \frac{A}{1 + \sqrt{\delta/(A^2 + \delta)}}. \tag{4.5}$$

Old normals lie in $p^\perp$, so $v - w$ denotes the spatial part of the corresponding displayed Lorentz vector.

4.1.2 Opposite walls and common components

Let B be the original incident Gram matrix, $C = B^{-1}$, and

$$\omega_i = \sum_j C_{ij} n_j, \qquad d_i = C_{ii}, \qquad u_i = \omega_i/\sqrt{d_i}.$$

An actual opposite wall has normal

$$c = Ap - v, \quad A > 0, \quad v = \sum_i \beta_i \omega_i, \quad \beta_i = -\langle c, n_i\rangle \geq 0, \quad A^2 = \beta^T C \beta - 1.$$

Strict interiority gives

$$\langle x, v\rangle < A. \tag{4.6}$$

Along the original edge in direction $-u_i$, of length L_i, the original farthest comparison gives

$$\langle x, u_i\rangle \geq \tanh(L_i/2). \tag{4.7}$$

Whole-component projection energies sum to $\|x\|^2$.

Proposition 4.4 (Reciprocal shadows; internal geometric interface)**.** *For an actual cap put* $T_i = (C\beta)_i/(A\sqrt{d_i})$*. If* $T_i > 1$*, it is the opposite wall of edge* i*, and* $\tanh L_i = 1/T_i$*. If it is not that opposite wall, then* $\tanh L_i \geq T_i$*. The value* $T_i = 1$ *is incompatible with a finite endpoint. For distinct actual opposite walls,*

$$\tanh L_i \tanh L_j \geq K_{ij}, \qquad K_{ij} = C_{ij}/\sqrt{d_i d_j}. \tag{4.8}$$

Opposite walls of orthogonal original directions cannot coincide.

Proof. The Schur identity says that $C - C_{\cdot i}C_{i\cdot}/d_i$ vanishes on row and column i; elsewhere it is the inverse of the retained old Gram matrix. It is entrywise nonnegative. Let $d = Dp - \sum_j \gamma_j \omega_j$ be the actual opposite of edge i, distinct from c. Acuteness and the Schur identity give

$$AD \geq \beta^T C \gamma \geq (C\beta)_i (C\gamma)_i/d_i.$$

The actual endpoint equation is $\tanh L_i = D\sqrt{d_i}/(C\gamma)_i$, so $\tanh L_i \geq T_i$. If $T_i > 1$ this alternative is impossible, forcing $c = d$; its endpoint equation gives $\tanh L_i = 1/T_i$. If $T_i = 1$, the distinct alternative is impossible and the same-wall equation would give an ideal endpoint. This is excluded by compactness. For the pair inequality, first use the contacts of the actual opposite of i to get $T_j \geq K_{ij}/\tanh L_i$; the distinct alternative at edge j yields the assertion.

Finally suppose the same wall closes orthogonal directions i, j. Clearing every other old normal adds positive coefficients and gives

$$c_F = Ap - \frac{A}{r_i} u_i - \frac{A}{r_j} u_j, \qquad r_i = \tanh L_i, \quad r_j = \tanh L_j.$$

Its product with the strict interior point remains negative, so $\langle x, u_i\rangle/r_i + \langle x, u_j\rangle/r_j < 1$. But (4.7) makes this sum at least $1 + (\tau_i^2 + \tau_j^2)/2 > 1$, where $\tau_i = \tanh(L_i/2)$. This contradiction proves the orthogonal assertion. □

The displayed pair inequality concerns full tangents, not half tangents. For an opposite wall, its retained foreign component can have at most one contact: two would make a cycle in its actual finite replacement. A positive foreign cost is $q = \beta_j^2 C_{jj} \geq 1/4$. In a nontrivial component $C_{jj} \geq 4/3$, so the only cost below $1/3$ is a simple 3-contact on an A_1, of cost $1/4$. Moreover

$$q < A^2. \tag{4.9}$$

Otherwise the actual reciprocal test would close that foreign direction too, violating the orthogonal assertion, with the equality case giving a null endpoint. Set $F(t) = 2t/(1+t^2)$. For unit directions of correlation k in a component of energy at most e, positive heights satisfy

$$\tau_i + \tau_j \leq \sqrt{2e(1+k)}, \qquad F(\tau_i)F(\tau_j) \leq \frac{4s}{(1+s)^2}, \quad s = \frac{e(1+k)}{2}. \tag{4.10}$$

The second inequality follows from concavity of $\log F$ on $(0,1)$. If its upper bound is smaller than K_{ij}, Proposition 4.4 forces a common actual opposite. Its finite-white extension or positive affine kernel then gives additional actual pressures.

4.2 Quadratic estimates from reference vectors

The elementary estimate behind every certificate is

$$\|x\|^2 \geq 2\langle x, W\rangle - \|W\|^2. \tag{4.11}$$

Every term in $\langle x, W\rangle$ must be justified. Write $w = \sum_i b_i P_i$, $b_i \geq 0$, with actual pressures $\langle x, P_i\rangle > a_i$, and demand $A_w = \sum_i b_i a_i$. For a new actual pressure $\langle x, Z\rangle \geq h$, choose

$$Q = Z - \sum_i t_i P_i, \qquad W = w + \mu Q.$$

Before using pressures require $\mu \geq 0$ and check every $b_i - \mu t_i \geq 0$. Then

$$\langle x, W\rangle \geq A_w + \mu(h - \sum_i t_i a_i), \quad \|W\|^2 = \|w\|^2 + 2\mu\langle w, Q\rangle + \mu^2\|Q\|^2. \tag{4.12}$$

The available price b_i is individual to anchor i. The first inequality is strict if some strictly constrained old anchor retains a positive coefficient, or if $\mu > 0$ and the new pressure is strict. For multiple corrections every new weight must be nonnegative; their combined debit at each anchor and all mutual cross terms must be retained.

For $P_i = e_1 + \cdots + e_i$, increasing concave interpolation through $(0,0)$ and actual floors $\langle x, P_i\rangle > f_i$ gives a reference with successive slopes as its coordinates. Its coefficients in the P_i are nonnegative slope decreases, and its demand equals its square. Deleting a knot between intervals of lengths a, b and slopes s_1, s_2 loses $ab(s_1 - s_2)^2/(a+b)$. For unit adjacent spacings this is half the squared slope difference. For the $\sqrt{i}-$constant floors at $i \geq 5$, such a unit-neighbor loss is $< 1/3200$; nonadjacent omissions have disjoint intervals and their losses add. End omissions use their exact chord costs. An omitted floor has coefficient zero.

References are Euclidean vectors: actual feasible points prove the pressures, and nonnegative coefficients combine them in (4.11).

4.3 Component estimates

4.3.1 Ordinary components and exceptional actual defects

All bounds below concern the same original projection, with arbitrary finite spectators.

Component	Energy greater than	Component	Energy greater than
ordinary nonshort A_1	1/10	ordinary A_2	319/2000
ordinary $I_2(m), m \geq 4$	1/7	A_3	1/4
B_3, H_3	2/7	A_4	2/7
B_4, D_4, F_4	1/3	H_4	11/36
A_5	3/8	A_6	3/7
B_5, B_6	3/7	D_5, D_6	5/14, 3/7
B_7, D_7	1/2	B_8, D_8	4/7
E_6, E_7, E_8	9/20, 5/9, 2/3		

An actual isolated dotted short A_1 is denoted I. A second short-wall profile J uses two old A_1 factors whose joint energy exceeds .34. The two typed rank-two exceptions, each of energy $> .09$, are

$$N_7 = (\text{old } A_2, \text{ new isolated native 7}), \qquad O_7 = (\text{old } I_2(7), \text{ new isolated native 3}).$$

A bare $I_2(7)$ is not automatically O_7. There is at most one short I/J profile or one typed exception; they do not coexist. Acuteness of their actual isolated walls is essential: two on orthogonal supports would have product $-\sqrt{h_1 h_2}$ strictly between $-1/2$ and 0. Appendix B.1 proves the endpoint estimates, the J case and the ordinary rates.

Useful joint bounds, counting I only once, are

$$I + A_1 > 1/5, \quad I + 2A_1 > 3/10, \quad I + T_2 > 21/100, \quad I + T_3 > 2/7. \tag{4.13}$$

Here T_j is ordinary irreducible of rank j. Both typed exceptions force another T_2 to cost $> 1/5$. Another A_1 costs $> 1/5$ beside N_7 and $> 2/15$ beside O_7. For the latter, isolated height $h = \frac{1}{4}\csc^2(\pi/7) - 1 < 1/3$ gives squared full tangent greater than 3/7. Squared half tangent exceeds $2/15$, since $4(2/15)/(1+2/15)^2 = 120/289 < 3/7$. The complete contact and defect proofs, including the actual-wall height constraints behind (4.13), are given in Appendix B.1. The uniform actual B/D floor is proved in Appendix B.2.

4.3.2 Common-wall forcing and the short defects

For the following small-component calculations, let T be the actual whole component, $C_T = B_T^{-1}$ its inverse Gram matrix, and β_T the cap's contact vector on T. Write

$$q = \beta_T^T C_T \beta_T, \qquad Q = \sum_{U \neq T} \beta_U^T C_U \beta_U, \qquad h = A^2 = q + Q - 1 > 0.$$

Thus q denotes the own-component cost and Q the full foreign cost.

Lemma 4.5 (Common-wall forcing and isolation)**.** *Suppose the energy of a component is at most e, with $0 \leq e < 1$. A pair for which the right side of* (4.10) *is below K_{ij} has one actual opposite. If one wall closes every direction of the component, then, with $v_i = (C\beta)_i/\sqrt{d_i}$ and $R_e^2 = 4e/(1+e)^2$,*

$$q - 1 \leq h \leq R_e^2 \min_i v_i^2. \tag{4.14}$$

If $h < 1/4$, it has no foreign contact. In a path, a wall closing every direction can have multiple contacts only at the two endpoints; in a D_4 star it has at most one.

Proof. The first assertion is (4.10). For a closed direction, $\tanh L_i = A/v_i$ and $\tau_i^2 \le e$, which gives (4.14). Every nonzero foreign cost is at least $1/4$ and is strictly below h by (4.9); hence the isolation assertion. If the old path between two contacts omits a direction closed by the wall, deleting that direction leaves a cycle in an actual finite replacement. This is impossible. The same argument applies to every two contacts in a D_4 star. Finally, a single-contact own cost $q = 1$ gives an affine principal set, impossible even in the presence of further spectators. □

4.3.3 The short directions and their exact debits

Call an A_1 direction short when $\tanh^2 L < 1/3$. For a short A_1 opposite put $b = \beta^2$, $h = A^2$, and let Q be its full foreign cost. Then

$$h = b + Q - 1, \qquad h < b/3, \qquad Q < 1 - 2b/3.$$

Equation (4.9) rules out two foreign components. With one, its cost must be the sole value below $1/3$, namely an A_1 contact 3 of cost $1/4$. Thus the only possibilities are

$$I:\ \text{isolated dotted } A_1, \quad 0 < h < 1/2; \qquad J:\ \text{dotted } A_1 + A_{1,3}, \quad 1/3 < h < 3/8.$$

For completeness, with one foreign component, $Q < h = b + Q - 1$ forces $b > 1$, and $Q < 1/3$ then forces $Q = 1/4$. For J, the heavy edge contributes more than $A/2$ to $x \cdot v$. The light shadow contributes at least $A(1 - \sqrt{1 - 1/(4h)})$; this is at least $A/2$ if $h \le 1/3$. Strict (4.6) therefore gives the asserted lower bound. Native acuteness excludes two short walls. Indeed their height product is below $1/2$, so their product must be zero. Disjoint supports give a negative product, heavy overlaps give a spatial product above $1/2$, and a sole light/light overlap of two J walls gives $1/4$, below their height product. None is zero. A J heavy direction has $\tanh^2 L > 4/13$, hence energy $> 9/100$; its light direction has $\tanh^2 L > 2/3 > 16/25$, hence energy $> 1/4$. Thus J funds both A_1 factors above their combined ordinary baseline $2/10$. Every nonshort direction pays at least $5 - 2\sqrt{6} > 1/10$, since $\sqrt{6} < 49/20$. We may conservatively discard the one short direction, but only the isolated type I can create an unfunded debit.

The rank-two exceptions are

old block / new isolated contact	$h = A^2$	energy
$A_2/7$	$(4/3)\cos^2(\pi/7) - 1 \in (0, 1/12)$	$> 9/100$
$I_2(7)/3$	$(1/4)\csc^2(\pi/7) - 1 \in (0, 1/3)$	$> 9/100.$

(4.15)

Their spatial supports lie entirely in their own blocks. Two defects in different blocks, or a defect and a short A_1, would have actual wall product $-\sqrt{h_1 h_2} \in (-1/2, 0)$. That interval is forbidden: distinct acute unit walls have product 0 or at most $-1/2$. Therefore

$$\text{at most one short } A_1, \quad \text{or one rank-two defect, but never both.} \tag{4.16}$$

This argument is independent of every other component type.

4.3.4 The A-chain branch

For A_r, use the zero-sum subspace of $\mathbb{R}^{r+1}$ with roots $(e_i - e_{i+1})/\sqrt{2}$. The inverse Gram matrix is

$$C_{ij} = \frac{2\min(i,j)(r + 1 - \max(i,j))}{r+1}.$$

A finite-white classification of actual opposite walls gives strong dual floors except for a short list of literal helper caps. Adjacent weak helpers are excluded using their actual mutual product and a positive two-cap projection. Concave interpolation uses only strong floors; helper pressures are paid at surviving anchors. The small end variance yields stronger component bounds:

r	$r \geq 7$	$r \geq 8$	$r \geq 13$	16	17
$e(A_r) >$	1/2	4/7	5/7	199/250	6/7

$e(A_{18}) > 91/100.$

Later local refinements give $e(A_{12}) > 13/20$ and $e(A_{15}) > 413/500$, uniformly with arbitrary spectators. The shared actual feet, common-pair inventories, uniform $1/2, 4/7, 5/7$ bounds, and the $N = 17, 18, 19$ central-window and paid-reference arguments are proved in Appendix B.3. The newer $A_{16} > .86$ proof in particular implies the older .796 bound without requiring its separate historical reference. The fresh smaller-rank proofs are in Sections B.4.5 and B.4.4. Their full native-cosine inventories and individual actual-helper payments are given there.

4.3.5 The common B/D prefix branch

For B_r, use long roots $(e_i - e_{i+1})/\sqrt{2}$, $i < r$, and short root e_r. For D_r, use the long chain and the fork root $(e_{r-1} + e_r)/\sqrt{2}$. On a long initial segment, $\omega_i = \sqrt{2}P_i$, $C_{ij} = 2\min(i, j)$. In type D, $x_1 > \cdots > x_{r-1} > |x_r|$; the last coordinate is signed and the forks are distinct actual duals.

Proposition 4.6 (Generic prefix theorem)**.** *Let $6 \leq M \leq 16$. Suppose an actual component is B_r, $r \geq M + 5$, or D_r, $r \geq M + 6$. Put*

$$f_i = \sqrt{i} - \frac{61}{100\sqrt{2}}, \quad E_M = \frac{f_5^2}{5} + \sum_{i=5}^{M-1} (\sqrt{i+1} - \sqrt{i})^2, \quad H_M = \left\lfloor \frac{M-5}{2} \right\rfloor.$$

With arbitrary finite spectators, $\sum_{j=1}^{M+1} x_j^2 > E_M - H_M/3200$.

Proof. Appendix B.5 proves that a weak site $5 \leq i \leq M + 1$ has a pure cap

$$v = \gamma\omega_i + \omega_1/2, \quad A^2 = 2i\gamma^2 + 2\gamma - \tfrac{1}{2}, \quad 101/125 < \gamma < 6/5,$$

with own label 5, finite label at least 7, or dotted own contact, and no spectator contact. Actual weak sites are nonadjacent. This is proved at these ranks, not transferred from a whole-component exclusion. If 5 is strong, choose a strong $K = M$ or $M + 1$. Interpolate through actual strong knots. Each missing internal floor loses $< 1/3200$; at $K = M + 1$ the extra increment $(\sqrt{M+1} - \sqrt{M})^2 > 1/(4M + 2) \geq 1/66$ pays the possible extra omission.

If 5 is weak, the first-six-root point table in Appendix B.5 gives $x_1 > .5965$, and 6 is strong, with $S_6 > 2.01815$. Interpolate through $(1, .5965), (6, 2.01815)$ and a later strong endpoint. The slopes decrease: $.5965, .28433$, then $1/(\sqrt{K} + \sqrt{6}) < 1/5$. Its energy exceeds

$$L_M = .7600299945 + \frac{(\sqrt{M} - \sqrt{6})^2}{M - 6},$$

the last term being zero at $M = 6$. Now $E_6 < .7$. The $k = M - 6$ increments from 6 onward lie in $(1/8, 1/5)$, so their variance is at most $k(1/5 - 1/8)^2/4 \leq 9/640$. Thus $L_M - E_M > .76 - .7 - 9/640 > 0$, directly also at $M = 6$. This proves the weak-5 branch without first assuming strong 5. □

Appendix B.5 supplies the complete finite-white inventory, zero-own repair, actual two-cap nonadjacency, all four weak-5 pressures and continuous seals, with every spectator retained. Its original rank inequalities are part of the theorem. A short-tail continuation at an excluded rank still requires its own new local proof.

4.4 The remaining spherical diagrams

Proposition 4.7 (Compact rank cover and its consumers)**.** *At an original farthest vertex of a compact Coxeter polytope of dimension* 18*, the complete necessary cover is the one displayed below. Its proper-component rows and its whole A_{18} and B_{18} rows are excluded by the indicated case proofs. The higher-dimensional exclusions are separate statements at their actual ranks.*

The accounting leading to the cover is given below, followed by exact locations of the longer row proofs. None of the retained rows asserts a realized polytope.

4.4.1 A fresh cover in exact dimension 18

Call A_r, $r \geq 7$, or B_r, D_r, $r \geq 9$, dominant. Each costs $> 1/2$, so at most one is possible. With none the ordinary rank-over-14 baseline, even allowing the unique defect, exceeds 1. Exactly one is therefore present. The fresh total-rank-18 necessary cover is:

Dominant	All retained spectator alternatives
A_{12}	$I + (A_5$ or $D_5)$, or $N_7/O_7 + T_4$
A_{15}	ordinary A_3; $T_2 + A_1$; $I + T_2$; or $O_7 + A_1$
A_{16}	ordinary T_2; two ordinary A_1; $I + A_1$; or N_7/O_7
A_{17}	one ordinary nonshort A_1, or I
A_{18}	none
B_{16}	N_7/O_7
B_{17}	I
B_{18}	none
D_{16}	N_7/O_7
D_{17}	one ordinary nonshort A_1, or I
D_{18}	none

Here ordinary $T_2 = I_2(m)$, $m \geq 3$, and $T_4 \in \{A_4, B_4, D_4, F_4, H_4\}$. H_4 and native 5 are not discarded by field arguments.

The reduction uses a finite rank-partition argument, with each energy bound valid at its actual component rank and with arbitrary spectators. For example, $A_8, \ldots, A_{11}$ cost more than $4/7$ and leave enough ordinary spectator ranks to exceed the common budget. A single short or typed defect is counted once through the joint bounds, rather than being repeated in overlapping sums. The B/D ranks require their own short-tail inventories before a generic prefix estimate can be extended to them. These are the distinctions enforced by the full C1 forest proof.

The first retained rows illustrate how a strengthened component theorem is used without changing the forest domain:

$$A_{12}+I+(A_5/D_5) > 13/20+5/14 = 141/140, \qquad A_{12}+N_7/O_7+T_4 > 13/20+.09+2/7 = 359/350.$$

The $413/500$ bound likewise closes every A_{15} row: ordinary total rank three costs $> 17/70$, $I+T_2 > .21$, and the O_7+A_1 row exceeds $787/750$. The remaining $A_{16}, A_{17}, B_{16}, B_{17}, D_{16}, D_{17}$ rows have the complete spectator-specific consumers in Addendum C, Sections B.3.5 and B.10.1–B.10.6. Whole A_{18} and whole B_{18} are treated in B.7 and B.8. Those proofs include the zero-own, finite, parallel and continuous dotted alternatives and all additional walls; the finite cover itself does not exclude a retained row.

Theorem 4.8 (Separate higher-dimensional exclusion)**.** *No compact hyperbolic Coxeter polytope has dimension at least* 19*.*

Theorem 4.8 is the conjunction of the actual-rank exclusions $19, \dots, 29$ in Addendum C, B.11–B.14, and Vinberg's published ceiling 29 above that range. In rank 19 the whole A_{19}, B_{19}, D_{19} cases have distinct endpoint and head proofs. C2 identifies their complete composition in B.14.22. This is independent of the exclusion of rank 18; it does not infer a Coxeter structure on an arbitrary face or delete a root from another rank's cover.

4.5 The final whole-D_{18} geometry

Proposition 4.9. *An original farthest vertex cannot have whole incident diagram D_{18}.*

The component cover and C1's complete row consumers leave this case. We give the continuous dotted-head and near/far mechanism, stating the remaining local inventory and pair exclusions from C2 exactly where used. We treat the final dotted-head case using the complete inventory proved in Appendix B.9. Keep the original p, z, x and use actual old roots

$$n_i = (e_i - e_{i+1})/\sqrt{2} \quad (1 \le i \le 17), \qquad n_{18} = (e_{17} + e_{18})/\sqrt{2}.$$

Write

$$P_i = e_1 + \cdots + e_i, \quad \Omega_i = \sqrt{2}P_i \ (i \le 16), \quad \Omega_{17} = (P_{17} - e_{18})/\sqrt{2}, \quad \Omega_{18} = (P_{17} + e_{18})/\sqrt{2}.$$

In particular x_{18} is signed and $x_1 > \cdots > x_{17} > |x_{18}|$. In this part only, set

$$S_i = \langle x, P_i \rangle, \qquad f_i = \sqrt{i} - \frac{.58}{\sqrt{2}}. \tag{4.17}$$

All terminating decimals in this paper denote exact rational numbers.

The head is the actual opposite wall of the long-end edge 1. Section B.9.1 proves that its normal is

$$H = A_H p - \eta\Omega_1, \qquad A_H^2 = 2\eta^2 - 1 > 0.$$

The same section gives the complete own-contact list. Sections B.9.8 and B.9.9 prove the strong terminal floor $S_{16} > f_{16}$, its funded reference append and the complete fork-role list. The finite-head cases close in Section B.9.22. We treat every dotted head $\eta > 1$ without a parameter cutoff.

Appendix B.9 proves these earlier inputs in geometric dependency order: actual head isolation, the fresh threshold window, strong 16, finite-head closure and the every-head pair exclusions. In particular it proves the changed-threshold strong-site statements afresh rather than inheriting them unconditionally.

4.5.1 A future point valid for every dotted head

Put $u = \eta^{-2} \in (0, 1)$, and consider

$$\Xi = H + n_1/\eta.$$

This is a positive combination of two actual normals. Its products with H, n_1 are respectively 0 and $-\eta + 1/\eta < 0$, and $\Xi^2 = -(1-u) < 0$. Its time is $A_H > 0$. Every other actual wall has nonpositive product with each summand. Hence its normalization is an actual future point of the original polytope, with no unknown wall omitted. Lemma 4.3 gives

$$\langle x, Z_u \rangle \ge h(u), \qquad Z_u = \frac{(2-u)e_1 + ue_2}{2}, \quad h(u) = \frac{\sqrt{2-u} - \sqrt{u(1-u)}}{\sqrt{2}}. \tag{4.18}$$

Indeed the spatial part of Ξ, divided by $\sqrt{2}\eta$, is Z_u. We will use

$$\sum_j (Z_u)_j = 1, \qquad R(u) = \|Z_u\|^2 = 1 - u + u^2/2.$$

Equation (4.18) is a point pressure, not an energy lower bound, and is valid on the entire open interval.

4.5.2 One reference shared by the head and both forks

The complete changed-threshold inventory (C2, B.9.2) leaves four initial rows: strong 5; an isolated dotted weak 5 followed by strong $J = 6$ or 7; or a pure far-left-three helper at 5. All other initial roles are strong or already excluded by actual positive cores. The reference lemmas B.9.4, B.9.8 and B.9.20 give the following quantified data.

Their common conclusion is the following reference contract: after incorporating the actual head pressure, each initial row has one positive pressure decomposition $x \cdot W > A_W$ such that

$$2A_W - \|W\|^2 > .992, \quad \operatorname{supp} W \subset \{1, \dots, 16\}, \quad W \cdot P_{16} = f_{16}, \quad \beta_{16} > .127 \text{ unused.} \tag{4.19}$$

Here A_W is the sum of its funded lower pressures; β_{16} is the retained individual coefficient of P_{16}. In particular the contract retains the available endpoint price, not merely a norm bound. The missing knots have coefficient zero, and all their chord losses are included before any price is asserted. This is C2's uniform reference lemma, using B.9.4, B.9.8 and B.9.20 together with Part II, "Bounds for the head and fork edges".

In the strong-five row its unheaded reference w is flat on the first five coordinates, has $x \cdot w > A_w$ and

$$2A_w - \|w\|^2 > .9565188, \qquad b < .3652, \qquad \beta_5 > .151.$$

It already has the endpoint sum and unused endpoint coefficient in (4.19). Put $Q_H = Z_u - P_5/5$. Then $w \cdot Q_H = 0$ and $\|Q_H\|^2 = R(u) - 1/5$.

Here is a complete example of the subsequent funding calculation. In the strong-five row choose weight $\mu = .33$ on Q_H. The debit $.33/5 = .066$ fits the individual price .151, and $w \cdot Q_H = 0$. Hence one reference gives

$$2A_{w+.33Q_H} - \|w + .33Q_H\|^2 > F(u) := .7372668 + .66h(u) - .1089R(u).$$

Put $a = \sqrt{2-u}$ and $b = \sqrt{u(1-u)}$. Differentiation gives

$$h''(u) = \frac{1/(4b^3) - 1/(4a^3)}{\sqrt{2}} > \frac{7}{4\sqrt{2}}, \qquad F''(u) > .7 \quad (0 < u < 1).$$

At $u_0 = 2/3$,

$$h(u_0) = \sqrt{2/3} - 1/3, \quad R(u_0) = 5/9, \quad F(u_0) > .9956502,$$
$$F'(u_0) = .2013 - .0825\sqrt{6}, \qquad |F'(u_0)| < .0008.$$

These inequalities follow from $.81649^2 < 2/3$ and $2.4494 < \sqrt{6} < 2.4495$. Strong convexity and completion of the square therefore give

$$F(u) > .9956502 - \frac{.0008^2}{1.4} > .99564 \qquad (0 < u < 1).$$

This seals the entire dotted-head interval, without a sampled minimum.

The other three rows use the same funding principle. The isolated-five rows spend the combined head and isolated-foot debit at their surviving P_J anchor and retain the mutual cross term. The helper row pays two separate anchors; replacing them by one pooled price would be invalid. Their exact coefficients, raw reference losses and continuous scalar seals are the remaining parts of the uniform reference lemma in C2. All corrections have sum zero and support at most 7, so the terminal coefficient in (4.19) is still available for either genuine fork. We now use the quantified contract in all four rows.

4.5.3 Both signed fork directions

Let $\Delta = S_{16} - f_{16} \geq 0$. The pressure decomposition of W keeps its original terminal coefficient. Therefore $\langle x, W\rangle > A_W + \beta_{16}\Delta$, and the same norm identity gives

$$\|x - W\|^2 < .008 - .254\Delta, \qquad x_{17}^2 + x_{18}^2 < .008 - .254\Delta. \tag{4.20}$$

The two true unit fork duals are

$$\frac{\Omega_f}{3} = \frac{P_{17} \pm e_{18}}{\sqrt{18}}, \qquad f = 17, 18.$$

For either actual fork height,

$$\begin{aligned} t_f := \langle x, \Omega_f/3\rangle &< \frac{f_{16} + \Delta + \sqrt{.016 - .508\Delta}}{\sqrt{18}} \\ &< \frac{3.59 + .1265}{\sqrt{18}} < .876. \end{aligned} \tag{4.21}$$

The numerator decreases with Δ: its derivative is $1 - .254/\sqrt{.016 - .508\Delta} < 0$ on its nonempty domain. The endpoint is handled by continuity. The seals are $f_{16} < 3.59$, $\sqrt{.016} < .1265$, $\sqrt{18} > 4.2426$, and $.876 \cdot 4.2426 = 3.7165176 > 3.7165$. No absolute value has replaced the signed last coordinate.

For an actual fork edge with distance cosine C, (4.7) gives $t_f \geq \sqrt{(C-1)/(C+1)}$. Hence

$$C < \frac{1 + .876^2}{1 - .876^2} < 7.6, \tag{4.22}$$

since $.876^2 = .767376$ and $7.6 \cdot .232624 = 1.7679424 > 1.767376$. In Section B.9.9, every far-next-three role has $g \geq \cos(\pi/5) > .809$ and exact cosine $C = 9g + 1 > 8.281$. Equation (4.22) removes that entire role, including all its previously classified native/null own alternatives.

4.6 An adjacent A_{18} vertex

Lemma 4.10. *Let q be an actual vertex of a compact Coxeter 18-polytope with whole incident diagram A_{18}. Write $z = E_q(q - y)$, $\|y\| < 1$. Suppose $-\langle z, r\rangle \leq \kappa E_q$ for every actual $r \in P$, where $\kappa < 11/8$. Then the height in its positive normalized endpoint-dual direction cannot exceed .78.*

Here q is not assumed farthest. The all-point comparison is an explicit hypothesis; the original x budget is never added to the local y budget.

The exact local inventory at this comparison is C2, Addendum C, B.4. It is proved before using any endpoint height. In the zero-sum model in $\mathbb{R}^{19}$ it leaves the following weak sites $5 \leq i \leq 14$ for the floors $y \cdot \omega_i > \sqrt{2i(19-i)/19} - .87$: near-four at the four literal side-order-five/six positions,

or pure far-end-three helpers of own coefficient $\varphi/2 \le \gamma < 17/16$. The latter are nonadjacent, and all isolated caps are strong. The full inventory includes zero, every finite, parallel and unbounded dotted own, and proves all-wall feasibility of the auxiliary points. These are necessary geometric premises of the local argument.

The proof has two geometric stages. If the endpoint height exceeded .78, each weak near-four would supply a genuine mate vertex, obtained by positive finite Gram inversion while retaining its actual cap. The selected products are checked explicitly and inverse positivity preserves every unselected original wall, including the reverse wall at q. The mate has a high normalized height and a small correlation with the endpoint direction. A Cauchy estimate excludes left near-fours; one positive combination of the endpoint, the right mate and an independent strong direction excludes right near-fours. These comparisons all use the stated factor $\kappa < 11/8$.

Only nonadjacent pure helpers then remain weak. Concave interpolation through the genuine strong floors builds a zero-sum reference in $\mathbb{R}^{19}$. An omitted weak knot is replaced by a chord, has coefficient zero, and contributes its exact loss. The two endpoint patterns have separate surviving individual anchor prices. Finally the presumed endpoint height pays a positive correction whose negative part debits only the appropriate available anchor; its cross term with the base reference is zero. The same local-y square exceeds one in both patterns. Catalogue C2 supplies both parts of the complete proof: Addendum C, Appendix B.4 proves the local inventory and helper nonadjacency; Part II, "A local A_{18} lemma at an adjacent actual vertex", gives the decisive .78 endpoint calculation, mate correlations, rational interpolation heights, chord losses and individual coefficient prices. Those technical calculations are essential to the lemma and are retained there.

4.7 The near/far pair and the final composition

4.7.1 Every actual near/far fork edge is short

For a near-three or far-three fork role, the complete finite-white classification in Section B.9.9 gives an actual other endpoint q with whole diagram A_{18}. The positive reverse direction at q is exactly its normalized new endpoint dual: p has nonzero product with only that new incident root. Let C be the original edge cosine, $\beta = \sqrt{C^2-1}/C$, and $t = t_f < .876$.

Suppose $C \ge 5.6$. Then

$$E_q/E = C - \sqrt{C^2-1}\,t > 5.6 - .876(5.51) = .77324.$$

The function increases for $C \ge 5.6$. Here $5.51^2 = 30.3601 > 30.36$ and $(11/8)(.77324) > 1$, so $E/E_q < 11/8$. Also $\beta > .9839$, because $31.36(1 - .9839^2) = 1.0016631744 > 1$. The positive reverse height satisfies

$$\langle y, u_{18}\rangle = \frac{\beta - t}{1 - \beta t} > .78.$$

Its denominator is positive and its limiting gap is

$$.9839 - .876 - .78(1 - .9839 \cdot .876) = .000179192 > 0.$$

The original maximum supplies all-point comparison at q. Lemma 4.10 excludes $C \ge 5.6$. Every actual near/far fork edge therefore has $C < 5.6$.

4.7.2 The entire near/far family

Suppose the actual fork opposites are near and far:

$$c_N = Np - \gamma\Omega_f - \Omega_o/2, \qquad c_F = Fp - \delta\Omega_o - \Omega_1/2,$$

with f, o the different actual forks. Their times and exact edge cosines are

$$N^2 = 9\gamma^2 + 8\gamma + 5/4, \quad F^2 = 9\delta^2 + \delta - 1/2, \quad C_N = \frac{18\gamma + 8}{\sqrt{19}}, \quad C_F = \frac{18\delta + 1}{\sqrt{19}}.$$

The short-edge bounds and $\sqrt{19} < 4.359$ give $\gamma < .912, \delta < 1.301$. Put $U = 3\gamma + 4/3, V = 3\delta + 1/6, a^2 = 19/36$. Then $N^2 = U^2 - a^2, F^2 = V^2 - a^2$, and

$$(UV - a^2)^2 - N^2F^2 = a^2(U - V)^2 \geq 0, \qquad UV - a^2 > 0.$$

The actual mutual magnitude consequently satisfies

$$\begin{aligned} \lambda = -\langle c_N, c_F \rangle &= NF - 8\gamma\delta - \gamma/2 - (9/2)\delta - 1/4 \\ &\leq \delta(\gamma - 1/2) - 5/9 \\ &< 1.301(.412) - 5/9 = .536012 - 5/9 < 0. \end{aligned}$$

This contradicts all-wall acuteness. If $\gamma \leq 1/2$ the upper bound is already negative; the accepted own list in fact has $\gamma \geq \cos(\pi/4) > 1/2$. Zero, every finite, parallel and unbounded dotted mutual are excluded at once, for all finite or dotted near/far owns.

4.7.3 Exhausting the actual head and fork covers

For a dotted head, (4.22) removes every far-next-three role from Section B.9.9. What remains is isolated, near-three, or far-three. Their six unordered pairs are exhausted by

Actual pair	Complete excluding argument
isolated/isolated	Sections B.9.14 and B.9.15
isolated/near	Section B.9.16
isolated/far	Sections B.9.17 and B.9.18
near/near	Section B.9.9
near/far	The complete argument above
far/far	Section B.9.18

Both assignments of the genuine signed forks are included. Every pair is impossible, so no dotted head remains.

The head cover also includes own zero, finite and parallel cases. Section B.9.1 removes own $0, 3, 4$ by time constraints and own 6 by a positive affine $[6, 3]$ kernel. At own 6 an actual null is $\sqrt{3}H + 2n_1 + n_2$; at parallel own it is $H + n_1$. The remaining finite labels $5, 7, 8, 9$ and every integer label at least 10 are excluded by the complete conjunction in Section B.9.22. Every dotted own has just been excluded. Thus the whole-D_{18} diagram is impossible.

Theorem 4.8 excludes every dimension at least 19. At exact dimension 18, the fresh complete cover and its row consumers, including the whole-B_{18} proof in Addendum C, B.8, leave only whole D_{18} at the same original farthest vertex. The head/fork argument excludes it. Hence dimension 18 is impossible, proving Theorem 4.1. Each step uses its own actual-rank cover and the same original vector within a case; no dimension heredity is used.

5 The compact arithmetic bound

The compact theorem gives $n \leq 17$. Arithmeticity supplies two further restrictions on the same actual walls: integral cyclic products, and positive definite conjugate forms at every nonphysical

embedding of the actual ground field. The proof uses those restrictions first to strengthen component bounds, then to reduce the remaining endpoint parameters to finite algebraic domains or a short list of dyadic possibilities.

The main ideas are the common-vector absorption of isolated constraints, actual-field descent, invariant-lattice recognition in the golden field, and the interaction of dyadic divisibility with geometric pressure. The complete component and field-case inventories remain in the pinned Addendum A, catalogue A0–A6. We state their precise interfaces before using them and work out the decisive algebraic and geometric mechanisms.

5.1 The actual arithmetic field

An arithmetic reflection lattice has an admissible quadratic-form model over its actual totally real field K; the structural reflection theorem is recorded as Lemma 5.2 of [13]. It supplies K-valued root lines and positive definite nonphysical forms. For roots r_i on those lines put

$$a_{ij} = 2(r_i, r_j)/(r_i, r_i).$$

Invariant cyclic products are algebraic integers and generate the actual ground field. Individual physical unit roots need not be K-valued. Disconnected roots need not have equal raw norm squareclasses.

If an actual principal Gram matrix, after its specified permitted K-rescaling, is defined over $k \subset K$ and indefinite physically, then $K = k$: an additional real embedding fixing k would preserve indefiniteness and contradict positive definiteness. A nonzero algebraic integer also has nonzero integral norm, so bounds at *every* nonphysical embedding can force a degree bound. Neither argument assumes a convenient overfield or an integer power basis.

The rational field is excluded in the dimensions used here: compactness requires anisotropy, while Meyer's theorem makes an indefinite rational form in at least five variables isotropic. These arithmetic inputs are reviewed in [14]. Finite/affine classification and acute wall geometry are external foundations; component estimates, forest reductions and computed propositions below are internal results.

For reference, the original adjacent-edge and completed-square inequalities are

$$x \cdot u \geq \tanh(L/2), \qquad \|x\|^2 \geq 2x \cdot w - \|w\|^2. \tag{5.1}$$

The first concerns a genuine edge and the second any one reference w. Every change of reference below records its individual available coefficients before these inequalities are applied.

5.2 Paying all isolated constraints together

The arithmetic component estimates use a common analytic step. It explains why overlapping local pressures can be combined without optimizing different vectors independently.

Lemma 5.1 (Simultaneous absorption)**.** *Let n_i be incident unit normals, ω_i their duals, and $w = \sum_i \beta_i \omega_i$ with $\beta_i = \langle w, n_i \rangle \geq 0$. Suppose $E_w = \|w\|^2 = \sum_i \beta_i H_i$. Outside a set J, assume $\langle x, \omega_i \rangle \geq H_i$. For $i \in J$ suppose $\beta_i > 0$, $0 < u_i < 1$, and*

$$\langle x, \omega_i - u_i n_i \rangle \geq H_i - \delta_i, \qquad 2\delta_i < \beta_i(2u_i - u_i^2).$$

Then $\|x\|^2 \geq E_w$, strictly if J is nonempty or some positively weighted retained pressure is strict.

Proof. Use the single vector $Z = w - \sum_{i\in J} \beta_i u_i n_i$. The pressures and $\langle n_i, n_j\rangle \le 0$ for $i \ne j$ give

$$\langle x, Z\rangle \ge E_w - \sum_{i\in J} \beta_i \delta_i, \qquad \|Z\|^2 \le E_w - \sum_{i\in J} \beta_i^2 (2u_i - u_i^2).$$

Substitution in (4.11) proves the assertion, including its strictness. In particular adjacent isolated sites are allowed. □

For an actual cap isolated within its component in the retained range $0 < u < 1$, retain its full foreign projection and write

$$c = Ap - \gamma\omega_i - v_{\text{for}}, \quad Q = \|v_{\text{for}}\|^2, \quad u = (1-Q)/\gamma^2 \in (0,1), \quad D_i = \|\omega_i\|^2.$$

Positive clearing gives the actual vector

$$\eta = c + v_{\text{for}} + \gamma u n_i = \gamma\{\sqrt{D_i - u}\, p - (\omega_i - u n_i)\},$$
$$\eta^2 = -\gamma^2 u(1-u).$$

It is future pointing, zero against the contacted foreign block and c, negative against n_i, and nonpositive against every other wall. Lemma 4.3 therefore gives

$$\langle x, \omega_i - u n_i\rangle \ge \sqrt{D_i - u} - \sqrt{u(1-u)}.$$

The loss from $\sqrt{D_i}$ is at most $a_i u + \sqrt{u(1-u)}$, where $a_i = (\sqrt{D_i} + \sqrt{D_i - 1})^{-1}$ in the chain ranges used. Thus a reference with $H_i + \ell < \sqrt{D_i}$, positive prices $\beta_i \ge b$, and $a_i \le a$ absorbs every such foot whenever

$$au + \sqrt{u(1-u)} - \frac{b}{2}(2u - u^2) < \ell \qquad (0 < u < 1).$$

The actual cap inventories, golden-helper exclusions and numerical reference coefficients are catalogue item A0. Their complete proofs are in Addendum A; the lemma proves the shared analytic combination used by those component estimates.

5.3 Why the golden field requires a lattice argument

Put $\varphi = (1+\sqrt{5})/2$. Bugaenko's theorem applies to the literal lattice $L = \mathbb{Z}[\varphi]^{n+1}$ with form

$$f = -\varphi t^2 + x_1^2 + \cdots + x_n^2 :$$

its full integral reflection group has an infinite-volume fundamental chamber for $n \ge 8$ [15]. Identifying the field alone does not place a chamber's reflection group in that integral group. The missing step is to show that every compatible invariant lattice gives reflections preserving L.

An actual whole-D_n long-end-five star supplies this information. Use the unit-equivalent form $f = -(\varphi-1)t^2 + \sum x_i^2$, old raw roots $-e_i + e_{i+1}$ and $-e_{n-1} - e_n$, and the cap $r = e_0 + \varphi e_1$, all of square two. Its full indefinite Gram forces the actual field to be $\mathbb{Q}(\varphi)$ by descent. The old group contains even spatial sign changes and permutations. Conjugating the cap reflection gives maps R_i with block

$$\begin{pmatrix} \varphi & -\varphi \\ 1 & -\varphi \end{pmatrix}$$

on coordinates $0, i$. The same configuration occurs in the whole-B_n head-five case. Arithmeticity supplies a full invariant $\mathcal{O}_K$-lattice Λ for the actual reflection group.

For distinct spatial indices i, j, k, even sign changes satisfy $(1-d_{ij})(1-d_{ik}) = 4E_{ii}$. Permutations and the displayed R_i then give every $4E_{ab}$ in their generated algebra; for example $4E_{ii}R_i = 4E_{i0} - 4\varphi E_{ii}$. The golden ring is a PID (its Minkowski bound is $\sqrt{5}/2 < 2$), so scaling the coordinate-content ideal to the unit ideal gives the sandwich

$$4L \subset \Lambda \subset L.$$

This is a statement about the actual invariant lattice, not just its rational span.

Here is the precise finite lattice-classification input from A0, Addendum A's "The actual golden long-end star and invariant lattices". Modulo two put

$$\ell(t, x) = \varphi t + \sum x_i, \qquad H = \ker \ell, \qquad v = (\varphi^2, 1, \ldots, 1),$$

and let E be the inverse image of H in L. The invariant subspaces of $L/2L$ are $0, \langle v \rangle, H, L/2L$, with $v \in H$ exactly when n is odd. Applying this classification to both residue flags $(\Lambda + 2L)/2L$ and $((\Lambda \cap 2L)/2)/2L$, and retaining every lift, gives the complete list

$$L, \quad E, \quad \mathcal{O}_K v + 2L, \quad \mathcal{O}_K(v + 2z) + 2E \quad (n \text{ odd}, \ z \in L).$$

The supplement proves the residue classification and the lift step; classifying only subspaces modulo two would not suffice.

The passage from this list to the literal lattice is intrinsic and illustrates the argument. We have

$$E = \{x \in L : f(x) \in 2\mathcal{O}_K\}, \qquad E^* = L + \mathcal{O}_K(v/2), \qquad f(v) = n - 1 - 2\varphi.$$

The dyadic valuation of $f(v)$ is zero for even n, one for odd n. Writing $x = y + av/2$ shows

$$\{x \in E^* : f(x) \in \mathcal{O}_K\} = L :$$

if a is not divisible by two, its final norm term has negative valuation while the other terms are integral. Thus any isometry of E, and by duality of $\mathcal{O}_K v + 2L = 2E^*$, preserves L. For a deep odd lattice, put $\lambda = v + 2z$. Its norm still has valuation one; the same calculation gives

$$\{x \in \Lambda/2 : f(x) \in \mathcal{O}_K\} = E.$$

Its isometries therefore preserve E and then L. Every lattice in the complete list has $O(f, \Lambda) \subset O(f, L)$. The actual reflection group consequently lies in the literal integral reflection group. Its full arrangement chamber refines the assumed compact chamber, contradicting the cited infinite-volume theorem. This excludes the actual whole-B/D isolated long-end-five stars.

5.4 From component bounds to actual endpoint parameters

The geometric counting starts afresh at total rank 17. As in the compact proof there is exactly one dominant component: two each cost more than $1/2$, while without one even discarding the single allowed rank-one defect leaves cost $> 16/14$. Typed defects give the stronger bound $> .09 + 15/14$. Exceptional types and all ordinary dihedrals are included in the component bounds; a field restriction is used only after it has been proved for the actual configuration.

Catalogue A0 supplies the strengthened, spectator-uniform arithmetic component estimates. In particular simultaneous absorption and the complete helper inventory give

$$e(A_{15}) > .8348, \qquad e(A_{16}) > .8703, \qquad e(A_{17}) > .9032.$$

Smaller chains, typed defects and short B/D tails have their separate actual-rank estimates. Those estimates feed the complete fourteen-group forest partition and its geometric consumers in A5. The following is the geometric interface needed for the algebraic part of the proof.

Proposition 5.2 (Rank-seventeen interface)**.** *At an original farthest vertex of a compact arithmetic Coxeter* 17*-polytope, all proper-component rows of the complete cover are impossible. The remaining whole diagrams are* A_{17}, B_{17}, D_{17}*. The* A_{17} *endpoint parameters obey Proposition 5.3 below. Each of the other two diagrams gives a pair in the quadratic domain of Proposition 5.4(iii). The two pairs stated there are geometrically impossible in each diagram.*

The proof is A5's "Complete rank-seventeen reduction" and "Every rank-seventeen row has a complete consumer". The whole-B_{17} and whole-D_{17} inclusions and consumers are distinct, even though their last algebraic domain is shared. The statement includes every finite, zero-own, parallel and dotted role in those actual-wall classifications; it is not an assertion that a bare abstract forest is realized.

Write $K_7 = \mathbb{Q}(2\cos(2\pi/7))$ and $K_9 = \mathbb{Q}(2\cos(2\pi/9))$ for the two real cubic cyclotomic fields used below.

Proposition 5.3 (Actual chain endpoints)**.** *If the whole incident diagram is* A_{17}*, both actual endpoint opposites are isolated within that component. Their integral parameters* U, V *generate the same actual field and satisfy*

$$36/17 < U, V < 3969/425, \qquad 0 < U^\sigma, V^\sigma < 36/17 \quad (\sigma \neq \mathrm{id}). \tag{5.2}$$

The field is nongolden, has degree at most five, and at least one endpoint is finite. In fact

$$K \in \{\mathbb{Q}(\sqrt{2}), \mathbb{Q}(\sqrt{3}), K_7, K_9\};$$

in the cubic fields a finite endpoint has label seven or eighteen, respectively.

The whole-A_{17} proof in Addendum A (catalogue A5) gives the complete endpoint inventory, golden-end cases, degree bound and finite-anchor reduction, before the algebraic computations are applied. It retains every finite-white and own case, including the golden time-compatible exception. One decisive step explains the simultaneous accounting. The common body has squared norm .9037838. Both dotted cores are paid at weight .3; the outer coefficients change from .1382 to .0782 and still fund every central foot. On the entire $0 < u < 1$ interval the extra contribution is

$$.6\{\sqrt{17/9 - u} - \sqrt{u(1-u)} - .4174\} - .09(8/5 - 2u + u^2) > .04969.$$

The complete circle bounds in that finite-anchor proof give $.9037838 + 2(.04969) > 1$ if both ends are dotted. No unavailable strong-five floor is used. The complete finite alphabet, including label 30 before its conjugate test, then gives the stated fields, rather than a prescribed field list.

In raw old-root normalization of square two, a cap has square $2U$ and time square $F(U)/18$, where $F(X) = X(17X - 36)$. The inverse endpoint entry is $1/18$. Every actual pair consequently satisfies

$$R = F(U)F(V) \in K^2, \qquad g = (UV - \sqrt{R})/18 \in \mathcal{O}_K, \qquad L = g^2/(UV) \in \mathcal{O}_K. \tag{5.3}$$

The root is physically positive. Acuteness gives $g \leq 0$. If $g \neq 0$, nonphysical positive definiteness gives $0 < L^\sigma < 4$. The branch $g = L = 0$ is retained. Physical owns and mutual parameters must be native or dotted; four would be a compactness-forbidden parallel pair. The endpoints need not be equal.

5.5 The four finite algebraic propositions

The computations check necessary algebraic domains. An empty domain excludes an actual configuration after geometric inclusion has been proved. A nonempty domain supplies necessary pairs, followed by a separate geometric contradiction, not examples.

5.5.1 Full quadratic integer rings

For squarefree $d > 1$ use $\omega = (1 + \sqrt{d})/2$ if $d \equiv 1 \pmod 4$, otherwise $\omega = \sqrt{d}$. Let Δ be the fundamental discriminant. If $Y = a + b\omega$ generates K, $Y > Y' > 0$, $Y < B$, then

$$0 < b\sqrt{\Delta} = Y - Y' < B.$$

Thus $\Delta < B^2$, and b and the strip-permitted integers a have explicit finite bounds. Squarefree trial division and these integer intervals cover the full maximal order.

Use rational pairs $a + b\sqrt{d}$. Integrality in the half basis means $2b, a - b \in \mathbb{Z}$; otherwise both coefficients are integral. Signs use exact rational square comparisons. For $b \neq 0$, a field square requires norm n^2 with rational $n \geq 0$ and one of

$$x^2 = (a + n)/2, \quad x^2 = (a - n)/2$$

a nonzero rational square. Set $y = b/(2x)$ and verify the full square identity. Both signed norm branches are necessary and sufficient. For $b = 0$ test rational squares and d times rational squares, including zero. A square norm alone is insufficient.

The complete positive quadratic finite native alphabet is

$$1, 2, 3, (3 + \sqrt{5})/2, (5 + \sqrt{5})/2, 2 + \sqrt{2}, 2 + \sqrt{3}.$$

Indeed $\phi(m) \leq 4$ leaves $m = 3, 4, 5, 6, 8, 10, 12$. Zero is orthogonality, values above four are dotted, and each radical value is retained only in its actual field.

Proposition 5.4 (Recorded finite outputs)**.** *In each quadratic domain below the parameters are algebraic integers generating the same actual quadratic field. At the indicated complete domains the exact outputs are as follows.*

1. *The quadratic A_{17} domain (5.2)–(5.3) has no survivor. The strips imply $\Delta < 88$. Its full enumeration and exact rejection certificates are catalogue A1.*

2. *The two cubic A_{17} finite-anchor domains are empty after the necessary tests. Their full integer rings are $\mathbb{Z}[U]$, where U is the largest root of $X^3 - 5X^2 + 6X - 1$ or $X^3 - 6X^2 + 9X - 3$, respectively. The complete coefficient boxes for $V = a + bU + cU^2$ are $[-4, 4] \times [-9, 9] \times [-3, 3]$ and $[-6, 6] \times [-9, 9] \times [-2, 2]$. The strips and cycles are those of the preceding actual endpoint reduction, with the full native alphabet at the physical embedding.*

3. *The quadratic domain*

$$4 < U < 11,\ 0 < U' < 2, \quad 2 < W < 676/17,\ 0 < W' < 4/17,$$

$K = \mathbb{Q}(U) = \mathbb{Q}(W)$,

$$g = UW - \sqrt{U(U-2)W(17W-4)}, \quad g,\ L = g^2/(2UW) \in \mathcal{O}_K, \quad 0 < L' < 4,$$

with the physically positive square root lying in K and the physical mutual parameter native or dotted, has $\Delta < 121$. Its only survivors are

K	U	W
$\mathbb{Q}(\sqrt{13})$	$(5 + \sqrt{13})/2$	$11 + 3\sqrt{13}$
$\mathbb{Q}(\sqrt{17})$	$(7 + \sqrt{17})/2$	$(21 + 5\sqrt{17})/2$.

4. The quadratic whole-B_{16} domain

$$2 < U < 12,\ 0 < U' < 2, \quad 4 < V < 64,\ 0 < V' < 1/4, \quad UV/2 \in \mathcal{O}_K,$$

$$k^2 = \frac{(U-2)V}{4U(4V-1)}, \quad k > 0, \quad G = 4Uk(4V-1) - UV, \quad L = G^2/(2UV),$$

with $k \in K$, integral G, L, actual acuteness and the following exact predicates, has $\Delta < 144$: $G \geq 0$; if $G = 0$ then $L = L' = 0$, whereas otherwise $0 < L' < 4$; and U and every nonzero L must belong to the native finite alphabet of their actual field or be greater than four. In particular $L = 4$ is excluded, and no continuous nonnative value below four is allowed. The identity

$$L + G = ((15U - 32)/2)V - 2U + 4$$

is checked for the physical positive square root k. The sole survivor is $U = 5 + 2\sqrt{6}$, $V = 10 + 4\sqrt{6}$.

These outputs are catalogue items A1–A4. The complete ring arguments, coefficient bounds, exact sign and field-square tests, termination proofs and computation records are given there. In particular both rational norm branches are retained; a square norm alone is never sufficient. The actual inclusions for whole B_{17} and D_{17} are distinct, even though their last algebraic domains coincide. Their two surviving pairs, and the one B_{16} pair, are excluded by the separate geometric consumers in catalogue A5. Proposition 5.2, the two empty A_{17} domains and the separate geometric consumers therefore exclude dimension seventeen.

5.6 The final whole-D_{16} configuration

Proposition 5.5 (Rank-sixteen interface)**.** *At an original farthest vertex of a compact arithmetic Coxeter 16-polytope, the complete nineteen-group cover and its geometric consumers leave only whole D_{16}, once the finite output Proposition 5.4(iv) is used for whole B_{16}.*

This is A5's "Complete rank-sixteen reduction" and its complete case dispatch. The proof uses a fresh total-rank-sixteen cover, including actual D_{11} and short-tail boundaries. Its conditioned A_{11} estimate is proved independently before the partition; the uniform 13/20 estimate would not imply it. Nothing is deduced by deleting a root from the rank-seventeen list.

Proposition 5.6 (Whole-D_{16} reduction)**.** *For this whole-D_{16} configuration, the actual field is quadratic. The actual long head and both genuine signed forks are isolated and dotted. There are integral generators $U, F, G \in \mathcal{O}_K$ satisfying*

$$4 < U < 98/9, \quad 0 < U' < 2, \qquad 4 < F, G < 121/2, \quad 0 < F', G' < 1/2, \tag{5.4}$$

$$UF/4,\ UG/4,\ FG/4 \in \mathcal{O}_K. \tag{5.5}$$

Their raw time squares are

$$T_H = U(U-2), \qquad T_F = 2F(2F-1), \qquad T_G = 2G(2G-1), \tag{5.6}$$

and every pair of time products is a square in K.

Its complete proof is A5's actual-rank wall, field and endpoint reduction. We state the geometric data on which its last steps depend, and explain their use below. A higher-rank head theorem or an algebraic search alone would not supply this inventory.

The original coordinates satisfy

$$x_1 > \cdots > x_{15} > |x_{16}|, \quad P_j = e_1 + \cdots + e_j, \quad S_j = x \cdot P_j, \quad f_j = \sqrt{j} - .61/\sqrt{2}.$$

The same reduction gives $S_j > f_j$ for $5 \le j \le 14$ and one positive prefix reference w with $x \cdot w > E_{14} = \|w\|^2$ and

$$E_{14} > .9084, \quad b < .361, \quad \beta_5 > .14, \quad \beta_{14} > .13, \quad f_{14} < 3.311, \qquad S_6 > 2.01815. \tag{5.7}$$

Here b is the reference's first-five mean. These coefficients are spent individually below. The head raw root has square $2U$, spatial part Ue_1; the fork raw roots have squares $2F, 2G$ and spatial parts $(F/2)(P_{15} + e_{16})$, $(G/2)(P_{15} - e_{16})$. They remain two distinct directions even if $F = G$.

Every original cycle remains mandatory even when only its necessary consequences are used below. The full field and endpoint case proofs are retained in A5.

Lemma 5.7 (Initial whole-D_{16} inventory)**.** *At the same original whole-D_{16} vertex, $S_j > f_j$ for $5 \le j \le 14$, and the positive reference has* (5.7)*. The actual long head is isolated, with finite own label at least seven or dotted own. Every actual fork has edge cosh less than twelve and is either isolated dotted or has exactly one simple-three contact at an endpoint of its retained A_{15} path, with positive own contact. Own zero and parallel fork roles are excluded. If both forks are isolated, the actual field has degree two or three.*

Proof location for the initial inventory. The initial inventory is catalogue A5's actual-rank geometric input. Its complete proof in Addendum A, "The last actual diagram," checks all original walls, closes the late spine and both signed fork inventories, and records coefficients before use. It is not a specialization of a higher-rank theorem. Starting from this inventory, A5 proves the necessary quadratic triple. The ideas behind its strengthened geometric bounds are illustrated next. □

5.6.1 Using a prefix reference without spending it twice

If $x_1 \ge .632$, replace only the first-five flat block of the reference in (5.7), preserving its sum. Its new coordinates are $.632, (f_5 - .632)/4$ four times. The next slope remains smaller, so all prefix coefficients remain nonnegative. The single-reference gain is

$$E_{14} + \frac{5}{4}(.632 - .361)^2 > 1.00020125,$$

which is impossible. Thus $x_1 < .632$. The actual head-edge comparison then gives head cosh $< 7/3$ and $U < 98/9$. This explains the narrow physical head strip; it is not an arbitrary search cutoff.

Once the head is dotted, put $u = 4/U$. The same positive actual core as in the compact proof yields

$$x \cdot Z \ge H(u), \quad Z = (1 - u/2)e_1 + (u/2)e_2, \quad Z^2 = R(u),$$

where $H(u) = (\sqrt{2 - u} - \sqrt{u(1 - u)})/\sqrt{2}$ and $R(u) = 1 - u + u^2/2$. The complete scalar estimate in A5 is $.8H(u) - .16R(u) > .295$ for $0 < u < 1$. The weight .4 on $Z - P_5/5$ spends $.08 < .14$ at P_5, with zero cross term against the flat block, and gains more than $.295 - .8(.361) + .032 = .0382$. One reference therefore exceeds .9466.

The P_{14} coefficient is still unused. If a genuine signed fork had cosh at least eleven, its pressure on $P_{15} + se_{16}$ would exceed 3.65. A weight .13 on $P_{15} + se_{16} - P_{14}$ has residual square two and zero cross with the head correction. It would give

$$\|x\|^2 > .9466 + .26(3.65 - 3.311) - .0338 = 1.00094.$$

Hence both fork coshes are below eleven, giving the strict fork bound $F, G < 121/2$. A5 uses these bounds to exclude cubic fields and every nonisolated endpoint role, with full field-square tests and path cycles. This is why the reduction produces the actual quadratic, isolated, dotted triple of Proposition 5.6.

The old connected simple-three system has one raw norm squareclass. After its permitted square-two normalization, the isolated head and fork raw roots have precisely the spatial parts displayed above. Their physically indefinite principal Grams are defined over $\mathbb{Q}(U)$ and $\mathbb{Q}(F)$, so actual-field descent proves the asserted field equalities. Their original cycles give integral g, L with

$$L_{HF} + g_{HF} = (15UF - 32F - 8U + 16)/4, \qquad L_{FG} + 7g_{FG} = 15FG/4 - 8(F + G) + 4,$$

and the analogous head/G identity. These yield all three quarter-products. The common actual K-time line yields the full field-square conditions. The geometric meaning of these variables is retained through the arithmetic argument.

5.6.2 Norm bounds, integer orders and valuations

Put

$$N_H = \mathrm{N}\, U, \quad T = \mathrm{Tr}\, U, \quad A = -\mathrm{N}(U - 2) = 2T - 4 - N_H.$$

The strict strips imply

$$5 \le T \le 12, \quad 1 \le N_H \le 21, \quad 1 \le A \le 17.$$

For a fork put $n = \mathrm{N}\, F$, $S = \mathrm{Tr}\, F$ and $B = 2S - 4n - 1 > 0$ odd. Then

$$1 \le n \le 30, \quad 5 \le S \le 60, \qquad \mathrm{N}\, T_H = -N_H A, \quad \mathrm{N}\, T_F = -4nB. \tag{5.8}$$

A nonsquare rational norm disproves a field square; a square norm still requires the full square test.

Use the full integer order of $K = \mathbb{Q}(\sqrt{d})$. The gap between the embeddings of U is a positive integral multiple of $\sqrt{\Delta}$, so

$$\Delta < (98/9)^2 < 119, \qquad \Delta \le 118. \tag{5.9}$$

For a prime $\mathfrak{p}$ over two, normalize $v_{\mathfrak{p}}(\mathfrak{p}) = 1$ and write

$$e = v_{\mathfrak{p}}(2),\ r = v_{\mathfrak{p}}(U),\ s = v_{\mathfrak{p}}(U - 2),\ k = v_{\mathfrak{p}}(F),\ l = v_{\mathfrak{p}}(G).$$

The factors $2F - 1, 2G - 1$ are units there. Quarter integrality and the time squares give

$$r + k, r + l, k + l \ge 2e, \qquad r + s + e + k,\ r + s + e + l \equiv 0 \pmod 2. \tag{5.10}$$

Also $s = r$ for $r < e$, $s = e$ for $r > e$, and $s \ge e$ for $r = e$. Norm valuations are $\sum f_{\mathfrak{p}} v_{\mathfrak{p}}$. The quadratic possibilities are inert $(e, f) = (1, 2)$, ramified $(2, 1)$, or two split primes $(1, 1)$. No PID or unramifiedness assumption enters.

5.7 A worked dyadic branch and the final exclusion

The possible behavior of two is inert, ramified, or split. In each case (5.10) forces divisibility of the fork parameters; their strict norm and conjugate bounds then leave short full-integer-ring lists. A square rational norm is only a necessary filter. Surviving square-norm products still require a square in the actual field, and surviving parameters still have to satisfy the original geometry.

We work out the odd-head-norm branch completely. One elementary strip calculation is useful. If $F/4$ is integral, then $16 \mid \mathrm{N}\, F < 30.25$, so $\mathrm{N}\, F = 16$. For $W = F/4$,

$$\mathrm{N}\, W = 1, \quad 8 < W < 121/8, \quad \mathrm{Tr}\, W = W + 1/W \in \{9, \dots, 15\}.$$

Its quadratic polynomial $X^2 - (\mathrm{Tr}\, W)X + 1$ and $\Delta \le 118$ leave the four possibilities

K	F
$\mathbb{Q}(\sqrt{77})$	$18 + 2\sqrt{77}$
$\mathbb{Q}(\sqrt{6})$	$20 + 8\sqrt{6}$
$\mathbb{Q}(\sqrt{13})$	$22 + 6\sqrt{13}$
$\mathbb{Q}(\sqrt{3})$	$28 + 16\sqrt{3}$.

Traces $12, 13, 15$ give discriminants $140, 165, 221$ and are excluded. This uses the full integer order, including half-integral coordinates.

5.7.1 Odd norm

If N_H is odd, at every dyadic prime $r = s = 0$. In an inert or split field, (5.10) makes both fork valuations odd and at least three; their norms are divisible by 64, contrary to $n \le 30$. Thus two ramifies. Now k, l are even and at least four. Both fork norms are sixteen and $F/4, G/4$ integral. The unit-fork table leaves only $\mathbb{Q}(\sqrt{3})$ or $\mathbb{Q}(\sqrt{6})$. The full head strips with odd norm give

K	U	$\mathrm{N}\, T_H$	$\mathrm{N}\, T_F$
$\mathbb{Q}(\sqrt{3})$	$5 + 2\sqrt{3}$	-39	-3008
$\mathbb{Q}(\sqrt{6})$	$3 + \sqrt{6},\ 5 + 2\sqrt{6}$	$-15,\ -15$	-960.

The radical coefficient is one or two; coefficient three exceeds the physical bound. The conjugate strip and norm parity remove the other rational coefficients. In $\mathbb{Q}(\sqrt{6})$, $6 + 2\sqrt{6} > 98/9$ and $4 + \sqrt{6}$ has even norm.

The $\sqrt{3}$ product norm is $64 \cdot 1833$, a nonsquare. For $U = 3 + \sqrt{6}$ the exact time product is $58200 + 23760\sqrt{6}$, of norm 120^2. Its possible rational square parts are 29160 and 29040, strictly between 170^2 and 171^2, so it is not a field square. The remaining triple is $U = 5 + 2\sqrt{6}$, $F = G = 4U$. The actual head and two genuine fork edges have coshes

$$C_H = (2 + \sqrt{6})/2 > 11/5, \qquad C_+ = C_- = 2(2 + \sqrt{6}) > 44/5,$$

using $\sqrt{6} > 12/5$. Equation (5.1) gives $x_1 \ge \sqrt{3/8} > .61$ and, for both $s = \pm 1$,

$$x \cdot (P_{15} + s e_{16}) > 4\sqrt{39}/7 > 3.56.$$

The last comparison follows from $6.23^2 < 39$. Averaging the two signed inequalities gives $S_{15} > 3.56$; Equation (5.7) gives $S_6 > 2.01815$. One positive reference gives

$$\|x\|^2 > .61^2 + 1.40815^2/5 + 1.54185^2/9 = 1.032821887 > 1.$$

The last pressure averages the two genuine fork edges, not a virtual root. This excludes odd norm.

5.7.2 The remaining dyadic cases

Catalogue A6 contains the complete even-norm proof, with its finite strip lemmas and all original geometric conditions. Its three branches are:

(i) Inert two: $v_{\mathfrak{p}}(U) = 1$ or 2, since higher valuation forces $\mathrm{N}\,U \geq 64 > 21$. At valuation one the exact fork strips and full field-square tests exclude every candidate. At valuation two the head is $U = 4\varphi^2$; a fork would have to be divisible by four, contradicting its complete norm-one strip in $\mathbb{Q}(\varphi)$.

(ii) Ramified two: the possible head valuations are $1, 2, 3, 4$. The first two are excluded by the full strip/square tests. At valuation three the head is $4 + 2\sqrt{2}$; fork norm and trace congruences exclude its remaining possibilities. Valuation four would give the golden head, whose dyadic prime is inert.

(iii) Split two: the discriminant bound leaves the complete squarefree list $17, 33, 41, 57, 65, 73, 89, 97, 105, 113$. The head strip is checked in $(a + b\sqrt{d})/2$, with full parity. Both dyadic primes are retained in the valuation argument. The odd-trace cases are separated by $v_2(\mathrm{N}\,U)$, and the sole even-trace head has a separate two-prime proof. The final necessary pairs are removed by full field squares or by one funded original-vector contradiction.

The complete field-case and coefficient tables are in A6, with the actual strip bounds and excluded endpoints. This proposition is a manual finite arithmetic proof using integer traces, norms, valuations and exact squares; it is not a new whole-D_{16} census assumption.

One split-case survivor shows why geometry must remain in the argument. The necessary pair is

$$K = \mathbb{Q}(\sqrt{57}), \quad U = (9 + \sqrt{57})/2, \quad F = 23 + 3\sqrt{57}.$$

Its head has $U > 8$, hence $x_1 > 1/\sqrt{3} > .577$. Rebuilding the first-five reference block with its sum fixed gives

$$E_{14} + \frac{5}{4}(.577 - .361)^2 > .96672.$$

A fork of cosh at least nine would have actual signed pressure > 3.577. The separate .13 coefficient at P_{14}, with residual square two and zero cross with the first-block change, would give

$$\|x\|^2 > .96672 + .26(3.577 - 3.311) - .0338 = 1.00208 > 1.$$

Thus its cosh is below nine and $F < 81/2$, contradicting $23 + 3\sqrt{57} > 44$. This illustrates the final geometric consumer of a necessary arithmetic survivor, without treating it as a polytope.

Proposition 5.8 (Dyadic exclusion)**.** *The actual whole-D_{16} configuration of Proposition 5.6, with its genuine signed walls and original-vector reference data, is impossible.*

Proof. Odd head norm was excluded above. For even head norm, the complete inert, ramified and split cases just specified are proved in A6, Addendum A, "Odd head norm and inert two", "Even head norm with ramified two" and "The complete split branch". These exhaust the quadratic fields. Every contradiction uses an original cycle, a full field-square condition or one funded norm estimate on the same original vector. □

The compact theorem first gives $n \leq 17$. Proposition 5.2 and A1–A3 exclude seventeen. Proposition 5.5, using A4, leaves only whole D_{16} in rank sixteen. Its actual-field and wall reduction gives Proposition 5.6, and the exhaustive dyadic argument excludes it. Therefore $n \leq 15$, proving Theorem 1.1(iii).

6 Proof catalogue and reproduction

The complete mathematical supplement is the repository edition fixed in [8]. Its Parts I and IV contain the finite-volume proof, Parts II and V the compact proof, and Parts III and VI the arithmetic proof. The technical propositions invoked here retain their full hypotheses and complete proofs there. In particular, the online geometric case analyses are as much a part of the argument as the finite computations.

The accompanying proof catalogue assigns FV1–FV4 to the finite-volume propositions, C1–C2 to the compact case results, and A0–A6 to the arithmetic component, finite-domain and terminal results. It gives exact source labels in the fixed supplement, so the reader can pass from a proposition in this article to its detailed proof without searching the entire collection. The catalogue also explains which central arguments are printed here and which technical proofs are supplied online.

A current companion for all three bounds. The entry point `repro/dimension_bounds/` contains a reader's guide and a machine-readable manifest. For each catalogue item it connects the statement, complete proof, exact checker, input records and recorded execution; where applicable it also identifies the Lean source, premise map, compilation and separate axiom audit. The manifest binds retained inputs by hashes and records the pinned Python and Lean environments. Its metadata check inspects those bindings without rerunning the mathematics. The historical manuscript catalogue keeps its original edition selections; in particular, its older compact identifiers are not presented as checks of the present bounds 17 and 15.

What can be checked. The companion separates reading the written proof, checking a retained certificate, regenerating a finite computation and checking a conditional formal deduction. The finite-volume current-rule conjunction has recorded execution for 259 scoped units, including all 64 selected application shards. This does not constitute a fresh combined replay of its older 146-job inventory or regeneration of every inherited classification. The public companion identifies the small-input checks available from a Git checkout and the operations whose larger inputs are not distributed. No asset from the separately indexed bulk collection is included in the arXiv source archive.

The Lean sources check selected conditional deductions. For compact 17, the scalar and composition modules have separate compilation and actual axiom-audit records for ten declarations; the dyadic kernel used for arithmetic 15 has corresponding records for fourteen declarations. Their ordinary hypotheses include the indicated geometric, analytic, field and finite-data propositions. Those hypotheses must be read in the source and premise maps: a printed list of foundational axioms does not list theorem parameters. The full written proofs supply the mathematics outside these kernels. This editorial revision preserves the existing execution records and introduces no new mathematical or Lean execution.

Acknowledgments. The author used OpenAI's GPT-6-astra and GPT-6.1-sol as research and coding assistants to investigate proof ideas, help strengthen the results, improve the exposition, and generate and audit the Python and Lean 4 code accompanying this work. The author reviewed the resulting mathematics and code and takes responsibility for the final content.

References

[1] M. N. Prokhorov. The absence of discrete reflection groups with noncompact fundamental polyhedron of finite volume in Lobachevskii space of large dimension. *Mathematics of the USSR-Izvestiya*, 28:401–411, 1987. https://www.mathnet.ru/eng/im1484.

[2] A. G. Khovanskii. Hyperplane sections of polyhedra, toroidal manifolds, and discrete groups in Lobachevskii space. *Functional Analysis and Its Applications*, 20:41–50, 1986. https://www.math.utoronto.ca/askold/1986-Faa-1-english.pdf.

[3] E. B. Vinberg. The absence of crystallographic groups of reflections in Lobachevsky spaces of large dimension. *Trudy Moskov. Mat. Obshch.*, 47:68–102, 1984. English translation, Trans. Moscow Math. Soc. (1985), 75–112. https://www.mathnet.ru/eng/mmo444.

[4] E. B. Vinberg. Hyperbolic reflection groups. *Russian Math. Surveys*, 40(1):31–75, 1985. doi:10.1070/RM1985v040n01ABEH003527.

[5] R. E. Borcherds. Automorphism groups of Lorentzian lattices. *Journal of Algebra*, 111:133–153, 1987. https://math.berkeley.edu/~reb/papers/al/al.pdf.

[6] Lizi Guo and Fangting Zheng. Dimension bounds for finite-volume hyperbolic Coxeter polytopes. arXiv:2611.01795v1, 2026. https://arxiv.org/abs/2611.01795.

[7] Alexander Kolpakov. Corner payments and a dimension bound of 57 for finite-volume hyperbolic Coxeter polytopes. Public repository manuscript, 21 October 2026, 2026. Initial public snapshot 32b68355. https://github.com/sashakolpakov/CoxeterPolytopes/blob/32b68355cdb13ca8629f3fa8b98d424dd70472c7/manuscripts/finite_volume57.tex.

[8] Alexander Kolpakov. Dimension bounds for hyperbolic Coxeter polytopes: complete proofs and addenda. Repository edition of 30 October 2026, snapshot 36eeaddf, 2026. Full manuscript, Parts I–VI. https://github.com/sashakolpakov/CoxeterPolytopes/blob/36eeaddf646eff25d4990c9ab730a20ada21f80f/arxiv/hyperbolic_coxeter_dimension_bounds/main.tex. The matching PDF and reproduction guides are linked from the repository README.

[9] A. Felikson and P. Tumarkin. On simple ideal hyperbolic Coxeter polytopes. arXiv:math/0502413. https://arxiv.org/pdf/math/0502413.

[10] M. Chein. Recherche des graphes des matrices de Coxeter hyperboliques d'ordre ≤ 10, 1969. https://www.numdam.org/item/M2AN_1969__3_3_3_0.pdf.

[11] V. V. Nikulin. Discrete reflection groups in Lobachevsky spaces and algebraic surfaces. In *Proceedings of the International Congress of Mathematicians*, volume 1, pages 654–671, Berkeley, 1986. https://www.mathunion.org/fileadmin/ICM/Proceedings/ICM1986.1/ICM1986.1.ocr.pdf.

[12] V. V. Nikulin. On ground fields of arithmetic hyperbolic reflection groups. In *Groups and Symmetries*, volume 47 of *CRM Proceedings and Lecture Notes*, pages 299–326, 2009. Corrected version arXiv:0708.3991v2. https://arxiv.org/pdf/0708.3991v2.

[13] I. Agol, M. Belolipetsky, P. Storm, and K. Whyte. Finiteness of arithmetic hyperbolic reflection groups. *Groups, Geometry, and Dynamics*, 2:481–498, 2008. https://doi.org/10.4171/GGD/47.

[14] M. Belolipetsky. Arithmetic hyperbolic reflection groups. https://arxiv.org/abs/1506.03111.

[15] V. O. Bugaenko. Groups of automorphisms of unimodular hyperbolic quadratic forms over $\mathbb{Z}[(\sqrt{5}+1)/2]$. *Vestnik Moskov. Univ. Ser. 1 Mat. Mekh.*, pages 6–12, 1984. No. 5. https://www.mathnet.ru/eng/vmumm3408.