Dimension bounds for hyperbolic Coxeter polytopes

Alexander Kolpakov

4 November 2026

Abstract

We prove that a finite-volume hyperbolic Coxeter polytope has dimension at most 57, that a compact one has dimension at most 17, and that a compact arithmetic one has dimension at most 15. The finite-volume argument compares corner weights with the geometry of vertex links. The compact arguments use projections of one vector associated with an original farthest vertex; arithmeticity further restricts the ground field and integral wall parameters. This concise edition presents the main geometric arguments and their exact local interfaces. The complete case proofs, computational proofs and reproduction instructions are supplied as versioned addenda in the accompanying repository.

1 Introduction

All polytopes in this article are convex, finite-sided and full-dimensional. A hyperbolic Coxeter polytope has dihedral angles that are integer submultiples of π; reflection in its facets generates a discrete group with the polytope as fundamental domain. Hyperbolic space has curvature -1.

Theorem 1.1. *Let $P \subset \mathbb{H}^n$ be a finite-volume hyperbolic Coxeter polytope. Then:*

(i) $n \leq 57$*;*
(ii) *if P is compact, then $n \leq 17$;*
(iii) *if P is compact and its reflection group is arithmetic, then $n \leq 15$.*

The classical finite-volume ceiling is 995, due to Prokhorov [1], using the combinatorial estimates of Khovanskii [2]. Vinberg obtained the compact ceiling 29 [3, 4]. The largest known finite-volume examples have dimension 21 [5]; the bounds here are upper bounds and make no new existence assertion. Compactness is essential in Theorem 1.1(iii).

Guo and Zheng recently obtained the general finite-volume bound 81 [6]. An earlier version of our bound 57, including its corner-weight method and higher-dimensional argument, was publicly posted on 21 October 2026 [7]. The present article condenses the three proof families assembled in the repository edition of 30 October [8].

The methods separate after the common local geometry. In the finite-volume case, triangles and quadrilaterals require corner payments that high-dimensional vertex diagrams cannot accommodate. Nonsimple ideal vertices are treated through the product structure of their links. In the compact case, local walls force quadratic lower bounds for components of a single vector of norm less than one. Arithmeticity enters only in the third argument, through integral cycles and positivity at the other real embeddings.

Detailed proofs and version. The full proof addenda [8] are fixed at repository commit `36eeaddf646eff25d4990c9ab730a20ada21f80f`. Parts I and IV contain the finite-volume argument,

Parts II and V the compact argument, and Parts III and VI the arithmetic argument. We refer to these as Addenda FV, C and A, respectively. The local propositions below cite the complete case proofs at that version; Section 6 gives the proof-catalogue entry points and reproduction package. Thus the long case analyses remain part of the proof, with stable references independent of subsequent research.

2 Actual walls and vertex diagrams

We use a Lorentz space of signature $(-,+,\dots,+)$ and the future unit hyperboloid model of $\mathbb{H}^n$. Outward wall normals are spacelike; the chamber inequalities are $\langle y, e_i\rangle \le 0$. Distinct wall normals have nonpositive products. Finite vertex diagrams are spherical. A horospherical link at an ideal vertex is a product

$$L_v = \prod_i \Delta^{r_i}, \qquad \sum_i r_i = n-1, \tag{2.1}$$

whose factors correspond to irreducible affine diagrams of orders $r_i + 1$. These standard face and link facts are used in their actual-wall form [4, 9].

Two elementary observations account for much of the local geometry.

Lemma 2.1. *A positive-definite Gram matrix with nonpositive off-diagonal entries has entrywise nonnegative inverse. A positive-definite principal set of actual chamber walls has a feasible finite common point.*

Proof. If B is the matrix, choose s larger than its largest eigenvalue. Then $M = sI - B$ is entrywise nonnegative and has spectral radius less than s, so $B^{-1} = s^{-1}\sum_{j\ge 0}(M/s)^j$.

Take a strict interior future timelike point o and put $b_i = -\langle o, e_i\rangle > 0$ for the selected walls. The vector

$$q = o + \sum_i (B^{-1}b)_i e_i$$

is orthogonal to the selected normals and has square $q^2 = o^2 - b^T B^{-1} b < 0$. Its product with o is negative, so it is future pointing. Every other chamber inequality is strict, because the coefficients are nonnegative and all other wall products are nonpositive. Normalize q to the hyperboloid. □

An irreducible affine principal set has a positive kernel vector (u_i). The root combination $\nu = \sum_i u_i e_i$ is null, nonzero since $\langle o, \nu\rangle < 0$, and feasible against every wall. It represents an actual ideal point. Irreducibility and the positive definiteness of proper deletions identify its whole affine factor. In contrast, a kernel vector in an indefinite core represents the zero vector: that vector is null and orthogonal to a timelike vector in the core span, whose perpendicular space is positive definite. Both interpretations are needed when controlling extensions.

At an actual two-face, the supporting walls form a positive-definite diagram of rank $n-2$. Adding one side gives an edge diagram, and adding two adjacent sides gives a positive-semidefinite corner diagram. For opposite sides of a quadrilateral, the corresponding principal diagram is indefinite. A positive or affine alternative would give a feasible common finite or ideal point of those sides. Minimal indefinite subdiagrams, inverse Gram matrices and the classification of finite and affine diagrams consequently reduce the local payment questions to explicit marked configurations.

3 The finite-volume bound

For a two-face F, let $s(F)$ be its number of sides and set

$$\mathcal{D}_5(P) = \sum_{\dim F=2} (5 - s(F)).$$

A corner rule assigns a nonnegative weight $w(v, F)$ to every incidence of a vertex with a two-face. Write $C(v) = \sum_{F \ni v} w(v, F)$.

Lemma 3.1 (Payment comparison)**.** *If every triangle receives total weight at least* 2 *and every quadrilateral at least* 1*, then* $\mathcal{D}_5(P) \le \sum_v C(v)$*.*

Proof. Each face receives at least $5 - s(F)$: this is the hypothesis for three and four sides, and follows from nonnegativity for at least five sides. Sum over all faces and interchange the two sums. □

3.1 The local rule and its finite maximum

The explicit rule is defined on whole marked vertex diagrams in Addendum FV, under "The intrinsic corner weights." It combines same-component pair weights, cross-component roles and contextual terms, with a prescribed priority among overlapping roles. Finite and affine components have separate tables. Whole-component conditions and all compensating payments are included in this one rule.

Proposition 3.2 (Local payment and finite charge)**.** *In ambient dimension* 58*, this simultaneous rule pays at least* 2 *on every triangular two-face and at least* 1 *on every quadrilateral. At every finite vertex its charge is at most* 377*.*

Construction and proof location. The geometric construction uses minimal indefinite diagrams for opposite pairs and for the three sides of a triangle. The bounded core algorithms include all permitted overlaps, cross bonds and additional supporting walls. Singular cores are controlled by their actual root relations, while nonsingular cores give inverse and Schur-complement bounds on the entire attached forest. The complete construction and extension proofs are Addendum FV's "Geometric coverage and exact local certificates" and "The dimension-specific final local rule."

The finite arithmetic maximizes an explicit function over all spherical forests of rank 58. If m_T is the multiplicity of type T, its charge, scaled by 64, has the form

$$\sum_T b(T) m_T + \sum_T j(T, T) \binom{m_T}{2} + \sum_{T<U} j(T, U) m_T m_U + \Psi(m). \tag{3.1}$$

The alphabet contains A_r, B_r, D_r at their allowed ranks through 58, the exceptional finite types, and a symbolic rank-two type for the remaining unbounded dihedral labels. Their common cost is established before that symbolic encoding is used. The tables and the complete context function are printed in Addendum FV. Catalogue item FV2 states the exact formal maximum 377 over this complete domain, including all contextual corrections. Its proof and finite certificates are in the repository. No realization of a maximizing forest is required. □

The geometric coverage theorem identifies every actual vertex with a member of this necessary finite domain. The catalogue supplies the whole-component tables, context states, complete maximization and its verification records.

3.2 The shape term

Suppose now that $n = 58$. For any vertex figure define

$$H(t) = \sum_{j=0}^{57} h_j t^j, \quad V = H(1), \quad G = \sum_{j=0}^{57} h_j \min(j, 57-j), \quad \kappa = \frac{5G - 57V}{2}. \tag{3.2}$$

For an ideal vertex, $H(t) = \prod_i (1 + t + \cdots + t^{r_i})$ by (2.1). For a finite vertex, $H(t) = 1 + t + \cdots + t^{57}$.

Lemma 3.3 (Shape curvature)**.** *For a proposed* 58*-dimensional polytope,*

$$\mathcal{D}_5(P) \geq 377 v_f + \sum_{v \text{ ideal}} \kappa(v) + 4205, \tag{3.3}$$

where v_f denotes the number of finite vertices.

Proof. Choose a generic linear function on the bounded Klein closure. Let E_v count the edges of the simple vertex figure, and let E_v^0 count those not cut by the induced level section. The incidence identities of Khovanskii [2, Theorem 9] give

$$\mathcal{D}_5(P) = \sum_v (5E_v^0/2 - E_v).$$

The index distribution of an oriented simple polytope is its h-vector. Assign each uncut edge below the section to its upper endpoint, and each above it to its lower endpoint. An endpoint of index j receives at least $\min(j, 57-j)$, so $E_v^0 \geq G_v$. Since each figure is simple of dimension 57, $E_v = 57V_v/2$.

At the global minimum and maximum the whole figure is uncut. Symmetry and $h_j \geq 1$ give, at each of these two vertices,

$$E_v - G_v = \sum_{j=29}^{57} h_j (2j - 57) \geq 841.$$

The resulting correction is $2(5/2)841 = 4205$. At a finite vertex, $G = 812$ and $V = 58$, hence $\kappa = 377$. Substitution gives (3.3). □

3.3 Ideal vertices and concentration

A marked pair in affine factor i occurs in $\prod_{j \neq i}(r_j + 1) = V/(r_i + 1)$ actual corner fibers. The all-family bounds in Addendum FV consequently give

$$C(v) \leq V \sum_i c(r_i), \tag{3.4}$$

where

$$c(r) = r/2 \ (1 \leq r \leq 8), \qquad c(r) = \frac{5r + 56}{r + 1} \ (24 \leq r \leq 57),$$

and the intervening values are

r	9	10	11	12	13	14	15	16
$c(r)$	71/16	53/11	31/6	11/2	40/7	367/60	411/64	227/34
r	17	18	19	20	21	22	23	
$c(r)$	119/18	128/19	137/20	146/21	155/22	163/23	85/12	

Proposition 3.4 (Positive ideal surplus)**.** *Every ideal vertex in dimension* 58 *satisfies* $\kappa(v) - C(v) \geq 36$.

Proof. For independent uniform variables U_r on $\{0, \dots, r\}$, put $S = \sum_i U_{r_i}$. Then

$$\frac{G}{V} = \frac{57 - \mathbb{E}|2S - 57|}{2}.$$

For $1 \leq a \leq b$, $U_{a-1} + U_{b+1}$ dominates $U_a + U_b$ in convex order. This follows by comparing the stop-loss functions at integer thresholds; the full elementary calculation is the "Concentrating two simplex ranks" lemma in Addendum FV. Conditioning proves preservation under adding a common independent summand.

Apply this transfer while two positive ranks are below eight. The sum of c-costs is unchanged, the normalized curvature cannot increase, and the product of factor orders decreases. The integer potential $\sum_i r_i^2$ strictly increases and is bounded by 57^2, so the process terminates. Its final domain consists of partitions of 57 with all parts at least eight, except possibly one part in $\{1, \dots, 7\}$.

Generate sorted partitions recursively by their first part and multiply $1 + t + \cdots + t^r$ by exact convolution. This finite calculation has 2552 terminal profiles and gives

$$\min\left(\frac{\kappa}{V} - \sum_i c(r_i)\right) = \frac{87}{10240} > 0, \qquad \min\left(\kappa - V\sum_i c(r_i)\right) = 36. \tag{3.5}$$

The complete domain, coefficients and exact comparisons are supplied with the "Complete terminal calculation" in Addendum FV. The recursion terminates because the remaining sum decreases; the unique first part proves that each partition is generated once.

Write the original normalized margin as m and the terminal one as m', with products V, V'. The transfer gives $m \geq m' > 0$ and $V \geq V'$. Thus $Vm \geq V'm' \geq 36$. Equation (3.4) proves the assertion for actual charges. □

3.4 The contradiction and the higher-dimensional tail

Proposition 3.2 and Lemma 3.1 give

$$\mathcal{D}_5(P) \leq 377v_f + \sum_{v \text{ ideal}} C(v).$$

Subtracting (3.3) and using Proposition 3.4 gives

$$0 \geq 4205 + \sum_{v \text{ ideal}} (\kappa(v) - C(v)) \geq 4205 + 36v_i > 0. \tag{3.6}$$

Hence dimension 58 is impossible, including the case $v_i = 0$.

Proposition 3.5 (Independent higher-dimensional exclusion)**.** *There is no finite-volume hyperbolic Coxeter polytope in any dimension* $n \geq 59$.

Proof and addendum reference. The range $59 \leq n \leq 163$ is treated by the dimension-specific payment families in Addendum FV's "The bounded higher-dimensional tail." Each family has its own finite maximum, affine envelopes and complete terminal-profile comparison. The ranges are 59–65, 66–97 and 98–163; all local replacement conditions are included before the charge comparison is applied.

For $n \geq 164$, the "Unbounded tail: actual framing walls and two budgets" uses Nikulin's narrow-facet theorem [10]. The proof constructs actual framing walls and proves a Lorentz inequality with

the two original endpoint budgets. Its point potentials, full-budget classical estimate and ordered packing control the complete unbounded range. The alternative in which the chosen narrow facet is compact has its own dimension estimate. These arguments concern the original chamber walls and do not require that an arbitrary facet be a Coxeter polytope. The full calculations, including their uniform rank reductions, are printed in the cited addendum. □

Combining Proposition 3.5 with (3.6) proves Theorem 1.1(i).

4 The compact bound

4.1 One original vector and legal reference vectors

Normalize wall normals to square one. The positive projections of Lemma 2.1 check every original wall. For compact P, fix z strict interior and a vertex p farthest from z. Retain this original choice throughout. Write

$$z = E(p - x), \quad E = -\langle z, p\rangle > 0, \quad x \in p^{\perp}, \quad \|x\|^2 = 1 - E^{-2} < 1. \tag{4.1}$$

The incident diagram has exactly n nodes. Its whole irreducible components T give orthogonal projections with

$$\sum_T e_T = \|x\|^2 < 1, \qquad e_T = \|x_T\|^2. \tag{4.2}$$

Every term uses this same original vector. On a geodesic the positive function $-\langle z, r\rangle$ has positive second derivative, so its maximum on a compact polytope occurs at a vertex. Consequently

$$-\langle z, r\rangle \le E \quad (r \in P). \tag{4.3}$$

If q is another vertex and $z = E_q(q - y)$, comparisons there use $\kappa = E/E_q$; q is not assumed farthest.

Lemma 4.1. *If $\xi = Ap - U$ is future timelike, $U \in p^{\perp}$, $\xi^2 = -k < 0$, and $\xi/\sqrt{k} \in P$, then for every Euclidean subspace $S \subset p^{\perp}$ containing U,*

$$\langle x, U\rangle \ge A - \sqrt{k}, \qquad \|x_S\|^2 \ge \frac{A - \sqrt{k}}{A + \sqrt{k}}. \tag{4.4}$$

At q the first bound is $\langle y, U\rangle \ge A - \kappa\sqrt{k}$.

Proof. Here x_S is the orthogonal projection of x to S. Substitute $\xi/\sqrt{k}$ into (4.3) and use $\|U\|^2 = A^2 - k$ and Cauchy's inequality. The zero-displacement case follows directly. □

Let B be the incident Gram matrix, $C = B^{-1}$, and $\omega_i = \sum_j C_{ij} n_j$, $d_i = C_{ii}$, $u_i = \omega_i/\sqrt{d_i}$. Every actual wall not through p has the form

$$c = Ap - v, \quad v = \sum_i \beta_i \omega_i, \quad \beta_i = -\langle c, n_i\rangle \ge 0, \quad A^2 = \beta^T C \beta - 1, \quad \langle x, v\rangle < A. \tag{4.5}$$

For an original edge of length L_i in direction $-u_i$, comparison at its endpoint gives

$$\langle x, u_i\rangle \ge \tanh(L_i/2). \tag{4.6}$$

Proposition 4.2 (Reciprocal shadows)**.** *Put* $T_i = (C\beta)_i/(A\sqrt{d_i})$*. If* c *closes edge* i*, then* $\tanh L_i = 1/T_i$*; otherwise* $\tanh L_i \geq T_i$*. Thus* $T_i > 1$ *forces closure and* $T_i = 1$ *is impossible. Distinct opposites of* i, j *satisfy*

$$\tanh L_i \tanh L_j \geq K_{ij}, \qquad K_{ij} = C_{ij}/\sqrt{d_i d_j}. \tag{4.7}$$

Opposites of orthogonal original directions are distinct.

Proof. For the actual opposite $d = Dp - \sum \gamma_j \omega_j$ of i, the endpoint equation gives $\tanh L_i = D\sqrt{d_i}/(C\gamma)_i$. If $c \neq d$, acuteness and the nonnegative Schur inverse give

$$AD \geq \beta^T C \gamma \geq (C\beta)_i (C\gamma)_i / d_i.$$

Here $C - C_{\cdot i} C_{i\cdot}/d_i$ is zero on row and column i and elsewhere is the inverse of the retained Gram matrix. This proves the alternative. For the opposite of i, the same Schur identity gives $T_j \geq K_{ij}/\tanh L_i$; distinctness at j gives the pair inequality. If one wall closed orthogonal i, j, clearing the other old walls by positive coefficients would give $c_F = Ap - Au_i/r_i - Au_j/r_j$, $r_i = \tanh L_i$. Positive clearing keeps $\langle z, c_F \rangle < 0$, so strict interiority requires $\langle x, u_i \rangle/r_i + \langle x, u_j \rangle/r_j < 1$. By (4.6) the sum is at least $1 + (\tau_i^2 + \tau_j^2)/2 > 1$, where $\tau_i = \tanh(L_i/2)$. □

An opposite wall contacts each retained foreign component at most once, since its finite replacement has no cycle. A positive foreign cost is $q = \beta_j^2 C_{jj} \geq 1/4$; the only cost below $1/3$ is a simple 3-contact on A_1, of cost $1/4$. Moreover

$$q < A^2 : \tag{4.8}$$

otherwise the reciprocal test closes that orthogonal direction too (or gives a null endpoint at equality). For unit directions with correlation k in a component of energy at most e, put $F(t) = 2t/(1+t^2)$. Cauchy and concavity of $\log F$ give

$$\tau_i + \tau_j \leq \sqrt{2e(1+k)}, \qquad F(\tau_i)F(\tau_j) \leq \frac{4s}{(1+s)^2}, \quad s = e(1+k)/2. \tag{4.9}$$

If this upper bound is below K_{ij}, the opposites coincide.

Actual points and simultaneous references. For an actual cap $c = Ap - v$, clear an old positive block J by $w = \sum_{i,j \in J} (B_J^{-1})_{ij} \beta_j n_i$. Set $\eta = c + w$, $\delta = \eta^2 = 1 - \beta^T B_J^{-1} \beta$. If $\delta < 0$, normalize η; if $\delta > 0$, use $p + (A/\delta)\eta$, of square $-1 - A^2/\delta$. Once selected products are nonpositive, all unselected products follow from the positive normal coefficients. The respective pressures are

$$\langle x, v - w \rangle \geq A - \sqrt{-\delta}, \qquad \langle x, v - w \rangle \geq \frac{A}{1 + \sqrt{\delta/(A^2 + \delta)}}. \tag{4.10}$$

A nonzero feasible future null vector contradicts compactness.

For any Euclidean reference W,

$$\|x\|^2 \geq 2\langle x, W \rangle - \|W\|^2. \tag{4.11}$$

Suppose $w = \sum b_i P_i$, $b_i \geq 0$, and actual pressures give $\langle x, P_i \rangle > a_i$. A correction $Q = Z - \sum t_i P_i$ with $\langle x, Z \rangle \geq h$ is legal at weight $\mu \geq 0$ only if every $b_i - \mu t_i \geq 0$. Then

$$\langle x, (w + \mu Q) \rangle \geq \sum b_i a_i + \mu(h - \sum t_i a_i), \qquad \|w + \mu Q\|^2 = \|w\|^2 + 2\mu\langle w, Q \rangle + \mu^2 \|Q\|^2. \tag{4.12}$$

For several corrections retain their total debit at each anchor and every mutual cross term. Strictness survives a positive remaining strict anchor or a positively weighted strict new pressure. All component sums below concern projections of the same original x.

For $P_i = e_1 + \cdots + e_i$, increasing concave interpolation through $(0,0)$ and actual floors $\langle x, P_i \rangle > f_i$ gives a reference with successive slopes as its coordinates. Its coefficients in the P_i are nonnegative slope decreases, and its demand equals its square. Deleting a knot between intervals of lengths a, b and slopes s_1, s_2 loses $ab(s_1 - s_2)^2/(a+b)$. For unit adjacent spacings this is half the squared slope difference. For the $\sqrt{i}-$constant floors at $i \geq 5$, such a unit-neighbor loss is $< 1/3200$; nonadjacent omissions have disjoint intervals and their losses add. End omissions use their exact chord costs. An omitted floor has coefficient zero.

4.2 Component estimates and the complete rank cover

Among the spectator-uniform component bounds of Addendum C are

Component	$e_T >$	Component	$e_T >$
ordinary A_1	$1/10$	ordinary $I_2(m)$	$1/7$
A_3	$1/4$	A_4	$2/7$
A_5, D_5	$3/8,\ 5/14$	E_6, E_7, E_8	$9/20,\ 5/9,\ 2/3$
A_{12}	$13/20$	A_{15}	$413/500$
A_{16}	$199/250$	A_{17}	$6/7$

An ordinary component excludes the explicitly classified short-wall defects. The exceptional rank-one type I is an actual isolated short dotted opposite wall. The typed rank-two exceptions are N_7, an old A_2 with a new isolated native-seven opposite, and O_7, an old $I_2(7)$ with a new isolated native-three opposite. Each has energy greater than .09. Their compatibility and the separate coupled rank-one exception are proved using the actual walls in Addendum C. A diagram label alone does not specify a defect.

Proposition 4.3 (Compact rank cover and its consumers)**.** *At an original farthest vertex in dimension* 18*, the necessary cover is the following. Here* $T_2 = I_2(m)$*,* $m \geq 3$*, and* $T_4 \in \{A_4, B_4, D_4, F_4, H_4\}$ *are ordinary components.*

Dominant component	*Retained spectators*
A_{12}	$I + (A_5$ *or* $D_5)$*, or* $(N_7$ *or* $O_7) + T_4$
A_{15}	A_3*,* $T_2 + A_1$*,* $I + T_2$*, or* $O_7 + A_1$
A_{16}	T_2*,* $2A_1$*,* $I + A_1$*, or* N_7/O_7
A_{17}	*one ordinary* A_1*, or* I
A_{18}	*none*
B_{16}, D_{16}	N_7/O_7
B_{17}	I
B_{18}	*none*
D_{17}	*one ordinary* A_1*, or* I
D_{18}	*none*

The complete row arguments exclude every entry except whole D_{18}*. Independent actual-rank covers exclude dimensions* $19, \ldots, 29$*; Vinberg's theorem excludes* $n \geq 30$*.*

Accounting and proof location. Call A_r, $r \geq 7$, and B_r, D_r, $r \geq 9$, dominant. Each costs more than $1/2$, so at most one occurs. Without one, the ordinary rank-over-14 baseline, allowing the unique

actual defect, exceeds one. Thus exactly one occurs. Partition the remaining rank using the ordinary and joint defect bounds, and then apply the actual-rank chain and prefix refinements. This gives precisely the displayed list. For example,

$$e_{A_{12}} + e_{I+(A_5/D_5)} > 13/20 + 5/14 = 141/140,$$

and $A_{15} > 413/500$ excludes its ordinary rank-three spectators and its short-wall alternatives using the joint bounds.

The complete accounting, the local continuation at each permitted rank, and the consumer for each retained row are printed under "The remaining spherical diagrams" in Addendum C. Whole A_{18} and B_{18} have separate endpoint and terminal-wall arguments. The higher-rank covers and the whole A_{19}, B_{19}, D_{19} proofs occur in Part V and are joined in its final complete conjunction. Every cover is necessary; no realization of its entries or dimension inheritance is used. □

4.3 The last whole component

The remaining whole D_{18} case illustrates the final geometric obstruction. Use $P_i = e_1 + \cdots + e_i$, and retain the signed coordinate in the two genuine fork duals. Addendum C's complete head and fork inventories, with their funded references, give the following terminal statements.

Proposition 4.4 (Whole-D_{18} terminal estimates)**.** *All finite and parallel head cases are excluded. For a dotted head there is a single legally funded reference supported in the first* 16 *coordinates with squared-demand gain greater than* .992*. Both signed fork heights* $\langle x, P_{17} \pm e_{18}\rangle/\sqrt{18}$ *are less than* .876*. The remaining fork types are isolated, near-three and far-three; the six unordered pairs are covered by the complete terminal classification. An actual near/far fork edge has cosine* $C = \cosh L < 5.6$*.*

Proof location and the last comparison. The four original reference cases and their individual anchor prices are given in Addendum C's "Bounds for the head and fork edges." The local A_{18} lemma at the adjacent actual vertex, with comparison factor 11/8, gives the short-edge conclusion. It uses the original maximum to compare distances at that vertex. The full head classification includes every finite label and the unbounded dotted parameter range.

For completeness, the final near/far calculation can be printed briefly. Let Ω_i be the spherical duals of the incident unit normals, so $\langle \Omega_i, n_j\rangle = \delta_{ij}$. Write the two actual opposite walls as

$$c_N = Np - \gamma\Omega_f - \Omega_o/2, \qquad c_F = Fp - \delta\Omega_o - \Omega_1/2,$$

where f, o are the distinct signed forks. Their time squares and edge cosines satisfy

$$N^2 = 9\gamma^2 + 8\gamma + 5/4, \quad F^2 = 9\delta^2 + \delta - 1/2,$$
$$C_N = (18\gamma + 8)/\sqrt{19}, \qquad C_F = (18\delta + 1)/\sqrt{19}.$$

The bound $C_N, C_F < 5.6$ gives $\gamma < .912$, $\delta < 1.301$. Put $U = 3\gamma + 4/3$, $V = 3\delta + 1/6$, $a^2 = 19/36$. Then $N^2 = U^2 - a^2$, $F^2 = V^2 - a^2$, and

$$(UV - a^2)^2 - N^2F^2 = a^2(U - V)^2 \geq 0, \qquad UV - a^2 > 0.$$

Consequently the actual mutual magnitude satisfies

$$\begin{aligned} -\langle c_N, c_F\rangle &= NF - 8\gamma\delta - \gamma/2 - (9/2)\delta - 1/4 \\ &\leq \delta(\gamma - 1/2) - 5/9 < 1.301(.412) - 5/9 < 0, \end{aligned}$$

contradicting acuteness. If $\gamma \leq 1/2$, the preceding upper bound is already negative. The other five fork pairs and both signed assignments are excluded in the complete head/fork proof of Addendum C. □

Propositions 4.3 and 4.4 exclude all $n \geq 18$, proving Theorem 1.1(ii).

5 The compact arithmetic bound

5.1 The actual arithmetic field

Arithmeticity supplies an admissible quadratic-form model over the actual totally real field K [11, Lemma 5.2]. Root lines are K-defined, invariant cyclic products are algebraic integers, and the nonphysical real forms are positive definite. For roots r_i put $a_{ij} = 2\langle r_i, r_j\rangle/\langle r_i, r_i\rangle$. The cycles in the a_{ij} are used without assuming that physical unit roots are themselves K-valued or that disconnected root lines have the same squareclass.

If an actually permitted rescaling defines an indefinite principal Gram over $k \subset K$, an additional embedding fixing k would preserve its indefiniteness. Nonphysical positivity therefore forces $K = k$. Norms of nonzero algebraic integers, together with bounds at every nonphysical embedding, give the degree reductions. The golden-field branch uses Bugaenko's published lattice theorem [12] after the invariant-lattice identification proved in Addendum A. The rational field is excluded by anisotropy and Meyer's theorem in the dimensions at issue [13].

5.2 Paying all isolated constraints together

The arithmetic component estimates use a common analytic step. It explains why overlapping local pressures can be combined without optimizing different vectors independently.

Lemma 5.1 (Simultaneous absorption)**.** *Let n_i be incident unit normals, ω_i their duals, and $w = \sum_i \beta_i\omega_i$ with $\beta_i = \langle w, n_i\rangle \geq 0$. Suppose $E_w = \|w\|^2 = \sum_i \beta_i H_i$. Outside a set J, assume $\langle x, \omega_i\rangle \geq H_i$. For $i \in J$ suppose $\beta_i > 0$, $0 < u_i < 1$, and*

$$\langle x, \omega_i - u_i n_i\rangle \geq H_i - \delta_i, \qquad 2\delta_i < \beta_i(2u_i - u_i^2).$$

Then $\|x\|^2 \geq E_w$, strictly if J is nonempty or some positively weighted retained pressure is strict.

Proof. Use the single vector $Z = w - \sum_{i\in J} \beta_i u_i n_i$. The pressures and $\langle n_i, n_j\rangle \leq 0$ for $i \neq j$ give

$$\langle x, Z\rangle \geq E_w - \sum_{i\in J} \beta_i\delta_i, \qquad \|Z\|^2 \leq E_w - \sum_{i\in J} \beta_i^2(2u_i - u_i^2).$$

Substitution in (4.11) proves the assertion, including its strictness. In particular adjacent isolated sites are allowed. □

For an actual cap isolated within its component in the retained range $0 < u < 1$, retain its full foreign projection and write

$$c = Ap - \gamma\omega_i - v_{\mathrm{for}}, \quad Q = \|v_{\mathrm{for}}\|^2, \quad u = (1 - Q)/\gamma^2 \in (0, 1), \quad D_i = \|\omega_i\|^2.$$

Positive clearing gives the actual vector

$$\eta = c + v_{\mathrm{for}} + \gamma u n_i = \gamma\{\sqrt{D_i - u}\, p - (\omega_i - u n_i)\},$$
$$\eta^2 = -\gamma^2 u(1 - u).$$

It is future pointing, zero against the contacted foreign block and c, negative against n_i, and nonpositive against every other wall. Lemma 4.1 therefore gives

$$\langle x, \omega_i - u n_i\rangle \geq \sqrt{D_i - u} - \sqrt{u(1 - u)}.$$

The loss from $\sqrt{D_i}$ is at most $a_i u + \sqrt{u(1-u)}$, where $a_i = (\sqrt{D_i} + \sqrt{D_i - 1})^{-1}$ in the chain ranges used. Thus a reference with $H_i + \ell < \sqrt{D_i}$, positive prices $\beta_i \geq b$, and $a_i \leq a$ absorbs every such foot whenever

$$au + \sqrt{u(1-u)} - \frac{b}{2}(2u - u^2) < \ell \qquad (0 < u < 1).$$

The actual cap inventories, golden-helper exclusions and numerical reference coefficients are catalogue item A0. Their complete proofs are in Addendum A; the lemma proves the shared analytic combination used by those component estimates.

5.3 The finite reductions

The compact theorem already gives $n \leq 17$. Addendum A constructs fresh necessary covers with fourteen groups at rank 17 and nineteen at rank 16. Their geometric consumers use the same original vector (4.1); arithmeticity strengthens the permitted wall parameters in those consumers.

Proposition 5.2 (Arithmetic reductions)**.** *At rank* 17*, every group except whole* A_{17} *is excluded. The two endpoint opposites of that component are isolated. Their integral generators* U, V *lie in the same actual field and satisfy*

$$36/17 < U, V < 3969/425, \qquad 0 < U^\sigma, V^\sigma < 36/17 \quad (\sigma \neq \mathrm{id}). \tag{5.1}$$

The field is $\mathbb{Q}(\sqrt{2})$*,* $\mathbb{Q}(\sqrt{3})$ *or one of the cubic fields generated by the largest roots of*

$$X^3 - 5X^2 + 6X - 1, \qquad X^3 - 6X^2 + 9X - 3.$$

At least one endpoint is finite. In rank 16*, all groups except whole* D_{16} *are excluded after the finite propositions below and their separate geometric consumers are applied.*

Proof location. The full covers and their row-by-row implications are printed in Addendum A under "The complete rank-seventeen reduction," "The complete rank-sixteen reduction," and the final arithmetic dispatches in Part VI. The whole A_{17} endpoint, field and finite-anchor proofs include every own and foreign-wall case. They prove the stated domains before any finite algebraic calculation is invoked. The whole B_{16}, B_{17}, D_{17} cases have distinct inclusions and consumers for their computed domains. □

For A_{17}, write $Q(X) = X(17X - 36)$. Necessary endpoint conditions are

$$R = Q(U)Q(V) \in K^2, \quad g = (UV - \sqrt{R})/18 \in \mathcal{O}_K, \quad L = g^2/(UV) \in \mathcal{O}_K. \tag{5.2}$$

The square root is physically positive, $g \leq 0$, and for $g \neq 0$ one has $0 < L^\sigma < 4$ at every other embedding. The zero case is retained. Physical parameters must be actual Coxeter parameters.

The four finite propositions used in the arithmetic argument have the following complete recorded outputs. Their full domains are printed in Addendum A's "Finite propositions and their complete algorithms."

Domain	Catalogue item	Output
quadratic A_{17}	A1	empty
cubic A_{17} anchors	A2	empty
shared quadratic B/D_{17}	A3	two pairs, in $\mathbb{Q}(\sqrt{13})$ and $\mathbb{Q}(\sqrt{17})$
whole quadratic B_{16}	A4	$U = 5 + 2\sqrt{6}$, $V = 10 + 4\sqrt{6}$

The survivors are necessary algebraic models; their actual geometric consumers are included in Proposition 5.2.

The full domains, maximal-order coefficient bounds, exact field-square tests and computational proofs of A1–A4 are stated in the GitHub proof catalogue and proved in the accompanying repository. The actual geometric inclusion is a separate assertion in Proposition 5.2; an algebraic survivor is not a realized polytope.

Equations (5.1)–(5.2) and the two empty endpoint domains exclude rank 17. It remains to exclude the last whole D_{16} configuration.

5.4 The head and the two genuine forks

In the original whole-D_{16} coordinates put $P_j = e_1 + \cdots + e_j$, $S_j = x \cdot P_j$ and $f_j = \sqrt{j} - .61/\sqrt{2}$.

Proposition 5.3 (Whole-D_{16} reduction)**.** *For the remaining actual diagram the field is quadratic, and the head and the two signed fork opposites are isolated and dotted. There are integral generators $U, F, G \in \mathcal{O}_K$ such that*

$$4 < U < 98/9, \quad 0 < U' < 2, \qquad 4 < F, G < 121/2, \quad 0 < F', G' < 1/2, \tag{5.3}$$

$$UF/4,\ UG/4,\ FG/4 \in \mathcal{O}_K. \tag{5.4}$$

Their time squares are

$$T_H = U(U-2), \quad T_F = 2F(2F-1), \quad T_G = 2G(2G-1), \tag{5.5}$$

and every pairwise product of these is a square in K. The original vector satisfies $S_j > f_j$ for $5 \le j \le 14$. The positive prefix reference w has $x \cdot w > E_{14} = \|w\|^2 > .9084$. The reference's first-five mean is less than .361, its remaining coefficient at P_{14} exceeds .13, and $S_6 = x \cdot P_6 > 2.01815$, $f_{14} < 3.311$.

The proof in Addendum A's "The last actual diagram" establishes the rank-16 wall inventory, simultaneous coefficient bounds, field descent and complete isolated endpoint reduction before the following dyadic argument. Both fork directions are retained even if their scalar parameters coincide.

Put $N_H = \mathrm{N}\, U$ and $A = -\mathrm{N}(U-2) = 2\,\mathrm{Tr}\, U - 4 - N_H$. For a fork put $n_F = \mathrm{N}\, F$, $S = \mathrm{Tr}\, F$ and $B = 2S - 4n_F - 1 > 0$, which is odd. The strict strips give

$$1 \le N_H \le 21, \quad 1 \le A \le 17, \quad 1 \le n_F \le 30, \quad 5 \le S \le 60,$$
$$\mathrm{N}\, T_H = -N_H A, \qquad \mathrm{N}\, T_F = -4n_F B, \qquad \Delta < (98/9)^2 < 119.$$

At a prime $\mathfrak{p}$ above two let $e = v_{\mathfrak{p}}(2)$, $r = v_{\mathfrak{p}}(U)$, $s = v_{\mathfrak{p}}(U-2)$, $k = v_{\mathfrak{p}}(F)$ and $l = v_{\mathfrak{p}}(G)$. Quarter integrality and the time squares imply

$$r + k, r + l, k + l \ge 2e, \qquad r + s + e + k \equiv r + s + e + l \equiv 0 \pmod 2. \tag{5.6}$$

Also $s = r$ for $r < e$, $s = e$ for $r > e$, and $s \ge e$ for $r = e$.

Proposition 5.4 (Dyadic terminal alternatives)**.** *The necessary integral conditions reduce any actual remaining configuration to one of the following two possibilities:*

(a) *$K = \mathbb{Q}(\sqrt{6})$, $U = 5 + 2\sqrt{6}$, and $F = G = 4U$;*

(b) *$K = \mathbb{Q}(\sqrt{57})$, $U = (9 + \sqrt{57})/2$, and $F = G = 23 + 3\sqrt{57}$.*

Integer-strip argument and proof location. Split first by the parity of N_H and then by the inert, ramified or split behavior of two. Equation (5.6) and the norm bounds leave bounded valuation possibilities. For example, if $F/4$ is integral, then $\mathrm{N}\,F = 16$; putting $W = F/4$ gives $\mathrm{N}\,W = 1$, $8 < W < 121/8$ and $\mathrm{Tr}\,W \in \{9, \dots, 15\}$. These seven traces and the discriminant bound give the complete unit-fork list. All other branches are treated by the full integer-order strips in Addendum A's "Odd head norm and inert two," "Even head norm with ramified two" and "The complete split branch." The split head list has seventeen rows. Its valuation and full field-square tests leave only (b); the odd-norm branch leaves only (a). The remaining inert, ramified and even-trace split cases are excluded by the printed integer-strip arguments. This is the manual dyadic proof, with its original geometric terminal cases retained, and requires no whole-D_{16} census. □

Both possibilities contradict the norm bound for the original vector. In (a) the actual head and fork edge cosines satisfy

$$C_H = (2+\sqrt{6})/2 > 11/5, \qquad C_+ = C_- = 2(2+\sqrt{6}) > 44/5.$$

Equation (4.6) gives $x_1 > .61$ and, for both signs,

$$x \cdot (P_{15} \pm e_{16}) > 4\sqrt{39}/7 > 3.56.$$

Averaging gives $S_{15} > 3.56$. Interpolating at indices $1, 6, 15$ with $S_6 > 2.01815$ yields

$$\|x\|^2 > .61^2 + \frac{1.40815^2}{5} + \frac{1.54185^2}{9} = 1.032821887 > 1.$$

In (b), $U > 8$ gives $x_1 > 1/\sqrt{3} > .577$. Rebuilding the first-five block of the reference in Proposition 5.3, with its sum fixed, gives gain greater than

$$E_{14} + \frac{5}{4}(.577 - .361)^2 > .96672.$$

If either actual fork had $\cosh L \geq 9$, its signed pressure would exceed 3.577. The separately available coefficient .13 at P_{14}, with residual square two and zero cross term with the first-block correction, would then give

$$\|x\|^2 > .96672 + .26(3.577 - 3.311) - .0338 = 1.00208 > 1.$$

Thus both fork cosines are below nine. In the isolated fork normalization $C_F^2 = 2F$, so $F < 81/2$, whereas $F = 23 + 3\sqrt{57} > 44$. This excludes (b).

There is no whole D_{16} configuration. Together with Proposition 5.2, this excludes ranks 16 and 17. The independent compact theorem supplies the preceding ceiling 17, and Theorem 1.1(iii) follows.

6 GitHub proof catalogue and reproduction

The versioned repository is the mathematical supplement to this concise edition. Its full manuscript [8] contains the complete proofs of every local proposition cited above. The following guide locates the essential supplementary material.

Result	Complete proof addenda	Reproduction guide
finite volume 57	Parts I, IV; intrinsic weights, coverage, finite maximum, cusp comparison and both tails	`docs/finite_volume57_reproduce.md`
compact 17	Parts II, V; actual component estimates, own-rank covers and all endpoint/head/fork cases	`docs/compact17_consolidated_proof_map_20261025.md`
compact arithmetic 15	Parts III, VI; field reductions, four finite propositions, rank covers and the manual dyadic finish	`docs/compact_arithmetic_consolidation_proof_map_20261025.md`

Each path is relative to the fixed repository version cited above. The accompanying `ADDENDA.md` maps the propositions of this edition to exact labels in that source. The package includes the algorithm specifications, exact checkers, available certificates, execution records and instructions for serial reproduction.

The GitHub proof catalogue is `arxiv/hyperbolic_coxeter_dimension_bounds/ADDENDA.md`. It assigns FV1–FV4 to the finite-volume propositions, C1–C2 to the compact case results, and A0–A6 to the arithmetic component, finite-domain and terminal results. Every item identifies its complete statement and proof in the fixed source [8], together with the corresponding checker and execution records. The catalogue and proof maps distinguish written arguments, exact computations and selected conditional Lean kernels. Those kernels do not formalize all geometric and literature inputs.

The availability and replay instructions are in `repro/PUBLIC_DATA.md`; not all separately indexed large inputs are distributed. This edition uses the retained proof and computation records and introduces no new mathematical or Lean execution.

Acknowledgments. The author used OpenAI's GPT-6-astra and GPT-6.1-sol as research and coding assistants to investigate proof ideas, help strengthen the results, improve the exposition, and generate and audit the Python and Lean 4 code accompanying this work. The author reviewed the resulting mathematics and code and takes responsibility for the final content.

References

[1] M. N. Prokhorov. The absence of discrete reflection groups with noncompact fundamental polyhedron of finite volume in Lobachevskii space of large dimension. *Mathematics of the USSR-Izvestiya*, 28:401–411, 1987. `https://www.mathnet.ru/eng/im1484`.

[2] A. G. Khovanskii. Hyperplane sections of polyhedra, toroidal manifolds, and discrete groups in Lobachevskii space. *Functional Analysis and Its Applications*, 20:41–50, 1986. `https://www.math.utoronto.ca/askold/1986-Faa-1-english.pdf`.

[3] E. B. Vinberg. The absence of crystallographic groups of reflections in Lobachevsky spaces of large dimension. *Trudy Moskov. Mat. Obshch.*, 47:68–102, 1984. English translation, Trans. Moscow Math. Soc. (1985), 75–112. `https://www.mathnet.ru/eng/mmo444`.

[4] E. B. Vinberg. Hyperbolic reflection groups. *Russian Math. Surveys*, 40(1):31–75, 1985. doi:10.1070/RM1985v040n01ABEH003527.

[5] R. E. Borcherds. Automorphism groups of Lorentzian lattices. *Journal of Algebra*, 111:133–153, 1987. `https://math.berkeley.edu/~reb/papers/al/al.pdf`.

[6] Lizi Guo and Fangting Zheng. Dimension bounds for finite-volume hyperbolic Coxeter polytopes. arXiv:2611.01795v1, 2026. https://arxiv.org/abs/2611.01795.

[7] Alexander Kolpakov. Corner payments and a dimension bound of 57 for finite-volume hyperbolic Coxeter polytopes. Public repository manuscript, 21 October 2026, 2026. Initial public snapshot 32b68355. https://github.com/sashakolpakov/CoxeterPolytopes/blob/32b68355cdb13ca8629f3fa8b98d424dd70472c7/manuscripts/finite_volume57.tex.

[8] Alexander Kolpakov. Dimension bounds for hyperbolic Coxeter polytopes: complete proofs and addenda. Repository edition of 30 October 2026, snapshot 36eeaddf, 2026. Full manuscript, Parts I–VI. https://github.com/sashakolpakov/CoxeterPolytopes/blob/36eeaddf646eff25d4990c9ab730a20ada21f80f/arxiv/hyperbolic_coxeter_dimension_bounds/main.tex. The matching PDF and reproduction guides are linked from the repository README.

[9] A. Felikson and P. Tumarkin. On simple ideal hyperbolic Coxeter polytopes. arXiv:math/0502413. https://arxiv.org/pdf/math/0502413.

[10] V. V. Nikulin. On ground fields of arithmetic hyperbolic reflection groups. In *Groups and Symmetries*, volume 47 of *CRM Proceedings and Lecture Notes*, pages 299–326, 2009. Corrected version arXiv:0708.3991v2. https://arxiv.org/pdf/0708.3991v2.

[11] I. Agol, M. Belolipetsky, P. Storm, and K. Whyte. Finiteness of arithmetic hyperbolic reflection groups. *Groups, Geometry, and Dynamics*, 2:481–498, 2008. https://doi.org/10.4171/GGD/47.

[12] V. O. Bugaenko. Groups of automorphisms of unimodular hyperbolic quadratic forms over $\mathbb{Z}[(\sqrt{5}+1)/2]$. *Vestnik Moskov. Univ. Ser. 1 Mat. Mekh.*, pages 6–12, 1984. No. 5. https://www.mathnet.ru/eng/vmumm3408.

[13] M. Belolipetsky. Arithmetic hyperbolic reflection groups. https://arxiv.org/abs/1506.03111.