Module A: Construction and classification of two-group polytopes

Application A of the marked-group umbrella theory

Private scientific draft, 10 October 2026. This volume is being independently challenged and is not an admitted replacement or a measured compression claim. It cites the companion umbrella manuscript **U** for general algebraic proofs. The exact U theorem signatures are listed in §1; the companion ledger binds the complete second author checkpoint of U, not just the cited statements.

The same local data can serve two different purposes: building a group with prescribed sections, and listing every group of a bounded order. In the first direction, small kernels are attached to subsets of the generator labels. In the second, a common nilpotent quotient reduces the search to normal subgroups of one finite group. This volume develops both directions for two-group polytopes and connects them through a class-two vertex-and-edge model.

The applications recover the declared principal results of five sources: Hou–Feng–Leemans on prescribed 2-power orders [HFLa] and on high-type rank-three groups [HFLb], Hou–Schulte on the common quotient [HS26], Cunningham–Feng–Hou–Schulte on prescribed square sections and extensions [CFHS25], and Gomi–Loyola–De Las Peñas on order 1024 [GLP18]. The bundle spans 2018–2026 in its specified editions. Earlier published geometric and cyclic-core theorems remain explicit inputs. The source results are attributed individually in §9; no source is credited merely because its terminology occurs here.

The central conclusions are these. Every finite marked two-group with a forest diagram and the intersection property has the same class-two quotient for that diagram (§2). The square lattice supplies a single family of local kernels (§3); U's supported-kernel theorem then gives prescribed edge orders, consecutive square sections and alternating assembly (§4). A bounded cyclic core reduces the high-type rank-three classification to eight explicit models (§5). Exact central descent and one invariant-kernel census complete the order-1024 classification, including presentation recognition and named facets (§§6–7). The free-extension conclusions use U's amalgam theorem and precisely stated geometric constructions (§8).

The mathematical text and its complete certificate artifacts form one charged proof package. The programs and all defining presentation data are external appendices, rather than repeated source code in the exposition. They are not unpriced black boxes: Appendix C specifies every input, algorithmic assumption, exhaustion argument, required check, output and replay command. Acquisition of U is charged once and every use of it in A has explicit hypothesis obligations. No page target, eventual twelve-paper bundle, or numerical saving is asserted before the complete package is rendered and reviewed.

1. Markings, geometric inputs and the interface to U

A marking is an ordered generating family of nonidentity involutions r_i, indexed by a finite set I. Write $G_J = \langle r_j : j \in J \rangle$. The intersection property is $G_J \cap G_K = G_{J \cap K}$ for every pair of subsets. A string C-group also has $[r_i, r_j] = 1$ for $|i - j| > 1$; nondegeneracy means that consecutive generators do not commute. The type is their consecutive product orders. Use $[x, y] = x^{-1}y^{-1}xy$, $x^y = y^{-1}xy$, and D_8 for the dihedral group of order eight. Ordered marked isomorphisms preserve every generator. The [GLP18] census instead identifies generator sets; for a nondegenerate string this adds precisely reversal.

We retain the regular-polytope/string-C-group correspondence, the marked epimorphism/cover correspondence and the coset description of faces and stabilizers. A deletion sends selected markers to themselves and all others to one. Its geometric interpretation gives the flat amalgamation property (FAP) at the corresponding initial or final cut. Flatness itself is a separate incidence conclusion: a factorization of the group by two relevant face stabilizers makes every face of the first kind incident with every face of the second kind. It is never inferred merely from IP.

The general facts used from U are listed with provisional exact signatures. Their complete proofs occur once in U, not in the application proof cost again.

- **U.K**: if a marked epimorphism $G \to B$ has kernel N and the base has IP, then G has IP exactly when the profiles $N \cap G_J$ have the same subset-intersection property.
- **U.Q**: for a quotient $q : G \to G/N$, $q(A) \cap q(B) = q(A \cap B)$ holds exactly when $N \cap BA = (N \cap B)(N \cap A)$. In particular $N \cap BA = 1$ preserves the intersection and embeds both ends; if $N \leq A$, then $q(A) \cap q(B) = q(A \cap B)$. The latter still requires checking that the displayed quotient marking on A is valid.
- **U.D**: a full Boolean family of genuine deletions implies IP and split parabolic retractions, including infinite groups.
- **U.SK**: exact base deletions and epimorphisms $E_S \to B_S$ injective on every proper-support parabolic give the full supported fiber group, with kernel profile $\ker \pi \cap H_J = \prod_{S \subseteq J} \ker(E_S \to B_S)$. It has IP, inherits Boolean deletions, and has the resulting parabolic order formula. Antichain supports embed their full local groups. If every local kernel is $\gamma_c(E_S)$ and B has class below c, the global kernel is $\gamma_c(H)$. A supported separator gives $H = H_J H_K$ when $B = B_J B_K$.
- **U.Nilpotent/U.Ladder**: for an involutory finitely generated presentation F with deletions, its class-two quotient Q inherits deletions and IP. Finite marked quotients with that same Q and order $|Q|2^k$ are classified by suitable kernels inside $\gamma_3(F/\gamma_{k+3}(F))$. Their IP is the exact all-parabolic kernel-profile condition. Every invariant index-two step is a hyperplane preimage in $M/(\Phi(M)[M, U])$, and iterating k steps exhausts every index-2^k invariant kernel.
- **U.Phi**: an irredundant generating marking of a finite two-group maps to a basis of its Frattini quotient; every binary coordinate assignment gives a character. All marked parabolics have the corresponding coordinate images.
- **U.Amalgam**: p-primary finite torsion is preserved by subgroups, finite p-group extensions and amalgamated free products. Proper embeddings in both factors make an amalgam infinite; finitely generated factors generate it.
- **U.Order**: finite normal factors and a finite quotient-index bound give the product upper bound; a marked finite quotient attaining that bound identifies the complete presentation.

U retains Burnside's basis theorem, lower-central graded Lie theory, finite nilpotency of finite two-groups and amalgam normal forms. Here we additionally retain the Gaussian-integer Euclidean algorithm, the precise cyclic-core theorem in §5 and the geometric assembly/extension theorems in §§4 and 8. The standard low-rank string-IP criteria, smooth square-quotient classification, universal geometric interpretation and exceptional infinite decomposition quoted by [GLP18] are stated where used. Exact GAP group and coset algorithms and NQ's universal nilpotent quotient algorithm are computational inputs. No finite classification catalogue supplies search completeness.

2. A canonical quotient and one finite classification problem

The application-specific bridge to U is a rigidity statement: a forest diagram forces independent edge commutators in degree two. Its proof uses actual IP in G; no intermediate quotient is presumed to retain IP.

Lemma A2.1 (descending lift). Let G be finite nilpotent and let $A \cap B = 1$. Put $D_j = \gamma_j(G)$, $A_j = A \cap D_j$, $B_j = B \cap D_j$. If the images of A_j, B_j generate D_j/D_{j+1} for every $j \geq m$, then $D_j = A_j B_j$ for every $j \geq m$.

Proof. Descend from a trivial term. Assuming $D_{j+1} = A_{j+1}B_{j+1}$, if $a \in A_j, b \in B_j$ have equal images in the next quotient, write $ab^{-1} = a'b'$ with $a' \in A_{j+1}, b' \in B_{j+1}$. Then $a'^{-1}a = b'b \in A \cap B$, so both images are trivial. Their images therefore form an internal direct sum in the abelian quotient. Counting gives $|D_j| = |A_j||B_j| = |A_jB_j|$, proving equality as sets. No prior normality or permutability of A, B is assumed. ■

Lemma A2.2 (graded support). Let a positively graded Lie algebra be generated in degree one by disjoint sets X, Y, with subalgebras L_X, L_Y. If $[x, y] \in L_Y$ for $x \in X, y \in Y$, then $L = L_X + L_Y$,

degree by degree.

Proof. The derivation ad x preserves L_Y because it does so on its generators. Iterated brackets of those derivations also preserve L_Y, by Jacobi. Hence $[L_X, L_Y] \subseteq L_Y$, and $L_X + L_Y$ is a Lie subalgebra containing all generators. The argument holds in characteristic two. ■

Theorem A2.3 (forest quotient). Suppose G is a finite two-group generated by involutions r_i, with marked IP, and its noncommuting pairs form a forest T on d vertices and m edges. Then edge commutators are independent in $G'/\gamma_3(G)$, and $G/\gamma_3(G)$ is the marked group

$$(v, z)(w, t) = (v + w, z + t + \beta(v, w)), \quad v, w \in \mathbb{F}_2^d,\ z, t \in \mathbb{F}_2^m, \tag{A-F}$$

where, after orienting each edge, its component of β is $v_{\text{head}} w_{\text{tail}}$. Its order is 2^{d+m}.

Proof. IP gives irredundancy. U.Phi gives $G/G' = C_2^d$ and $G' \cap \langle r_i \rangle = 1$. The lower-central graded Lie ring is generated by these degree-one elements; all its graded components are $\mathbb{F}_2$-spaces by bilinearity. Edge commutators span degree two. If they satisfy a nonzero linear relation, root each forest component and choose a participating edge whose child endpoint has maximal depth in its component. Removing it separates the descendant set X from all other vertices Y, and no other participating edge is internal to X. The relation therefore writes the only nonzero cross bracket as a sum of brackets internal to Y. Lemma A2.2 shows every graded component is generated by pure X- and pure Y-brackets. Their representatives lie in $G_X \cap \gamma_j(G)$ and $G_Y \cap \gamma_j(G)$.

Actual IP gives $G_X \cap G_Y = 1$, so Lemma A2.1 gives $G' = (G' \cap G_X)(G' \cap G_Y)$. For the crossing endpoints $x \in X, y \in Y$, write $[r_x, r_y] = uv$ accordingly. Then $u \in G_X \cap G_{\{x\} \cup Y} = \langle r_x \rangle$, whence $u = 1$; next $[r_x, r_y] \in G_Y \cap G_{\{x,y\}} = \langle r_y \rangle$, so it too is one. This contradicts that the edge is noncommuting. Independence follows.

Bilinearity makes (A-F) associative; its coordinate markers are involutions with independent central edge commutators. It is generated by those markers, and its parabolic on I consists exactly of vertex coordinates supported on I and edge coordinates supported on edges internal to I. Thus it has IP. Collection in the corresponding class-two presentation gives at most 2^{d+m} elements, and this coordinate model attains the bound. Independence in $G'/\gamma_3(G)$ gives the same order for $G/\gamma_3(G)$; hence the marked map is an isomorphism. ■

Corollary A2.4 (F013 1.1–1.2). For every finite nondegenerate two-group string C-group of rank $d \geq 2$, $G/\gamma_3(G) \cong G(C_d)$, of order 2^{2d-1}; the associated polytope covers C_d.

Proof. Its diagram is a path, so $m = d - 1$. Conder's marked presentation has involutions, distant commutation, adjacent order four, and adjacent squares central. The model (A-F) realizes that presentation with the maximal collected order, identifying it with $G(C_d)$. The retained marked covering correspondence gives the geometric assertion. This also proves uniqueness of the minimal object in the declared finite nondegenerate class. ■

Notice that no intermediate quotient was assumed to retain IP. Lemma A2.1 is the step that prevents that circular assumption.

Corollary A2.5 (rank and excess classify the finite search). Let T be a specified forest on d vertices and m edges, and let k≥0. Set

$$F_T = \langle x_i \mid x_i^2 = 1,\ [x_i, x_j] = 1 \text{ for nonedges} \rangle, \quad U_{T,k} = F_T/\gamma_{k+3}(F_T), \quad K = \gamma_3(U_{T,k}).$$

The ordered IP markings with graph T and order 2^{d+m+k} are exactly the quotients $U_{T,k}/N$, where N is normal, contained in K, $[K : N] = 2^k$, and the images $(N(K \cap (U_{T,k})_J))/N$ have IP for every pair J,L. Generator-set equivalence is exactly the action of Aut(T) on these kernels.

Proof. Killing any set of generators preserves every defining relator of F_T. Collection in its class-two quotient gives at most d+m binary coordinates, and (A-F) realizes all of them, so this quotient is precisely the marked model Q_T. Every target has that same quotient by A2.3. These are exactly U.Nilpotent's hypotheses. Its all-parabolic condition and U.Ladder give the stated bijection and exhaustive search. An ordered isomorphism must commute with the generating map, hence has the same kernel. A permutation of the generator set preserves the noncommutation graph and extends to an automorphism of F_T and $U_{T,k}$; this proves the last claim. ■

This is used twice below: the order-256 seed has (d,k)=(3,3), while all nondegenerate order-1024 markings have (d,k)=(3,5),(4,3),(5,1). Their actual terminal counts require the complete computation in Appendix C. The parameter theorem proves exhaustion and termination, not those output values.

3. The affine square seed

Let

$$a(x,y) = (-x,y), \quad b(x,y) = (y,x), \quad c(x,y) = (x,1-y).$$

These generate $W = \mathbb{Z}^2 \rtimes D_8$, of type $\{4,4\}$, with $ac = ca$. Its presentation is

$$W = \langle a,b,c \mid a^2 = b^2 = c^2 = (ab)^4 = (bc)^4 = (ac)^2 = 1 \rangle. \tag{A-S1}$$

Here is an algebraic normal-form proof. Put $d = bab$, $y_0 = cd$, and $x_0 = by_0b$. The relations give $[a,d] = 1$, $[a,y_0] = 1$, $dy_0d = y_0^{-1}$, $ax_0a = x_0^{-1}$, and conjugation by b swaps x_0, y_0. Substitution of $bc = x_0ab$ makes $(bc)^4 = 1$ equivalent to $[x_0,y_0] = 1$. Thus every word collects to $x_0^u y_0^v M$, $M \in \langle a,b \rangle = D_8$. In the affine model x_0, y_0 are the unit translations, so these forms are distinct. This proves (A-S1).

For $n \geq 5$, define the invariant lattice

$$\Lambda_n = \begin{cases} 2^e\mathbb{Z}^2, & n = 2e+3, \\ 2^e\langle (1,1),(1,-1) \rangle, & n = 2e+4, \end{cases} \qquad e \geq 1,$$

and put $H_n = W/\Lambda_n$. Its order is 2^n; the full linear D_8 coordinate is retained, even if its action on the vertex quotient is not faithful. Its pair orders are $4, 4, 2$.

Write $T = abcb$, $U = cbab$, translations by $(-1,0)$, $(0,1)$. The extra relators for H_n can be taken as T^{2^e}, U^{2^e} in the square case, and $T^{2^e}U^{2^e}, T^{2^e}U^{-2^e}$ in the diagonal case. After deleting any marker, both T, U have order at most two: deleting a leaves conjugates of c, deleting c leaves conjugates of a, and deleting b leaves ac, ca. Therefore every lattice relator survives every deletion. The even Coxeter relators also survive. Thus H_n has Boolean deletions and IP by U.D.

Moreover,

$$\gamma_3(W) = 2\mathbb{Z}^2, \qquad \gamma_3(H_n) = 2\mathbb{Z}^2/\Lambda_n, \qquad |\gamma_3(H_n)| = 2^{n-5}. \tag{A-S2}$$

Indeed reduction modulo $2\mathbb{Z}^2$ makes the adjacent squares central; conversely $[[a,b],c]$ and $[[b,c],a]$ give the two doubled unit translations, up to signs. Thus H_5 is class two of order 32. The squared translations are not always independent cyclic subgroups: in H_6 they coincide and have order two. Formula (A-S2) uses a lattice index and does not make the false independence assertion.

Lemma A3.1 (square-map recognition). Every finite two-group string C-group of type $\{4,4\}$ and order 2^n is marked-isomorphic to H_n; necessarily $n \geq 5$.

Proof. The two dihedral parabolics have orders eight and intersect in order two, so their product already has 32 elements. The universal group (A-S1) maps onto the given group. IP gives irredundancy, so U.Phi gives elementary abelianization of rank three. Hence word parity is a homomorphism onto C_2, and the kernel of (A-S1) lies in the subgroup of orientation-preserving plane isometries. It contains

no nonidentity torsion: a finite-order plane isometry fixes a point; the square tiling moves that point into a closed fundamental right triangle; its stabilizer is a subgroup of a vertex dihedral parabolic or an edge reflection group. Each embeds by the prescribed pair orders and IP. An orientation-preserving plane isometry without finite order is a translation. Thus the kernel is a D_8-invariant sublattice of $\mathbb{Z}^2$ of index 2^{n-3}. Identifying the lattice with an ideal of $\mathbb{Z}[i]$, its generator has norm a power of two. Unique factorization (from the Euclidean algorithm) makes it a unit times a power of $1+i$. Even powers give the square lattice and odd powers the diagonal lattice displayed above. There is exactly one such ideal at each index, proving the assertion. ■

The quotient $q_n : H_n \to H_5$ fixes the three named generators. Its kernel is $\gamma_3(H_n)$. Every proper marked parabolic has at most two generators, and its type is respectively 4,4 or2 both upstairs and downstairs. Each is the identical dihedral or order-four group; singleton and empty parabolics likewise embed. This verifies U.SK's full proper-support condition, rather than merely checking three involutions.

4. Prescribed local sections from supported kernels

Theorem A4.1 (edge inflation and exact orders). Let B have Boolean deletions and select pairs S={i,j} whose product orders q_S are even. For m_S divisible by q_S, attach the marked dihedral map $D_{2m_S} \to B_S = D_{2q_S}$. The resulting supported fiber group has IP, all deletions and

$$|H_J| = |B_J| \prod_{S \subseteq J} (m_S/q_S).$$

Every selected pair has product order m_S and every unselected pair retains its old order.

Proof. The dihedral quotient is injective on its proper parabolics, which are trivial or a single reflection. U.SK applies and its antichain clause identifies each selected local group. On an unselected pair every extra coordinate is a reflection or one; a base pair of odd order cannot have a single-reflection deletion. Thus the extra coordinates do not increase its base product order. ■

For d≥3, n≥5, k_i≥2 and $\sum k_i \le n-1$, set $\delta = n-1-\sum k_i$ and $B = C_2^{d-3} \times H_{\delta+5}$, with the elementary factors first. Its initial d−3 consecutive pair orders are 2 and its final two are 4. Attach $m_i = 2^{k_i}$. The seed supplies all deletions, so A4.1 gives

$$|H| = 2^{d-3+\delta+5} \prod_{i \le d-3} 2^{k_i-1} \prod_{i > d-3} 2^{k_i-2} = 2^n.$$

Distant markers commute in every coordinate, so the marking is string. The geometric correspondence proves **[HFLa] Theorem 1.2**. Taking d=3 proves **[HFLb] Theorem 1.2**, including its stated n≥10 range, without another construction. These existence results do not imply the classification of §5.

Theorem A4.2 (square kernels on a forest). Let T be a forest and choose distinct induced three-vertex paths S, with parameters $n_S \ge 5$. Attach $H_{n_S} \to H_5 = (Q_T)_S$ to the model (A-F). Then

$$|H_J| = 2^{|J|+|E(T[J])|+\sum_{S\subseteq J}(n_S-5)}, \qquad \gamma_3(H) = \prod_S \gamma_3(H_{n_S}), \qquad H/\gamma_3(H) = Q_T. \qquad \text{(A-P)}$$

It has Boolean deletions and IP, and each selected triple is exactly H_{n_S}.

Proof. Deleting coordinates in (A-F) gives all base deletions and its parabolic order $2^{|J|+|E(T[J])|}$. Section3 verifies every proper-support injection and the local γ3 kernel. U.SK, including its antichain and lower-central clauses, gives every assertion. ■

For the d-vertex path, take all consecutive triples with parameters n_3,…,n_d. Equation (A-P) gives the exact group order $2^{\sum n_j - 3(d-3)}$, all prescribed square sections and all FAP cuts. To prove flatness when d≥4, let L={0,…,d−2} and M={d−3,d−2,d−1}. Every support lies on one side. The vertex-and-edge counts in (A-F), together with IP, give $Q_T = (Q_T)_L (Q_T)_M$. U.SK therefore gives $H = H_L H_M$. The stabilizer of the base (d−4)-face contains H_M, while H_L is a facet stabilizer; taking inverses also gives $H = H_M H_L$. The coset face criterion proves (d−4,d−1)-flatness and hence flatness. The co-(−1)-face cut at d=3 is the tautological whole-polytope cut; the other two cuts are genuine seed deletions. This proves all clauses of **[CFHS25] Theorem 5.1**.

The universal presentation is a separate conclusion

Write G for the group H constructed in the path case above. The supported fiber group is not an arbitrary amalgam. To prove the source's universal-presentation claim, a further local normalization argument is needed.

Lemma A4.3 (four-marker centralization). Suppose a, b, c, d are involutions with $[a, c] = [a, d] = [b, d] = 1$ and $[(ab)^2, (cd)^2] = 1$. Then d commutes with $(cbab)^2$ and $(abcb)^2$.

Proof. Put $x = bab$, $E = (ab)^2 = ax$, $F = (cd)^2$. The element a commutes with F, so $[E, F] = 1$ implies $[x, F] = 1$. Also $[x, d] = 1$, hence $[x, Fd] = [x, cdc] = 1$. Conjugating by c gives $[cxc, d] = 1$. Both factors of $(cbab)^2 = (cxc)x$ commute with d. Finally $abcb = b(cbab)^{-1}b$, and b commutes with d, proving the second assertion. ■

First let P_0 have the string $[4, \dots, 4]$ relations and every prescribed triple presentation from §3. All its defining relators are stable under every deletion assignment: the string Coxeter relators have even exponents, and the local lattice relators survive every deletion of their three markers by §3. Thus the assignments define endomorphisms of P_0, and U.D proves its IP without assuming finiteness. The constructed quotient G shows that each local triple embeds with its exact prescribed marking and that every generator is nontrivial. Hence P_0 itself is a string C-group of the required local type. Every marked string C-group of that local type satisfies its defining relators, so is its quotient. The standard marked covering correspondence therefore constructs the universal regular polytope of that type directly. No separate assumption about a universal polytope's presentation or existence is needed.

Now let Q be the quotient of P_0 by

$$[(r_i r_{i+1})^2, (r_{i+2} r_{i+3})^2] = 1 \quad (0 \le i \le d-4). \tag{A-U}$$

The constructed group satisfies these relations. For (A-U), no tuple coordinate contains both disjoint edges; the edge missing one endpoint has square one in that coordinate. Hence $Q \twoheadrightarrow G$. Each triple embeds in Q, since its map onward to the embedded triple in G is injective. Its third lower-central subgroup K_i is generated by its two squared translations and has order $2^{n_{i+3}-5}$, by (A-S2). Lemma A4.3 and its reversed version show that the nearest outside markers centralize K_i, while the farther markers do so by string commutation. Thus every $K_i \lhd Q$. Their product K is finite of order at most $2^{\sum_j (n_j - 5)}$. Modulo it all triples are class two; each edge commutator commutes with its endpoints and nearest outside markers, and with farther markers by string commutation. Thus Q/K is class two. Collecting markers and central edge commutators gives $|Q/K| \le 2^{2d-1}$. Therefore

$$|Q| \le 2^{\sum_j (n_j - 5) + 2d - 1} = 2^{\sum_j n_j - 3(d-3)} = |G|.$$

U.Order, applied to the finite normal factors and the displayed quotient bound, identifies $Q = G$ as marked groups. Since P_0 is the universal type group just constructed, this is exactly its quotient by (A-U). The finite group presentation, the universal group and both intersection properties have all been proved using the same deletion calculus.

This proves **[CFHS25] Theorem 5.3**. No assertion about the order of the universal group P_0 itself is required.

Alternating assembly

We retain the following precise FAP case of Monson–Schulte's assembly theorem. If regular (d−1)-polytopes P,Q have a common facet K and FAP retractions $\Gamma(P) = N_P \rtimes \Gamma(K)$ and $\Gamma(Q) = N_Q \rtimes \Gamma(K)$, the tail-triangle group $(N_P \times N_Q) \rtimes \Gamma(K)$, with distinct final markers, gives an alternating semiregular d-polytope with these facets. Its full automorphism group is that group if P and Q are not isomorphic, and its extension by branch-swapping C2 when P=Q; in that case the polytope is regular. This is a retained geometric theorem, not a consequence of IP alone.

Apply A4.2 to the tree with common prefix and two final leaves, attaching the two lists $(n_3, \dots, n_{d-2}, n_{d-1})$ and $(n_3, \dots, n_{d-2}, m_{d-1})$. Write L,M for the facet label sets and C=L∩M. The two deletions map H into $H_L \times_{H_C} H_M$. They are jointly injective: every vertex coordinate, edge coordinate and supported kernel coordinate lies on one side and is detected there. Their image is the full fiber product, since (A-P) gives its order $|H_L||H_M|/|H_C|$. Split deletions identify it with the exact tail-triangle group in the retained theorem. For d=4 the common facet is a square; otherwise it has list $(n_3, \dots, n_{d-2})$. Its order is $2^{\sum_{j=3}^{d-2} n_j - 3(d-5)}$, including 8 for the empty list. The two facet kernels have orders $2^{n_{d-1}-3}$ and $2^{m_{d-1}-3}$. Thus the full automorphism order is

$$2^{\sum_{j=3}^{d-2} n_j + n_{d-1} + m_{d-1} - 3(d-3)}$$

when the final parameters differ, and twice that order when they agree. Unequal parameters give unequal facet orders and rule out an interchange; equal parameters give identical marked facets and their branch swap. These are exactly **[CFHS25] Theorem 6.1(a–b)**. No additional mixed co-(d−4)-section is prescribed.

5. Bounded cyclic cores and the eight high-type families

The supported constructions solve existence. Completeness at the high dihedral boundary needs a different local invariant, the cyclic core. Its bounded quotient is nevertheless recognized by the same square model, and its only finite seed is generated by the same U.Nilpotent/U.Ladder search as order 1024.

The retained **cyclic core theorem of Lucchini** says: if a cyclic proper subgroup M of a finite group H has core K, then $|M : K| < |H : M|$. Here K is the largest normal subgroup of H contained in M. We use the form recorded in Isaacs, *Finite Group Theory*, Theorem 2.20, also stated as L035 Proposition 2.9.

Theorem A5.1 (bounded cyclic quotient). Let $(G; a, b, c)$ be a rank-three string C-group of order 2^{t+s}, type $\{4, 2^t\}$, with $s \geq 3$ and $t \geq s - 2$. Put

$$r = bc, \quad m = 2^t, \quad d_0 = (ab)^2, \quad A = \langle r^{2^{s-2}} \rangle.$$

Then $A \lhd G$, and G/A is a string C-group of order 2^{2s-2} and type $\{4, 2^{s-2}\}$.

Proof. The even-word subgroup $H = \langle ab, bc \rangle$ has index at most two. IP and U.Phi give $G/G' = C_2^3$, where its image has dimension two, so the index is exactly two. For $M = \langle r \rangle$, the cyclic core theorem gives $|M : \mathrm{Core}_H M| < 2^{s-1}$, hence at most 2^{s-2}. Thus A is the subgroup of the cyclic core of its prescribed order, so is characteristic there and normal in H. The involution b inverts r; consequently $A \lhd G$. It meets $\langle a, b \rangle$ trivially by IP and lies in $\langle b, c \rangle$. Their quotient parabolics still intersect in $\langle bA \rangle$: equality of cosets gives membership in the original right parabolic, after multiplication by an element of A. The left parabolic embeds and the right has its stated dihedral order. This proves IP, independence, type and order in the quotient. ■

Since $\mathrm{Aut}(A)$ is abelian, every commutator centralizes A. In particular $[d_0, r^{2^{s-2}}] = 1$. If $A \neq 1$, its unique involution $z = r^{m/2}$ is central in G. The universal identities

$$r^a = d_0 r, \quad d_0^a = d_0, \quad r^c = r^{-1}, \quad d_0^c = d_0^r \qquad \text{(A-C1)}$$

follow from the involution, string and order-four relations.

Hereafter $s \in \{3,4,5\}$, $n = t + s \geq 10$, so m is divisible by 32. The following eight models will realize every class. Let z be a distinguished central involution of the finite group D:

Branch	Group D	Right-conjugation automorphism T, fixing z
$s = 3$	$\langle d_0, z\rangle = C_2^2$	$T(d_0) = d_0 z^\epsilon$
$s = 4$	$\langle d_0, d_1, z\rangle = C_2^3$	$T(d_0) = d_1,\ T(d_1) = d_0 z^\epsilon$
$s = 5$	Four involutions d_0, d_1, d_2, z, z central, $[d_0, d_2] = 1$, $[d_0, d_1] = [d_1, d_2] = z^\beta$	$T(d_0) = d_1,\ T(d_1) = d_2,\ T(d_2) = d_0 d_1 d_2 z^\nu$

Here $\epsilon \in \{0,1\}$, or $(\beta, \nu) \in \{0,1\}^2$. For the last row an explicit group law on four binary coordinates is

$$(u, v, w, e)(u', v', w', e') = (u + u', v + v', w + w', e + e' + \beta(vu' + wv')). \tag{A-C2}$$

Its bilinear final term proves associativity and gives all sixteen normal forms $d_0^u d_1^v d_2^w z^e$. Thus the three D-orders are four, eight and sixteen. The displayed maps preserve their involution and commutator relations and have fourth power one. For example in the last row the two nontrivial commutator contributions cancel in $(d_0 d_1 d_2)^2$, and applying T four times to d_0 returns d_0; its orbit generates D together with z.

Adjoin a cyclic r of order m, with $x^r = T(x)$, and take the central quotient

$$N = (D \rtimes \langle r\rangle)/\langle z r^{-m/2}\rangle. \tag{A-C3}$$

The identified subgroup has order two and meets both D and $\langle r\rangle$ trivially, since $T^{m/2} = 1$. Hence D embeds, r retains order m, and $|N| = |D| m/2$.

Define automorphisms $\mathcal{A}, \mathcal{C}$ by

$$\mathcal{A}(r) = d_0 r, \quad \mathcal{A}(d_0) = d_0, \qquad \mathcal{C}(r) = r^{-1}, \quad \mathcal{C}(d_i) = T^{1-i}(d_0), \quad \mathcal{C}(z) = z.$$

In the first two rows $\mathcal{A}$ fixes D; in the third it fixes d_0, d_1, z and sends d_2 to $d_2 z^\beta$. For an explicit verification, substituting the listed generator images into the defining involutions and commutators of D preserves each relation. On its generators the compatibility equalities are

$$\mathcal{A}(Tx) = T(\mathcal{A}(x)^{d_0}), \qquad \mathcal{C}(Tx) = T^{-1}\mathcal{C}(x).$$

They express precisely preservation of $x^r = T(x)$. Collection gives $(d_0 r)^4 = r^4$ in the first two rows and $r^4 z^\nu$ in the last. Since eight divides $m/2$, this preserves $r^m = 1$ and $r^{m/2} = z$. Finally $\mathcal{A}^2 = \mathcal{C}^2 = 1$ and $\mathcal{A}\mathcal{C} = \mathcal{C}\mathcal{A}$ on r, d_0, which generate N. For instance both composites send r to $r^{-1} d_0$. These checks prove the automorphism assertions for every admissible m, not only for a computational sample.

Set $G = N \rtimes \langle a, c\rangle$, where a, c are commuting involutions inducing these automorphisms, and put $b = rc$. Then a, b, c are involutions, $[a, c] = 1$, $bc = r$, and $(ab)^2 = d_0$. They generate G, of order $2^s m$, with pair orders $4, m, 2$. In G/N, the images of a, b are independent whereas the right parabolic has image $\langle cN\rangle$. Hence the only candidates for their intersection are $1, b, d_0, d_0 b$. The normal form (A-C3) gives $d_0 \notin \langle r\rangle$, leaving exactly $\langle b\rangle$. The three generators are independent (already $a \notin \langle b, c\rangle$, and the pair orders exclude the others), so this proves rank-three IP. Direct substitution yields

$$\begin{aligned} &s = 3: \quad [d_0, c] = z^\epsilon, \\ &s = 4: \quad [d_0, r^2] = z^\epsilon, \\ &s = 5: \quad [a, r^2]^2 = z^\beta, \qquad [a, r^4] = z^\nu. \end{aligned} \tag{A-C4}$$

They are invariant words of the ordered marking and distinguish the two, two and four models within their branches.

Completeness and bare presentation recognition

The eight source presentations have common relations

$$a^2 = b^2 = c^2 = (ac)^2 = (ab)^4 = r^m = 1, \qquad r = bc,$$

and the following additional relations, with $z = r^{m/2}$:

Source name	m	Extra relations
G_1, G_2	2^{n-3}	$[d_0, c] = z^\epsilon$, $\epsilon = 0, 1$ respectively
G_3, G_4	2^{n-4}	$[d_0, r^2] = z^\epsilon$, $\epsilon = 0, 1$ respectively
G_5, G_6, G_7, G_8	2^{n-5}	$[a, r^2]^2 = z^\beta$, $[a, r^4] = z^\nu$, with $(\beta, \nu) =$ $(0,0), (1,0), (0,1), (1,1)$

Theorem A5.2 (high-type completeness). Every group in the three classes of L035 Theorem 1.3 is exactly one of the marked models of §5; each displayed source presentation defines that model.

Proof, first branch. For $s = 3$, Theorem A5.1 gives a quotient of type $\{4, 2\}$ and order sixteen, necessarily $D_8 \times C_2$. Thus $h = [d_0, r] \in A = \langle r^2 \rangle$. Since it commutes with r, and d_0 centralizes A, $h^2 = [d_0, r^2] = 1$. Hence $h = 1$ or z, and (A-C1) makes this exactly $[d_0, c] = z^\epsilon$. The conjugates of d_0 obey $d_1 = d_0 z^\epsilon$; (A-C1) forces all the first model's actions and normal forms. The resulting surjection from that model onto the actual group is between groups of the same order, hence an isomorphism.

Second branch. For $s = 4$, the quotient of order 64 and type $\{4, 4\}$ is H_6, by Lemma A3.1. In that affine quotient $[d_0, r^2] = 1$, so upstairs $h = [d_0, r^2] \in \langle r^4 \rangle$. Again $h^2 = [d_0, r^4] = 1$, giving $d_2 = d_0 z^\epsilon$, where $d_i = d_0^{r^i}$. Conjugation by a fixes d_0, d_1. Applying it to the equation for d_2, with z central, gives $d_2^a = (d_1^{d_0})^r = d_2$, so $[d_0, d_1] = 1$. All second-model relations follow, and the same order comparison proves uniqueness.

The bounded seed for the third branch. The same universal-kernel theorem used for the order-1024 classification gives the following finite fact: every order-256 marked string C-group of type $\{4, 8\}$, with $r = bc, d_0 = (ab)^2$, satisfies exactly one row.

Row	Relation pattern	Ordered marked classes
I	$[a, r^2]^2 = 1,\ [a, r^4] = 1$	1
II	$[a, r^2]^2 = r^4,\ [a, r^4] = 1$	1
III	$[d_0, r^2] = r^4$	1
IV	$[(r^2)^a, r] = r^4$	1

Apply U.Nilpotent and U.Ladder, as specialized in A2.5 with d=3 and k=3. The universal class-five group has order 2^{18} and third lower-central subgroup of order 2^{13}. Its invariant index-two ladder has 7, 23 and 61 kernels. Testing all 64 subset pairs for each final candidate gives 37 ordered IP markings, four with the indicated orientation of type {4,8}. The census program in Appendix C evaluates every displayed word on all four and verifies that each belongs to exactly one row, one marking per row. Section 7 states the complete output and the shared finite-search proof. Thus this bounded input is now conditional on exact GAP/NQ algorithms, with no abstract-group catalogue. No SmallGroups identifier or catalogue multiplicity enters these relation implications.

Third-branch lifts. For $s = 5$, Theorem A5.1 gives $A = \langle r^8 \rangle$ and the table applies to G/A. In row III we would have $[d_0, r^2] = r^e$, $e \equiv 4 \pmod 8$. It commutes with r, so $[d_0, r^8] = r^{4e} = 1$, impossible for $m \geq 32$, since $v_2(4e) = 4$. In row IV put $x = (r^2)^a$. The relation would give $[x, r] = r^e$, $e \equiv 4 \pmod 8$, hence $r^x = r^{1-e}$. The unit $1 - e \equiv 5 \pmod 8$ has order $m/4$ modulo m: successive squaring gives $v_2((1-e)^{2^j} - 1) = j + 2$. But $x^{m/8} = (r^{m/4})^a \in A$, by normality, so it centralizes r, contradicting that order. Thus III and IV cannot lift.

In I and II, $[a, r^4] \in A$, so a maps r^4 to an odd power of itself and $M = \langle r^4 \rangle \lhd G$. Quotienting gives a marked $\{4, 4\}$ group of order 128, hence $H_7 = W/4\mathbb{Z}^2$. Let $N = \langle d_0, r \rangle$. Equations (A-C1) make it normal, with index four: modulo N, a, c commute, and their independence follows in the rank-three Frattini quotient, where N has the one-dimensional image spanned by r. The cyclic normal subgroup M is central in N, because d_0 is a commutator and r centralizes its own powers.

In H_7, the conjugates d_i, $0 \leq i \leq 3$, have linear part $-I$ and translation parts $(0,0), (0,2), (2,2), (2,0)$, up to the choice of coordinate directions. They generate C_2^3, with $d_3 = d_0 d_1 d_2$. Upstairs, every $[d_i, d_j]$ is in central M and has order at most two. Put $[d_0, d_1] = z^\beta$; conjugating by r gives $[d_1, d_2] = z^\beta$. Write $d_3 = d_0 d_1 d_2 u$ with $u \in M$. Conjugation invariance gives

$$[d_1, d_2] = [d_2, d_3] = [d_2, d_0][d_2, d_1],$$

so $[d_0, d_2] = 1$. Consequently $(d_0 d_1 d_2)^2 = 1$; because d_3 is an involution, $u^2 = 1$, giving $u = z^\nu$. One further conjugation gives $d_4 = d_0$. These are precisely the final row of the table in §5, and (A-C1) forces the actions of the commuting complement $\langle a, c \rangle$. The model maps onto G and has its order $32m$, so is isomorphic. This proves completeness of the eight models.

It remains to recognize the **bare presentations**, where the target order cannot yet be assumed. Let P be one of them and $z = r^{m/2}$, initially of order at most two. For G_1, G_2, if $\epsilon = 1$, a centralizes $z = [d_0, c]$ since it centralizes d_0, c; b, c centralize it since they invert r. If $\epsilon = 0$, d_0 is central; (A-C1) and the evenness of $m/2$ then make z central too. Collection gives at most $8m$ normal forms, attained by the first model.

For G_3, G_4, $d_2 = d_0 z^\epsilon$. If $\epsilon = 0$, applying a as in the second-branch argument gives $[d_0, d_1] = 1$, after which the model relations and centrality of z follow. If $\epsilon = 1$, $[d_0, r^2] = z$ implies that conjugation by d_0 inverts z; as $z^2 = 1$, it centralizes z. Thus z commutes with r and all d_i. Write $j = [d_0, d_1]$. Applying a to the equation for d_2, and using (A-C1), gives $z^a = jz$. Since a fixes d_0, d_1, it fixes j, and $a^2 = 1$ gives $j^2 = 1$. Hence d_0, d_1 centralize j. Also $j^r = [d_1, d_2] = [d_1, d_0 z] = j^{-1} = j$, and c exchanges d_0, d_1, so j is central. Collection yields $(d_0 r)^4 = jr^4$. Since $m/8$ is even, $z^a = (d_0 r)^{m/2} = z$, so $j = 1$. The second model relations now give at most $16m$ forms, all attained by that model.

For G_5–G_8, put $w = [a, r^2]$. If $\beta = 1$, conjugation by a inverts w, and so fixes $z = w^2$. If $\nu = 1$, it inverts $[a, r^4] = z$, again fixing z. If both are zero, a centralizes r^4 and therefore z. Thus z is central in every case, and $[a, r^4] = z^\nu$ makes $M = \langle r^4 \rangle$ normal of order at most $m/4$. Modulo M, the presentation is W with the extra relation $[a, r^2]^2 = 1$. In the affine model $[a, r^2]$ is a doubled unit translation, so its square and conjugates kill precisely $4\mathbb{Z}^2$. Thus the quotient has order 128, and $|P| \leq 128(m/4) = 32m$. The constructed model attains this bound and satisfies every relator, so the order comparison principle identifies P with it. This completes presentation recognition and the theorem. ■

The preceding theorem proves the full marked interpretation of L035 Theorem 1.3; the listed word invariants distinguish all the declared families. It does not classify arbitrary two-groups of order 2^n, other types, or the source's unnumbered exceptional $n < 10$ discussion.

6. Central descent and complete binary lifts

This branch concerns the principal nondegenerate classification in Gomi–Loyola–De Las Peñas [GLP18], with the marking retained. Its equivalence sends the distinguished generator set onto the other set. For

a connected string diagram this permits precisely the two orders related by reversal. It does not identify arbitrary markings of an abstractly isomorphic group. The rank-two case is the unique dihedral marking of order 1024. The three higher-rank totals below are 125, 105 and 8; all stated types, facets and universal orders are part of the conclusion.

The specifically retained prior results in this branch are the standard string intersection criteria quoted as [GLP18] Lemmas 2.2–2.4, the universal geometric interpretation quoted as its Theorem 4.2, the finite smooth square-quotient classification quoted as its Lemma 5.1, and the exceptional infinite universal decomposition quoted as its Theorem 5.3. Their relevant precise forms are stated below. Exact finite-group and complete coset-enumeration algorithms in GAP, and NQ's universal nilpotent quotient algorithm, are computational inputs. The Atlas supplies definitions of named marked presentations, all included in the defining-data appendix; neither its completeness nor a SmallGroups catalogue is assumed.

6.1. Preliminary adapters and exact quotient intersections

Proposition N1 (endpoint cosets). A finite string C-group of rank d and adjacent orders $m_1,\dots,m_{(d-1)}$ has order at least $2\prod m_i$. In particular, a nondegenerate string two-group of order 1024 has rank at most five.

Proof. Put $a=r_0$, $b=r_1$ and $H=\langle r_1,\dots,r_{(d-1)}\rangle$. IP gives $H\cap\langle a,b\rangle=\langle b\rangle$. If $H(ab)^i=H(ab)^j$, then $(ab)^{i-j}$ lies in the intersection of the rotation subgroup with $\langle b\rangle$, which is trivial. The m_1 such cosets, with $0\le i<m_1$, are therefore distinct. Iterating gives the bound. Each m_i is at least four in the stated two-group case, so $2^{2d-1}\le 1024$, forcing $d\le 5$. ■

The smooth Coxeter quotient assertion in source Theorem 2.7 is the homomorphism from the string Coxeter presentation sending each distinguished generator to its marked involution; the specified adjacent orders make it smooth. A disconnected string diagram yields a direct product of its component subgroups: they commute and IP makes each meet the subgroup on the other components trivially. This gives the source's preliminary reduction; a separate census of every degenerate product is not added to the principal contract. Source Lemma 2.8 is the coordinate-character consequence of Burnside's basis theorem: an irredundant generating list of a finite two-group maps to a basis of $G/\Phi(G)$, and every binary assignment extends to a homomorphism to C_2. No infinite-group assertion is made here.

For rank three write $L=\langle a,b\rangle$ and $R=\langle b,c\rangle$. The retained rank-three criterion says that these two rank-two marked string C-groups, together with $[a,c]=1$ and $L\cap R=\langle b\rangle$, give full IP. Precisely, for d=4 or 5, the versions quoted in source Lemmas 2.3–2.4 require the prefix $H=\langle r_0,\dots,r_{(d-2)}\rangle$ to have IP and each equality $H\cap\langle r_j,\dots,r_{(d-1)}\rangle=\langle r_j,\dots,r_{(d-2)}\rangle$ for $j=1,\dots,d-1$, with the final right side trivial. These are respectively all four and all five source hypotheses. All finite certificates below instead test every subset pair directly.

If $N\lhd G$ and $N\cap RL=1$, U.Q says the quotient facets have exactly their old intersection and both embed; their product orders are unchanged. This is source Lemma 3.1. If $N\le L$ and the displayed images of a,b still give a rank-two marking on L/N, then $N\cap R\le\langle b\rangle$ and $b\notin N$, so R embeds. The last clause of U.Q gives the same facet intersection and the rank-three criterion gives IP. This is source Lemma 3.2, including its symmetric case. An unrelated marking on the abstract quotient L/N would not suffice.

6.2. A safe central quotient and the three complete lift patterns

Assume throughout this subsection that G is a finite marked string two-group $\langle a,b,c\rangle$, with $m=|ab|\ge 4$ and $n=|bc|\ge 4$. Put $x=(ab)^{m/2}$ and $y=(bc)^{n/2}$.

Lemma N2 (central elements in an end product).

$$Z(G)\cap RL\subseteq\{1,x,y,yx,ybx\}. \tag{N1}$$

If yx is central, x and y are separately central. The element ybx is never central.

Proof. Every element of RL has a unique expression $h(ab)^j$ with $h \in R$ and $0 \le j < m$. Equality of two expressions would put a rotation in $L \cap R = \langle b \rangle$. Conjugation by b negates j, so centrality implies j=0 or m/2. The former case gives an element of $Z(R) = \{1, y\}$. In the latter case write z=hx; since x commutes with b, h lies in $C_R(b) = \{1, b, y, yb\}$. The choice h=b fails because the reflection bx does not commute with a when $m \ge 4$. This proves (N1).

If yx commutes with a, then y does, because x already does; commuting with c similarly makes x commute with c. Thus both half-rotations are central. Finally $x, y \in \Phi(G)$, since they are even powers of rotations. If z=ybx were central, then a,z,c would have the same Frattini basis as a,b,c and would generate G. A central z is an involution here: $z \in C_G(x)$ implies y commutes with x, and y,b,x are commuting involutions. Thus a,z,c are three commuting involutions, contradicting $|ab| \ge 4$. ■

Theorem N3 (central descent with IP). G has a central order-two subgroup N such that the displayed images of a,b,c give a rank-three string C-group on G/N. Successive such descents, allowing degenerate intermediate markings, stop at C_2^3.

Proof. If x is central, take $N = \langle x \rangle$. The marked dihedral quotient L/N has rotation order $m/2 \ge 2$ and R embeds because $N \cap R = 1$. The $N \le L$ clause of U.Q proves the quotient intersection. The same reasoning applies to central y. If neither half-rotation is central, choose any central involution z, which exists in every nontrivial finite two-group. Lemma N2 puts z outside RL; now the injective-ends clause of U.Q proves IP and preserves both end groups. A degenerate rank-three marking is C_2 times a dihedral marking. Unless it is C_2^3, the central rotation involution of its nonabelian dihedral factor can be factored out while preserving rank three. Orders strictly decrease by two, so this process terminates exactly at C_2^3. ■

This proves source Corollary 3.3, Lemmas 3.4–3.5 and Theorem 3.6. The bottom boundary is necessary in source Corollary 3.7: a rank-three order-eight group cannot have a rank-three order-four quotient. Its literal unrestricted wording is not asserted.

Theorem N4 (three lift patterns with complete presentations). Let Q be a finite rank-three marked string C-group with a complete presentation in its involutions a_0, b_0, c_0, including Coxeter relations and additional relators $R_1, \ldots, R_t$. Let $G = \langle a, b, c \rangle$ be a central extension with kernel $\langle z \rangle$ of order two, mapping those involutions respectively to a_0, b_0, c_0. Suppose the lifts are involutions, $[a, c] = 1$ and G has IP. Exactly three order patterns are possible: the left adjacent order doubles, the right adjacent order doubles, or neither doubles. In each case a complete presentation is obtained by adjoining central z with $z^2 = 1$, each defining Coxeter word of Q set equal to its actual value 1 or z (in particular $(ab)^p = z$ when its quotient order p doubles), and equations $R_i = z^{\epsilon_i}$, $\epsilon_i \in \{0, 1\}$. At least one relation expresses z as a word in a,b,c, so z can be eliminated.

Proof. Each adjacent order is unchanged or doubled. If both doubled, z would be the halfway rotation in both L and R and hence in their intersection $\langle b \rangle$, impossible for a nontrivial dihedral rotation. Every defining word of Q lifts to 1 or z. The indicated presentation P maps onto G and has $P/\langle z \rangle = Q$; centrality and $z^2 = 1$ imply $|P| \le 2|Q| = |G|$, proving that map an isomorphism. When an adjacent order doubles, its half-power is z. If neither does and every additional relator lifted to 1, the generating tuple would define a surjection $Q \to G$, contradicting the orders. ■

This is source Theorem 3.8 and Table 1. It classifies successful extensions; it does not assert that every binary lift choice has the required order or IP. Those properties and equality of ordered markings must still be checked.

7. Exhaustive bounded classification and named objects

Apply A2.5 (U.Nilpotent/U.Ladder) to the path on d vertices with $k = \log_2|G| - (2d - 1)$. The census program of Appendix C is a single parameterized program for the four cases in the following table. NQ

computes the universal nilpotent quotient of class k+2. The original generators are transported through its epimorphism and pc isomorphism; they are never replaced by an arbitrary generating list. Every normal kernel in the invariant ladder is generated, with only equal subgroups deduplicated. The binary canonical subgroup key is used only after all ambient relative pc orders have been checked to be two.

d	k	Order of U	Order of $\gamma_3(U)$	Kernels	IP	Fixed	Classes	Meets
3	3	2^{18}	2^{13}	61	37	3	20	3904
3	5	2^{38}	2^{33}	401	235	15	125	25664
4	3	2^{34}	2^{27}	1483	199	11	105	379648
5	1	2^{16}	2^{7}	127	15	1	8	130048

Here Kernels counts terminal normal subgroups, IP counts ordered markings satisfying every intersection test, Fixed counts reversal-fixed markings, and Meets counts actual subgroup-intersection checks. The ladder sizes are respectively (7,23,61), (7,23,61,147,401), (31,267,1483) and (127). These are computed outputs, never candidate filters. All 539264 subset-pair intersections are tested, including every candidate rejected by the kernel-profile criterion, and the two IP decisions agree for every candidate. Burnside's orbit formula gives (ordered+fixed)/2 reversal classes. The four ordered type-{4,8} markings in the first case satisfy the four distinct word rows in §5, one each; this is the same finite input used in the eight-family completeness proof. The last three cases prove the classification at order 1024, with the rank bound in Proposition N1 proving that no further nondegenerate rank is omitted.

For rank three the twelve type counts, up to reversal, are:

Type	Count	Type	Count
{4,4}	1	{8,8}	22
{4,8}	8	{8,16}	40
{4,16}	14	{8,32}	8
{4,32}	4	{8,64}	4
{4,64}	2	{16,16}	10
{4,128}	2	{16,32}	10

These are exactly source Table 5. The complete presentations defining all named facets, and the full 107 rows of Table 6, appear as literal mathematical data in the complete defining-data artifact of Appendix C. For every one of the 199 rank-four and 15 rank-five ordered markings, the presentation checker of Appendix C reconstructs the quotient from its exact new kernel key and proves unique generator-preserving isomorphisms from both named end groups: 428 exact facet identifications. It computes dual names from reversed generators, rather than assuming that equal-looking types or letter suffixes are self-dual. Under reversal a facet pair (A,B) becomes (dual(B),dual(A)). Counts are compared after this operation for each of the 99 rank-four and eight rank-five rows individually.

The four printed type-{4,16,4} rows in Table 6 have right-hand type {4,16}; their ordered vertex-figure markings necessarily have type {16,4}. The data use precisely these duals, preserving every letter suffix and count. This qualification was checked on printed page 16; it is not an inferred change to class counts.

For each of the 106 finite rows, the complete universal presentation is the union of the two complete facet presentations, shifted to overlapping generator lists, together with the distant-end commutation relation. Completed Todd–Coxeter enumeration at the trivial subgroup gives 101 orders of 1024 and five orders of 2048. The finite checker also verifies actual IP and both exact facet embeddings. The remaining row is the explicitly retained infinite universal of §7.4; it is never sent to an unbounded finite-order test.

7.2. Every printed rank-three presentation and its corrections

The complete finite-presentation protocol in the presentation checker of Appendix C takes the bare displayed relators, performs complete trivial-subgroup Todd–Coxeter enumeration, and checks that the resulting permutation action has the table's full order. It evaluates every relator, proves generation by the specified marked involutions, and tests all 4^d marked-subgroup intersections. Thus its order bounds concern the raw presented groups, not just finite images or nilpotent truncations.

Write C(p,q) for the rank-three Coxeter relations of type {p,q}. In source Table 2, adjoin to C(4,8) the following pairs, set equal to 1:

Name	R_1	R_2
512a	$(abcb)^4$	$(ca(bc)^3ab)^2$
512b	$(cab)^8$	$(a(bc)^3b)^2$
512c	$(c(abcb)^2)^2$	$(ca(bc)^3ab)^2$
512d	$(a(bc)^3b)^2$	$(cab)^5c(ba)^2(cb)^2ab$

All four raw groups have order 512 and are pairwise inequivalent as ordered markings; their complete primary Atlas presentations are identified exactly. Class I of source Example 3.9 would require an order-512 quotient of C(2,8), whose order is 32, so contributes none. Class II uses C(4,4) with $(abcb)^8=1$, of order 512; its two lifts have central $\tau=(bc)^4$ and $(abcb)^8=1$ or τ. They define the two distinct targets called 1024a and 1024b. For each Table 2 row, the three Class III choices $(R_1,R_2)=(\tau,1),(1,\tau),(\tau,\tau)$, with central $\tau^2=1$, yield exactly:

Base	$(\tau,1)$	$(1,\tau)$	(τ,τ)
512a	1024c	1024d	1024e
512b	1024a	1024c	1024f
512c	1024c	1024g	1024h
512d	512d	1024a	512d

Quotienting by the actual τ subgroup recovers the named base marking in every case. In the two order-512 outcomes $\tau=1$. All eight order-1024 targets are pairwise inequivalent as ordered markings and match the eight type-{4,8} quotients in the unfiltered census bijectively, proving completeness independent of any source table assertion.

Separately evaluating every literal Table 4 row gives the ordered-label map

$$(a,b,c,d,e,f,g,h)_{\text{printed}} \longmapsto (a,b,c,d,c,a,g,c)_{\text{Table 3}}. \tag{N2}$$

To obtain its intended eight targets, the second displayed additional relation in each of printed rows e,f,h must have right-hand side τ instead of 1. These are exactly the already checked (τ,τ) lifts of Table 2 rows a,b,c. No other row or count changes. Printed page 10 was inspected visually; the diagnosis is about the literal source, without claiming priority or unique editorial intent for the repair.

7.3. Universal presentations and all finite examples

For source Definition 4.1 use the combined facet presentation described in §7. The retained geometric universal principle, source Theorem 4.2, states that a nonempty class of regular polytopes with those specified facet and vertex-figure groups has that marked universal string C-group and every member is

its smooth quotient. At the group-presentation level the epimorphism is immediate from the defining words; the abstract-polytope interpretation remains an explicit background input.

The bare presentations in Examples 4.3–4.5 give:

Object	Raw order	Type	Ordered end labels
Example 4.3 universal	512	{4,4,4}	{4,4}*32, {4,4}*128
Example 4.4 universal	1024	{4,4,4}	{4,4}*32, {4,4}*256
Example 4.5 universal	2048	{4,4,8}	{4,4}*64, {4,8}*128a
Example 4.5 first quotient	1024	{4,4,8}	same two labels
Example 4.5 second quotient	1024	{4,4,8}	same two labels
Example 4.5 third quotient	1024	{4,4,4}	not in the prescribed smooth class

The first two universal presentations add $(abcb)^2=(bcdc)^4=1$ and $(abcb)^2=(dbc)^8=1$ respectively to the type-{4,4,4} Coxeter relations. The third adds

$$(cab)^4 = 1, \qquad dbcd(cb)^2(dc)^2bc = 1 \tag{N3}$$

to the type-{4,4,8} relations. The complete named facet words in the complete defining-data artifact of Appendix C differ, where necessary, only by cyclic conjugation from these displayed source words. Their exact generator-preserving isomorphisms and the full raw order/IP tests verify each row above.

For the order-2048 group the three printed central-involution words are

$$bcdcbab(dc)^2dba, \qquad (bcdcba)^2, \qquad (cd)^4. \tag{N4}$$

The checker independently enumerates all central involutions and obtains exactly these three distinct elements. Adding each as a raw relator gives the three quotient rows above. The first two are inequivalent marked groups, with unchanged type and end markings. Every normal order-two subgroup is central, so these three exhaust all order-1024 quotients of this universal, and exactly two belong to the prescribed smooth class. Example 4.3 has no quotient of order 1024; Example 4.4 has only itself. These statements concern these exact orders and do not extrapolate centrality to larger kernels.

7.4. Square-lattice centers and the exceptional infinite row

Retain the finite smooth square-quotient classification quoted in source Lemma 5.1: a finite smooth quotient of [4,4] lies in one of the two families obtained by adding $(abcb)^s$=1, of order $8s^2$, or $(abc)^{2s}$=1, of order $16s^2$, for a positive integer s. The former has |abcb|=s and |abc|=2s; the latter has |abcb|=2s and |abc|=2s. The retained decompositions are $(\langle a,c^b\rangle\times\langle c,a^b\rangle)\rtimes\langle b\rangle$ in the first case, and $(\langle b,b^a\rangle\times\langle b^c,b^{ac}\rangle)\rtimes\langle a,c\rangle$ in the second. Each displayed factor is dihedral of order 2s; the complements have orders two and four respectively. These structural identifications are part of the quoted prior theorem, rather than inferred from the orders. Only the finite scope is used; the infinite Coxeter group itself is not asserted to admit such an s.

Use the common affine model of §3 after reversing its endpoints and conjugating all three generators by b. Since the lattices are invariant under the coordinate swap, the same square and diagonal quotients have coordinates

$$a(x,y) = (1-x,y), \quad b(x,y) = (y,x), \quad c(x,y) = (x,-y). \tag{N5}$$

Its group is W=$\mathbb{Z}^2 \rtimes D_8$; t=abcb translates by (1,0), (abc)² translates by (1,1), (bc)² is (0,−I), and (ab)² is ((1,1),−I). The two finite families are W/s$\mathbb{Z}^2$ and W/L_s, where L_s=⟨(s,s),(s,−s)⟩. The §3 affine normal forms establish these identifications; equivalently the displayed source relations give surjections onto these affine images of the stated orders, hence isomorphisms.

Proposition N5 (all source center formulas). For s=2^m, in W/s$\mathbb{Z}^2$ the center is ⟨(ab)²,(bc)²⟩ when m=1 and $\langle (abc)^s \rangle$ when m≥2. In W/L_s it is ⟨ac,(ab)²,(bc)²⟩ when m=0 and $\langle (abcb)^s \rangle$ when m≥1.

Proof. In a central affine pair (v,M), commutation with all translations forces M to act trivially on T=$\mathbb{Z}^2$/L. If the signed-permutation action on T is faithful, M=I and the center is T^{D_8}.

For L=s$\mathbb{Z}^2$, s≥4, faithfulness is immediate on the two coordinate vectors. Invariance gives 2x=2y=0 and x=y modulo s, leaving precisely 0 and (s/2,s/2). For s=2, the action kernel is the diagonal sign subgroup; to commute also with the linear D_8 subgroup, M must lie in {I,−I} and v in {0,(1,1)}. This gives the first exceptional center. No first-family formula at m=0 is claimed in the source or here.

For L=L_s with s≥2, a nontrivial diagonal sign change cannot act trivially, since (2,0) and (0,2) do not lie in L_s. A signed swap cannot act trivially because its difference on a coordinate vector has a coordinate ±1, not divisible by s. A fixed translation satisfies (2x,0),(0,2y)∈L_s, hence s divides both x,y; modulo L_s its class is 0 or (s,0), both swap-invariant. For s=1 the lattice consists of vectors with even coordinate sum, so T=C_2 with trivial D_8 action and the group is $C_2 \times D_8$. Here ac=(t,−I), (bc)²=(0,−I), and (ab)²=(bc)² in the quotient, giving exactly the claimed generating set. Its three displayed generators are not independent. ■

For the sole infinite row of Table 6 retain the exact source Theorem 5.3, attributed there to McMullen–Schulte: the universal type-{4,4,4} group with extra relations (abcb)⁴=(bcd)⁴=1 has

$$U = (W_1 \times W_2) \rtimes \langle b,d \rangle, \quad \langle b,d \rangle \cong C_2^2, \quad W_1 = \langle c^{bd}, a, c^b \rangle \cong [4,4], \quad W_2 = \langle c, a^b, c^d \rangle = W_1^b. \tag{N6}$$

In particular it is infinite. This theorem, including its marked decomposition, is retained and is not a result of finite coset enumeration.

Proposition N6 (the exceptional kernel location). If π:U→G is a marked epimorphism to a finite string two-group with IP, then ker$\pi \le W_1 \times W_2$ and

$$G \cong ((W_1 \times W_2)/\ker\pi) \rtimes \langle \pi(b), \pi(d) \rangle. \tag{N7}$$

Proof. IP makes the four marked generators irredundant. Their images form a basis of G/Φ(G). The b- and d-coordinate characters, composed with π, agree on every generator with the projection U→C_2^2 in (N6): the displayed W_i generators have even b- and d-parities. Thus every element of kerπ lies in $W_1 \times W_2$. The image of ⟨b,d⟩ still has order four by these same coordinate characters and meets the image of $W_1 \times W_2$ trivially, proving the split quotient formula. ■

This proves source Lemma 5.4 without assuming quotients preserve IP. At order 1024 the first factor in (N7) has order 256. Finally the two source Theorem 5.5 presentations share the type-{4,4,4} Coxeter relators and (abcb)⁴=(bcd)⁴=1. The first adds (dcbabdc)²=1. The second adds

$$(bdcdba)^2(cbab)^2 = 1, \qquad (abcb)^2(babdcd)^2 = 1. \tag{N8}$$

the presentation checker of Appendix C proves directly that these two raw groups have order 1024 and full IP, and identifies them with the two and only two census markings whose ordered ends are {4,4}*128 and {4,4}*64. Unequal end orders rule out their becoming equal under reversal. This proves the exact exceptional classification without repeating the source's separate case analysis or presuming its presentations finite.

The finite execution in the presentation checker of Appendix C totals 214 complete presentations and 43840 actual IP meets: 72 defining Atlas models, 34 source example/literal-table presentations, the two exceptional presentations, and 106 finite universal presentations. Its distinct logical role from the census program of Appendix C is preserved: the universal-kernel proof gives exhaustion, while complete raw coset tables and exact marked isomorphisms give presentation and name recognition. All code and mathematical data are included as complete external appendices with the bindings in Appendix C. These are exact finite computations conditional on the stated algorithms, not Lean proofs or experimental extrapolations. The concluding source suggestion about geometric realizations, a separate classification of all degenerate products, and historical novelty of the corrections are outside the contract.

8. Free extensions and two-primary torsion

For the infinite statements we retain the following previously established **free-extension identifications**, including their geometric existence, universality and symmetry conclusions. All input ranks here satisfy the source's standing convention $d \geq 3$:

1. If K is regular with facet F, its free extension is regular, of rank one higher, with group $\Gamma(K) *_{\Gamma(F)} (\Gamma(F) \times C_2)$.
2. If K is oriented chiral with directly regular facets F, its free extension is oriented chiral, of rank one higher, with group $\Gamma(K) *_{\Gamma^+(F)} \Gamma(F)$, where $\Gamma(F) = \Gamma^+(F) \rtimes C_2$.
3. If regular d-polytopes P, Q have isomorphic facets K, their universal alternating semiregular $(d+1)$-polytope has group $\Gamma(P) *_{\Gamma(K)} \Gamma(Q)$, with an additional branch-swapping C_2 precisely if $P \cong Q$.

The first is the standard regular free-extension construction; the second is Schulte–Weiss, *Free extensions of chiral polytopes* (1995); the third is the universal alternating construction of Monson–Schulte (2012, 2021). The embeddings are the indicated face or facet-rotation subgroups, not an assumed embedding of a full facet group in a chiral automorphism group.

In each of the three free-extension identifications the common subgroup is proper in both factors, as the face/flag action or the new reflection shows. In the second factor of the first two constructions its index is two. Every factor has two-primary torsion: in the chiral case the **rotation subgroup** of the facet embeds in $\Gamma(K)$, and its extension by C_2 suffices to prove the property for $\Gamma(F)$. All factors are finitely generated by their finite marked sets. U.Amalgam and, when needed, its finite-extension clause prove exactly **F016 Theorems 7.1, 7.2 and 7.3**, including infinite input groups. No assertion that all elements have finite order is made.

9. Source contracts, corrections and attribution

The F013 common-quotient theorem is recovered by a different kernel. Its specified auxiliary subgroup is false in general. We include the proof of the correction so that no false historical claim is counted as a corollary of this manuscript.

In the path model (A-F), name the central edge coordinates $c_1, \ldots, c_{d-1}$, and put

$$\sigma_i = r_{i-1}r_i, \quad A_i = \langle \sigma_1, \ldots, \sigma_{i-1}, \sigma_i^2, \sigma_{i+1}, \ldots, \sigma_{d-1} \rangle, \quad C = \bigcap_i \Phi(A_i). \tag{A-X1}$$

These are exactly the source's equations (7)–(9), and $\sigma_i^2 = c_i$. The vertex images of the rotations are independent and span the even-weight subspace. Each A_i contains all central coordinates, with vertex-image dimension $d-2$. Thus $|A_i| = 2^{2d-3}$. Squaring a collected class-two word shows its square subgroup is generated by the squares and pair commutators of the generators; Burnside identifies it with $\Phi(A_i)$. The endpoint computations give

$$\Phi(A_1) = \langle c_2, \dots, c_{d-1} \rangle, \qquad \Phi(A_{d-1}) = \langle c_1, \dots, c_{d-2} \rangle.$$

All listed coordinates are squares; no missing endpoint coordinate can arise because every remaining rotation has the missing endpoint vertex coordinate zero. For an internal i, all coordinates except c_i are squares, and $[\sigma_{i-1}, \sigma_{i+1}] = c_i$. Hence $\Phi(A_i) = \langle c_1, \dots, c_{d-1} \rangle$ internally and

$$C = \langle c_2, \dots, c_{d-2} \rangle, \qquad |C| = 2^{d-3} \quad (d \geq 3). \tag{A-X2}$$

For $d = 2$, separately, $C = 1$. For every $d \geq 4$, this model has order 2^{2d-1} but $C \neq 1 = \gamma_3(G)$. It contradicts F013 v1 Theorem 4.2(c)'s kernel order, (d)'s quotient order and prescribed identification, and (e)'s equality with $\gamma_3(G)$. The actual quotient order is 2^{d+2}; it still has IP, with interior adjacent orders two. Thus this counterexample does not refute the separate main common-quotient existence conclusion, nor every clause of 4.2(d).

Two F016 proof repairs also remain explicit. In a diagonal square-map quotient the perpendicular squared translation subgroups can coincide, so their orders cannot be multiplied as in its page-11 argument; (A-S2) uses the correct lattice index. In its chiral extension proof the full facet automorphism group need not embed in the chiral group; §8 only uses its rotation subgroup and a finite index-two extension.

The exact intake is the five specified editions in the references. The following is the principal-result map; the companion ledger expands every normalized statement, multipart clause and all 33 [GLP18] numbered/table records.

Source	Target contract	Derivation and actual U call
[HFLa], Theorem 1.2	Every admissible rank, order and sequence of 2-power adjacent orders	A4.1; square seed; U.SK with two-generator proper-parabolic checks
[HFLb], Theorem 1.2	Rank-three existence in the stated n≥10 range	d=3 specialization of A4.1
[HFLb], Theorem 1.3	All 2+2+4 high-type families; completeness, bare orders/IP and marking distinction	§5; A3.1; cyclic-core input; A2.5/U.Nilpotent/U.Ladder seed and exact relation checks
[HS26], Theorems 1.1–1.2	Canonical marked quotient and geometric cover for every finite nondegenerate rank	A2.1–A2.4; U.Phi; retained cover correspondence
[CFHS25], Theorem 5.1(a–d)	Exact prescribed sections/order, flatness and all required FAP cuts	A4.2 path binding; U.SK antichain, γ3, separator and deletion clauses
[CFHS25], Theorem 5.3	Exact quotient of the universal local presentation	A4.3 normalization, actual local embeddings and finite order bound
[CFHS25], Theorem 6.1(a–b)	Alternating semiregular assembly and exact full automorphism order	A4.2 fork binding; two-deletion fiber isomorphism; retained assembly theorem
[CFHS25], Theorems 7.1–7.3	Regular, chiral and alternating free extensions, including infinite input	§8; all geometric embeddings/properness checked; U.Amalgam

Source	Target contract	Derivation and actual U call
[GLP18], §§2–3/Table 1	Marked quotient criteria, central descent and all binary lift patterns	§6; U.Q/U.Phi; explicit center-in-product argument
[GLP18], Example 3.9/Tables 2–4	Every stated raw presentation and intended named target; diagnosed literal discrepancies	§7.2; complete coset tables and generator-preserving isomorphisms
[GLP18], Examples 4.3–4.5	Exact universals and their order 1024 smooth quotients	§7.3; complete raw presentation orders/IP and all central involutions
[GLP18], Lemmas 5.1–5.4/Theorem 5.5	Lattice centers, retained infinite decomposition, kernel location and exactly two exceptional targets	§7.4; A3.1 affine model; U.Phi; exact census and presentations
[GLP18], Tables 5–6	All 125 rank 3,105 rank 4 and 8 rank 5 reversal classes; all 107 facet rows and universal orders	A2.5/U.Nilpotent/U.Ladder plus all-parabolic census and exact named-object recognition

The [HS26] auxiliary kernel assertions are corrected above; its two principal existence conclusions remain recovered. The [CFHS25] diagonal-translation and chiral-facet proof steps use the corrected arguments stated here. In [GLP18], Corollary 3.7 stops at order8, Table 4's literal rows remain distinguished from their repaired targets, and reversed facet markings in Table 6 are explicit. The retained inputs are part of the contract, not silently counted as newly proved results. No open question, historical novelty claim, unnumbered small-n classification or separate census of degenerate products is included. The power-polytope recollections in [CFHS25] are not extra source targets.

This scientific organization and the supported-kernel applications are repository work, with the support-inclusion simplification credited to independent reviewer 122. The source authors receive credit for the original results and constructions. The classical ingredients of U and the new assembly carry no historical-priority claim. Removing one declared U dependency only invalidates the recorded derivation; it does not show that the conclusion has no other proof.

Appendix C. Complete computational proof interface

There are four complete external artifacts, all part of this volume's proof cost: `phase78_n009_integration_census.g`, `phase78_n009_integration_finite.g`, `phase78_n009_integration_data.g` and `phase78_n009_integration_run.py`. Their legacy filenames identify established reusable artifacts; no surrounding manuscript is imported by those filenames. The first three contain every executed mathematical instruction and all 72 named Atlas presentations,107 Table 6 rows and source relators. The fourth is the complete no-clobber orchestration and output checker. Source editions and artifacts are hash-bound in the companion ledger before independent review.

The exact algorithms assumed are GAP's finite permutation/pc group arithmetic, subgroup intersection and complete Todd–Coxeter enumeration, and NQ's universal nilpotent quotient algorithm. A completed coset table at the trivial subgroup establishes the raw presentation order; a finite image alone would not suffice. No source count is used to prune candidate generation, and no saved certificate, private primary file, permutation-array atlas or SmallGroups catalogue is read.

For every (d,k), the census constructs $U_{T,k}$ from its original marked nilpotent quotient map, transports those same generators, and enumerates every U-normal index-two hyperplane preimage k times. Only equal subgroups are deduplicated. Its binary subgroup key is used only after checking that every ambient relative pc order is two. A2.5 supplies completeness. Every terminal candidate is subjected both to the

kernel-profile test and to all 4^d actual marked subgroup meets, including rejected candidates. Reversal is computed on the markings; the observed fixed count and the two-element orbit formula give the generator-set classes. The order 256 seed evaluates the four distinct word patterns on each oriented type-{4,8} marking, so its use in §5 is an explicit finite implication with no hidden classification assumption.

The presentation checker constructs all raw groups from complete relators, finishes the trivial-subgroup coset table, evaluates all defining relations, checks generation, exact involution/product orders and all subset intersections. It verifies each facet name by an actual generator-preserving isomorphism. Dual names are computed using reversed generators. It enumerates all central involutions in the order 2048 example and checks every quotient. For each finite Table 6 row it combines both complete facet presentations and distant-end commutation, proves the raw universal order and verifies both facet embeddings. The one infinite row invokes the precisely stated retained decomposition; it is never sent to an unbounded finite-order computation.

The complete run requires 539264 census meets,214 raw presentations and 43840 additional meets,428 exact facet identifications and all 107 Table 6 rows. These are expected results checked after exhaustive generation, not assumptions proving it. Successful replay establishes the finite conditional obligations, not a Lean formalization or a substitute for the parameter proofs.

From the research directory's parent, run:

```
python3 -B -O research/leemans/phase78_n009_integration_run.py --output-dir
/private/tmp/NEW-two-group-certificate
```

The output path must not exist. The artifact manifest records source/program hashes, installed versions, complete logs and all generated mathematical data. The application ledger charges the four source files in full, plus the actual U dependency closure and all written hypotheses in A. Rendered mathematical pages, certificate bytes and any cumulative multi-volume accounting will be reported separately after complete review; the supported-kernel acquisition cost is not waived because it has several users.

References and source-version boundary

[HFLa] D.-D. Hou, Y.-Q. Feng and D. Leemans, *On regular polytopes of 2-power order*, revised author PDF dated 8 November 2019, journal 2020, DOI 10.1007/s00454-019-00119-5. Exact retained primary file: `sources/catalogue_L032.pdf`.

[HFLb] D.-D. Hou, Y.-Q. Feng and D. Leemans, *Existence of regular 3-polytopes of order* 2^n, arXiv:1802.08947v2, 22 January 2019; journal catalogue title *2-groups behaving as automorphism groups of regular 3-polytopes*. Exact retained primary file: `sources/catalogue_L035.pdf`.

[HS26] D.-D. Hou and E. Schulte, *String C-groups of 2-power order project onto a common string C-group*, arXiv:2605.20691v1, submitted 20 May 2026, PDF dated 21 May. Exact retained primary file: `sources/catalogue_F013.pdf`.

[CFHS25] G. Cunningham, Y.-Q. Feng, D.-D. Hou and E. Schulte, *Symmetric polytopes whose automorphism groups are 2-groups*, arXiv:2512.15511v1, submitted 17 December 2025, PDF dated 18 December. Exact retained primary file: `sources/symmetric_2groups2025.pdf`.

[GLP18] Y. Gomi, M. L. Loyola and M. L. A. N. De Las Peñas, *String C-groups of order 1024*, Contributions to Discrete Mathematics **13**(1) (2018), 1–22, DOI 10.55016/ojs/cdm.v13i1.62475. Primary journal PDF, SHA256 `0618a57cb5ff35c13587877818a2e77bce442cf2c561f204b6d5e5e095fa2fda`.

Its quoted standard intersection criteria are attributed to M. Conder and D. Oliveros, *The intersection condition for regular polytopes*, JCTA **120** (2013), 1291–1304. The universal and finite square-quotient inputs are quoted from McMullen–Schulte, *Abstract Regular Polytopes* (2002), the latter on pp. 364–365, 368. Its exceptional infinite decomposition is quoted from McMullen–Schulte, *Regular polytopes from twisted Coxeter groups and unitary reflexion groups*, Advances in Mathematics **82** (1990), 35–87. Those

earlier proofs are retained in the precise quoted forms; no claim of fresh acquisition or rederivation is made.

The named-object data are credited to M. I. Hartley's Atlas of Small Regular Polytopes; the complete defining-data artifact records the exact primary URL and snapshot hash for every defining presentation. The URL may contain a catalogue identifier, but replay neither calls SmallGroups nor assumes a catalogue is complete. The entire new finite search and recognition were run with GAP 4.15.1 and NQ 2.5.11.

The retained cyclic-core result is Lucchini's theorem in the form recorded by Isaacs, *Finite Group Theory*, Theorem 2.20. The retained alternating assembly is Monson–Schulte, *The assembly problem for alternating semiregular polytopes* (2020), Theorem 5.3 in its FAP case. The chiral free-extension theorem is Schulte–Weiss, *Free extensions of chiral polytopes* (1995); the universal alternating construction is Monson–Schulte (2012,2021), in the exact forms stated in §8. These prior proofs are retained rather than included as additional compressed sources. The regular-polytope and marked-cover correspondences are the standard McMullen–Schulte framework, with the scopes stated in §1.