The third Janko row in the Higman–Sims comparison table

A source join of the acquired finite incidence certificates

Put $G = J_3$ and $N = 50\,232\,960$. The source of the comparison is Table 1, pages 232–233, of D. Leemans, *The residually weakly primitive geometries of HS*, Australas. J. Combin. **33** (2005), 231–236 (L091). Its Janko classification is expressly attributed to reference [9], the earlier Janko paper (L097). We distinguish the finite results below from that earlier classification's coverage premise.

Proposition 1. *The seven acquired rank-two prescriptions are eligible and mutually nonisomorphic, even allowing a permutation of the types. Exactly two are truncations of a supplied rank-three geometry:* 2.3 *is the truncation of active row* 2 *on types* $1, 3$*, and* 2.6 *is the truncation of active row* 1 *on those types. The other five are not truncations of any of the twenty-one supplied rank-three geometries.*

Proof. Eligibility of $2.1, 2.2, 2.4, 2.5, 2.7$ is already established by the accepted literal realizations. For 2.3 and 2.6 we use the actual subgroups P_1, P_3 of their accepted parent. The incidence maps are the identity on the retained right-coset sets, so preserve both incidence and nonincidence. Their intersections and indices are

	$\lvert P_1\rvert$	$\lvert P_3\rvert$	$\lvert P_1 \cap P_3\rvert$	$[P_1 : P_1 \cap P_3]$	$[P_3 : P_1 \cap P_3]$
2.3	8160	2880	480	17	6
2.6	1080	2880	180	6	16

The complete actual local coset actions are doubly transitive, hence primitive, and the two subgroups generate the accepted ambient group. Thus these geometries are connected and firm; all rank-one residues are nonempty. Rank-two coset incidence is flag-transitive because an incident pair has a common representative. For weak primitivity at the empty flag, retain the parent's already checked primitive type: type 1 in active row 2, and type 3 in active row 1. These are the same actual subgroups and ambient actions, not merely subgroups of the same order. Together with the primitive local actions this proves residual weak primitivity.

For completeness, the rank-two intersection test uses all double cosets. Put $B = P_1 \cap P_3$. The neighbours of P_1 are the cosets P_3h, where h runs over a right transversal of B in P_1. Two distinct objects of the first type with a common object of the second type can be transported to P_1, P_1x with $x \in P_3 \setminus B$. The orbits of such pairs are indexed by the nonidentity (B, B) double cosets in P_3. For every representative the two complete neighbour sets intersect in exactly one object, as checked in the actual finite records. Hence both geometries satisfy IP_2. No implication from local double transitivity to this intersection assertion is assumed.

The seven unordered pairs of object cardinalities are

2.1	17442	43605	2.2	17442	17442
2.3	6156	17442	2.4	26163	43605
2.5	23256	25840	2.6	17442	46512
2.7	14688	14688			

They are distinct, proving bare nonisomorphism. For a flag-transitive rank-two coset geometry, its two object counts and its number of incident pairs are $N/|P|$, $N/|Q|$, and $N/|P \cap Q|$. These are intrinsic even if an isomorphism does not preserve the specified action. We compare the two parabolic orders and their full intersection order for each of the three pairs in each of the twenty-one supplied rank-three families. Among all sixty-three comparisons, only active row 1, types $1, 3$, matches 2.6, and only active row 2, types $1, 3$, matches 2.3. All other candidate truncations are excluded by the intrinsic counts. The two matches are actual isomorphisms by the earlier literal coset maps, completing the assertion. □

The comparison row and its retained premises

The accepted named table of marks gives 137 conjugacy classes of subgroups and 71 564 248 subgroups of G. The supplied rank-three roster has nineteen bare incidence classes by the separate accepted flag-count and point-graph argument. Consequently the finite data give seven distinct rank-two geometries, two explicit truncations and nineteen distinct rank-three geometries. To promote these to the global nontruncation row

$$(\text{rank } 2, 3, 4, 5, 6) = (5, 19, 0, 0, 0),$$

we retain exactly the following inputs from L091 reference [9]: every eligible geometry appears in the supplied earlier classification, and there is no eligible geometry of rank at least four. Under these inputs, every possible rank-two truncation is among the sixty-three tested pairs, and no rank-three geometry can be a truncation of a higher rank one. The printed rank-three value seventeen is replaced by nineteen; the seventeen diagram labels of the supplement do not count its bare classes. The correction is explicit, while an independent global Janko census is not claimed.

The reference [9] original is outside the active 105 mathematical manuscripts, and its author supplement is an auxiliary source. No L097 recovery credit is awarded. This join does not import every metric, abstract subgroup name or automorphism assertion printed in that separate supplement. The new HS classification and the Hoffman–Singleton constructions of L091 have their own accepted proofs. Their integration, correction ledger and distinct whole-source review remain the final L091 admission steps. GAP and the complete table of marks retain their explicit software and classification premises. No new conjecture, Lean or compression result is claimed by this join.