Marked geometry of strict Conway compositions

Abstract

An ordered expression in crossing-circle tangles, products and physical turns determines a link diagram and a real wire polytope. We construct the complete hyperbolic metric whenever that polytope has a strict state. Four separate boundary-edge labels allow the pyramid-and-prism construction to pass through turned elementary children and arbitrary strict closing pairs. We prove convexity, retain every original shaded triangle, and identify the closed assembly with the literal projection cut of the specified link. Whole complex face maps then give its marked deformation, with parabolic main cusps. The original primitive filling equations have a unique small solution for independently long signed slopes. Smooth completion and an absolute fundamental-chain comparison identify the volume with a critical value of a classical cell potential. No asymptotic formula for a quantum invariant is used or asserted.

1 The fixed diagram and the geometric statements

A crossing circle is an auxiliary unknotted component bounding a disk punctured twice by the main strands. We retain its disk and its original framing. Let h be the physical quarter-turn of this elementary two-strand tangle, with full twists removed. A construction is a finite expression in distinct copies of h, ordered products and physical clockwise quarter-turns T. No reduced algebraic expression replaces the original diagram in this paper.

The four endpoints of a box are numbered $0, 1, 2, 3$ cyclically, starting at the bottom left. In the stack FG, with F below G, join F_2 to G_1 and F_3 to G_0. Its exposed tuple is (F_0, F_1, G_2, G_3). A turn replaces a tuple by (C_1, C_2, C_3, C_0) and moves the entire box and all its disks. Cylindrical closure C joins endpoints 0 to 3 and 1 to 2. Our closed diagrams are $C(FG)$ with arbitrary such F, G subject to the following condition.

Definition 1.1. A strict wire state assigns a leaf variable $a \in (0, 1/2)$ and an output variable to every open subexpression. A turn complements its variable. Every product retained below the closing root satisfies

$$\max(x, y) < z < x + y < 1. \tag{1.1}$$

The two closing outputs are $b, 1 - b$. The final forced gate is removed, but this complementary relation is retained. An expression is strict if the resulting real polytope has a strict state.

Thus strictness is a condition on this particular presentation. A turned leaf or a bare leaf may occur at a closing child. This is larger syntax than the completed/completed closing grammar of [2]; its global metric theorem will not be applied here.

Fix $\varepsilon_i \in \{-1, 0, 1\}$ at the H original leaves. It specifies the retained signed half twist; $\varepsilon_i = 0$ is the pure disk pairing. Let Y_E^{ε} be the exterior in S^3 of the resulting augmented link, retaining all H crossing circles and every main component. The source-independent functions

used below are

$$J(u) = \int_0^u \log(2\sin \pi t)\,\mathrm{d}t \quad (0 \le u \le 1), \qquad J(1-u) = -J(u), \tag{1.2}$$

$$g(a) = J(2a) - 4J(a), \tag{1.3}$$

$$\begin{aligned} m(x,y,z) = {} & J(x+y) + J(z) - J(x) - J(y) \\ & - J(z-x) - J(z-y) - J(x+y-z). \end{aligned} \tag{1.4}$$

Let $\mathcal{F}$ be the sum of g over leaves and m over retained products, and let F_0 denote its maximum. The variational theorem of [1] supplies the unique strict optimizer and its negative definite Hessian in independent wires. We state the precise input in Section 2; its proof does not use a global metric or a quantum noncancellation theorem.

Theorem 1.2 (Complete marked exterior)**.** *For every fixed strict expression and retained half-twist pattern, Y_E^ε has a complete finite-volume hyperbolic metric assembled from two copies of a convex right-angled ideal polyhedron. Its volume is*

$$\mathrm{Vol}(Y_E^\varepsilon) = 2\pi F_0. \tag{1.5}$$

At each crossing circle the original meridian μ_i and disk longitude λ_i are a primitive ordered basis. Their complete longitude-to-meridian translation ratio is

$$\tau_i = \frac{2\mathrm{i}c_i}{1 - \varepsilon_i \mathrm{i}c_i}, \qquad c_i = \cot(\pi a_i) > 0, \qquad \operatorname{Im}\tau_i > 0. \tag{1.6}$$

All original main components have complete cusp ends.

The primitive basis fixes the filling convention. For a subset $I \subseteq \{1,\dots,H\}$ and integers p_i, fill precisely the auxiliary slopes $\mu_i + p_i\lambda_i$, $i \in I$. Denote the original filled exterior by $Y_E^\varepsilon(\mathbf{p}; I)$. Cutting the original disk, rotating through p_i full turns and removing its crossing circle inserts the physical power $\varepsilon_i - 2p_i$ in that same disk frame. The disjoint disk neighborhoods show directly that the ambient manifold is still S^3. We never fill a main cusp.

Theorem 1.3 (Primitive signed fillings)**.** *There is a holomorphic classical potential $\Phi_\varepsilon(\boldsymbol{\chi})$, with $\Phi_\varepsilon(0) = F_0$, and a framed representation of the original group near its oriented complete holonomy, whose primitive similarity logarithms are*

$$U_i = -2\pi\mathrm{i}\chi_i, \qquad V_i = -2\partial_i\Phi_\varepsilon. \tag{1.7}$$

Every main peripheral group remains parabolic. Set $\chi_i = 0$ outside I. There is a bound $B_{E,\varepsilon,I}$ such that, whenever all $|p_i| > B_{E,\varepsilon,I}$, the restricted action

$$S_{\varepsilon,\mathbf{p}}(\boldsymbol{\chi}) = \Phi_\varepsilon(\boldsymbol{\chi}) + \frac{\pi\mathrm{i}}{2}\sum_{i\in I}\frac{(\chi_i - 1)^2}{p_i} \tag{1.8}$$

has a unique critical point in a fixed neighborhood of zero. The representation there completes to a smooth complete finite-volume hyperbolic metric on the original $Y_E^\varepsilon(\mathbf{p}; I)$, and

$$\mathrm{Vol}(Y_E^\varepsilon(\mathbf{p}; I)) = 2\pi \operatorname{Re} S_{\varepsilon,\mathbf{p}}(\boldsymbol{\chi}_*). \tag{1.9}$$

The signs and relative sizes of the filling integers are independent. If I contains all auxiliaries and the literal strand pairing has one component, the right side is the volume of that knot complement.

The uniqueness assertion is local in this prescribed family, not a classification of the character variety. The potential is defined from the cells in Section 6; no selected quantum action is an input. A maximum over the finitely many ε, I gives one bound for a fixed strict expression if desired. An empty filling subset recovers Theorem 1.2.

The classical geometry of fully augmented links and their half-twist partners is well established; see [5]. Dehn filling and holomorphic peripheral potentials belong to the framework of [6]. The work here is the ordered individual-edge assembly and its marked continuation for strict expressions, with the original manifold and the absolute normalization proved explicitly.

2 Variational input and local cells

We use the following specific conclusions of the strict-profile companion [1, Lemma 2.2, Theorem 3.3, Lemma 4.1 and Proposition 4.2]. For each nonempty strict fixed-output domain, the maximum over the internal wires is strict, unique and nondegenerate. Its profile $f_E(b)$ is real analytic on that domain. The analogous closed statement holds on Definition 1.1's reduced affine constraint space. The exact open types and product table are

$$L = (0, 1/2), \quad H = (1/2, 1), \quad A = (0, 1), \qquad \begin{array}{c|ccc} & L & H & A \\ \hline L & A & H & A \\ H & H & \varnothing & H \\ A & A & H & A \end{array}. \tag{2.1}$$

Every point of each indicated interval is attained. A leaf has type L; a turn exchanges L, H and preserves A. The nonempty closing pairs are AA, AL, AH, LH and their reversals.

For an optimized output define

$$B = \cot \pi b, \quad \varkappa = e^{-f_E'(b)}, \qquad X^2 = \frac{1 + B^2}{1 + \varkappa}, \quad Y^2 = \varkappa X^2, \quad X, Y > 0. \tag{2.2}$$

Then $X^2 + Y^2 - B^2 = 1$, and the imported profile identities give

$$L : (B, X, Y) = (B, B, 1), \qquad H : (B, X, Y) = (-Y, 1, Y), \qquad A : -Y < B < X. \tag{2.3}$$

A turn sends (B, X, Y) to $(-B, Y, X)$. At a stationary product the two children have a common X, and, writing $S = B_1 + B_2 > 0$,

$$B_o = \frac{B_1 B_2 + X^2 - 1}{S}, \qquad Y_o = \frac{Y_1 Y_2}{S}. \tag{2.4}$$

These are scalar variational identities; they do not identify an open quadrilateral with a physical four-punctured tangle boundary.

The local geometric inputs are the explicit cells of [2, Section 2]. They require only the strict inequalities, not the completed grammar. A leaf a is the convex ideal pyramid

$$O = \infty, \qquad (C_0, C_1, C_2, C_3) = (1, E, -1, -E), \qquad E = e^{2\pi \mathrm{i} a}. \tag{2.5}$$

Its base is an ordered quadrilateral gluing face. The side triangles (O, C_1, C_2) and (O, C_3, C_0) are shaded; the other two are white. The four apex edges are right angled. On the base, edges $01, 23$ have angle πa and edges $12, 30$ have angle $\pi(1/2 - a)$. The pyramid volume is $\pi g(a)$.

For a product put $p = z - x$, $q = z - y$, $r = x + y - z$, $w = 1 - x - y$. All are positive. The prism has ideal vertices

$$v_0 = 1, \qquad v_1 = e^{2\pi \mathrm{i} p}, \quad v_2 = e^{2\pi \mathrm{i}(p+r)}, \quad v_3 = e^{2\pi \mathrm{i}(p+r+q)}, \tag{2.6}$$
$$P = v_1 + \frac{\sin \pi x \sin \pi r}{\sin \pi w \sin \pi p}(v_1 - v_0),$$

and ∞. Its two triangular faces are white. Its ordered attachment and output tuples are

$$F : C \mapsto (v_0, v_1, P, \infty), \quad G : C \mapsto (\infty, P, v_2, v_3), \quad C^o = (v_0, v_1, v_2, v_3). \tag{2.7}$$

The angle table from the explicit supporting planes is

edge	$Pv_1, \infty v_0$	$Pv_2, \infty v_3$	v_1v_2, v_0v_3	$P\infty$	$v_0v_1,\ v_2v_3$
angle/π	x	y	$1-z$	w	$p,\ q$

. (2.8)

The prism volume is $\pi m(x, y, z)$. These volume normalizations follow by subdivision into positive ideal tetrahedra and the Lobachevsky formula, [2, equation (2.1)].

For an ordered quadrilateral let

$$\varkappa(a, b, c, d) = \frac{(a-b)(c-d)}{(a-d)(b-c)}.$$

It determines the whole ideal face, including shear. The local calculation [2, equations (3.2)–(3.3)] gives input cross ratios e^{-m_x}, e^{-m_y} and output e^{-m_z} in the canonical prism order. The first two assignments in (2.7) invert the canonical input ratios. Consequently $f'_F(x) + m_x = 0$ and $f'_G(y) + m_y = 0$ match all four ordered vertices. Full reversal $C^F_j \mapsto C^G_{3-j}$ preserves $\varkappa$, so the closing equation $f'_F(b) = f'_G(1-b)$ matches the entire closing face as well. No comparison of absolute values replaces these isometries.

3 Four boundary labels and convex assembly

An open cell has one ordered quadrilateral gluing face $Q = (C_0, C_1, C_2, C_3)$. Number its edges $j = (j, j+1)$ modulo four. Edges $0, 2$ are called free, and $1, 3$ cap edges. These names record the original leaf order, not an unmarked symmetry of the whole tangle. Set $\eta_j = 1$ if the face beside edge j, other than Q, is shaded, and set $\eta_j = 0$ if it is white. The four bits $\boldsymbol{\eta} = (\eta_0, \eta_1, \eta_2, \eta_3)$ will be called the boundary mask.

Proposition 3.1. *At every open subexpression evaluated at its strict stationary wires there is a convex ideal cell with a maximal, nondegenerate ordered quadrilateral output. Its boundary-edge angles are*

$$\frac{\theta_j}{\pi} = \begin{cases} b - \eta_j/2, & j = 0, 2, \\ 1 - b - \eta_j/2, & j = 1, 3. \end{cases} \tag{3.1}$$

Every other surviving edge is right angled and separates a white face from a shaded triangle. The possible supports satisfy

$$L : \eta_0 = \eta_2 = 0, \qquad H : \eta_1 = \eta_3 = 0, \qquad A : \boldsymbol{\eta} = 0. \tag{3.2}$$

Every original shaded triangle remains an entire face. The four output corners are four active original terminal singletons. Every other main ideal vertex is a permanently consumed pair of terminals; crossing-circle apices stay separate.

The statement permits a type H output with only one shaded free edge. Replacing its two free angles by a common angle would therefore lose information essential to the next gluing.




Figure 1: Individual boundary labels in an HL product. A colored edge has a shaded adjacent face; the other edges have a white adjacent face. The turned leaf has mask $(1, 0, 1, 0)$, the upper leaf has mask $(0, 1, 0, 1)$, and the product has mask $(1, 0, 0, 0)$. Its two free angles are $\pi(z - 1/2)$ and πz. The pictures display the ordered quadrilateral gluing faces.

3.1 Every seam angle

For a leaf the mask is $(0, 1, 0, 1)$, so (3.1) is the pyramid angle calculation. A turn rotates the mask by one and replaces b by $1 - b$. This proves the assertion under turns and exchanges the support conditions for L, H.

At a product the output mask is exactly

$$\boldsymbol{\eta}^o = (\eta_0^F, 0, \eta_2^G, 0). \tag{3.3}$$

Indeed the two output cap sides adjoin the prism's white triangles, and each output free side inherits its indicated child side. A child cap seam has angle $\pi(1 - \eta_j/2)$: a zero bit merges two white pieces flatly, while a one bit leaves a right shaded–white edge. At the edge common to both prism inputs, the total angle is

$$\pi \left[w + x - \frac{\eta_2^F}{2} + y - \frac{\eta_0^G}{2} \right] = \pi \left[1 - \frac{\eta_2^F + \eta_0^G}{2} \right]. \tag{3.4}$$

Both bits could be one only with two type H inputs, forbidden by (2.1). Thus this seam is either flat white–white or right shaded–white. The remaining free output angles are $\pi(p + x - \eta_0^F/2)$ and $\pi(q + y - \eta_2^G/2)$; the cap angles are $\pi(1 - z)$. This gives (3.1).

The support assertion also follows case by case from (2.1). An output of type A has no type H child, so both inherited free bits vanish. An output of type H has zero cap bits by (3.3). Turns give all type L cases. We assert these support inclusions, not equality with the two-bit leaf masks.

For successive convex attachments one must check the intermediate stage, not only the final totals. Attach F to the prism first. Its free seams have angles

$$\pi(z - \eta_0^F/2), \qquad \pi(1 - y - \eta_2^F/2).$$

If the first bit is one, F has type H, hence $z > x > 1/2$. If the second bit is one, $x > 1/2$ and $x + y < 1$ give $y < 1/2$. Both displayed angles are therefore positive and strictly less than π. In particular the other prism input remains a supported nondegenerate quadrilateral. The cap seams have already been checked. The second attachment is now justified by (3.4).

3.2 Convexity includes the ideal boundary

Here is the geometric argument used at each attachment. Place two convex finite-faced ideal polyhedra on opposite sides of the plane of the matched quadrilateral. Their finite

intersection is that face. The finite union is closed, connected and locally convex: the only new boundary points lie on the seam edges, where the total angle is at most π. There are no finite seam vertices.

Two finite points in the union have a shortest path in it. Indeed a length-bounded minimizing sequence stays in a compact ambient hyperbolic ball. Constant-speed parametrization and Arzelà–Ascoli give a limiting path; closedness keeps it in the union. Local convexity makes each small subarc the ambient geodesic. The complete path is therefore the unique ambient geodesic segment. This proves convexity of the union.

There is a further ideal-point issue. The two ideal closures can meet only on the boundary circle of the common supporting plane. The intersection of either cell with that plane is exactly its maximal quadrilateral face. An additional ideal vertex on the circle would enlarge that face past one of its edges, making the edge flat; all four supporting angles there are strictly nonflat. Equivalently, in the Klein model the closed quadrilateral meets the sphere at infinity only in its four vertices. Thus the only ideal coincidences are the four specified corner identifications. This argument applies again to the second input, which the intermediate angle check kept strictly supported.

Every output angle in (3.1) lies in $(0, \pi)$ on its actual type interval. Hence the output remains a maximal supported quadrilateral and the induction continues. Every shaded-triangle seam is right, never flat. Only white faces merge: no shaded edge or triangle is deleted, split or merged with another shaded face. A leaf apex is never in an attaching quadrilateral. It retains its four right edges and its two original shaded sectors.

3.3 Terminal pairs never become chains

Give leaf i the four distinct terminal symbols $(i, 0), \dots, (i, 3)$ and the distinct apex symbol O_i. The purely literal endpoint rules prove the following invariant independently of the metric. An open expression has four active singleton terminals; all other terminals form disjoint consumed two-element pairs. A turn only permutes the active tuple. Product consumes exactly

$$F_2 \sim G_1, \qquad F_3 \sim G_0, \tag{3.5}$$

and keeps F_0, F_1, G_2, G_3 active. The child leaf sets are disjoint. A consumed pair is never subsequently attached or exposed. Thus no class grows to three terminals. The geometric maps (2.7) make exactly these pairs; the ideal-point argument excludes any additional identification.

At an exposed corner the two adjacent mask bits have sum at most one. If their sum is zero, its cyclic faces are the gluing quadrilateral, white, shaded, white; if the sum is one there are only the quadrilateral, white and shaded faces in the retained order. This also follows by taking the ideal Euclidean link: the two quadrilateral-edge angles sum to π or $\pi/2$, and the remaining edge angles are right. Each exposed corner therefore carries exactly one original shaded sector. At (3.5), the two sectors concatenate in the original ribbon order. This proves every assertion of Proposition 3.1.

4 The closed polyhedron and the original link

At the closing root glue the entire face by

$$C_j^F \longmapsto C_{3-j}^G. \tag{4.1}$$

The profile equation already matches its ordered metric. An edge j on the first child matches edge $2-j$ on the second. Its total angle is π minus half the sum of the two mask bits, in units

of π. Two shaded free bits would require an HH closing pair; two shaded cap bits would require LL. Neither is strict. The convex-union proof therefore applies, and all remaining edges of the closed polyhedron P_E are right shaded–white edges.

The four closing pairs in (4.1) are between distinct root-child leaf sets. Together with (3.5) they pair every original terminal once, never with a terminal of the same leaf. Hence each leaf still has four distinct corner vertices. Its two shaded triangles are distinct and meet only at their original apex. This rules out a transitive corner collapse at closure.

The boundary of the actual convex polyhedron is a sphere. Its white faces are maximal convex ideal polygons. They cannot be bigons or zero-width regions. No two shaded triangles share an edge: every surviving edge has a white side. These observations will matter for both the topological cut and the cusp returns.

4.1 Recovering every colored face from the literal wires

Compile the same boundary without prisms. At leaf i put the two shaded triangles

$$(O_i, C_{i1}, C_{i2}), \qquad (O_i, C_{i3}, C_{i0})$$

in its original oriented projection chart. At a terminal record its outgoing-next and incoming-previous shaded neighbors. Joining two actual identity-wire endpoints concatenates those two ordered sectors. To see the order directly, take a thin planar neighborhood of the joining arc: its two white sides join the complementary sectors, with no crossing or reflected child. Turns rotate the charts; the final closure uses (4.1). This constructs the full oriented ribbon graph, not merely its abstract incidence or component count.

All white face cycles are now determined. A facial walk entering along (u, v) leaves v along the neighbor preceding u in its cyclic order. The original shaded triples are some of these walks; the remaining walks are exactly the white regions. The prism maps (2.7) induce the same two pairings and the same sector concatenation. The mask proof shows that every other seam is flat white–white and every shaded edge remains. Thus the metric boundary has precisely this ribbon graph and every one of its colored facial cycles.

4.2 The reversible disk cut

Each original auxiliary circle lies in its own leaf ball, perpendicular to the projection sphere, with its recorded twice-punctured disk. The disks and their collars remain disjoint under physical turns and products. In either projection ball each auxiliary semicircle is boundary parallel by the corresponding half disk. Drilling the semicircle introduces one handle; cutting along that same bridge half disk cancels the handle in its explicit collar. The disjoint cancellations leave a ball. The main segments lie on the projection boundary; removing their neighborhoods creates ideal ends and no interior handle.

The cut ball has one ideal vertex for each paired main segment and one for each auxiliary semicircle. The exposed half disks give exactly the two shaded triangles above. The remaining regions have precisely the compiled white facial walks. Identify this labeled boundary sphere with ∂P_E and extend over the balls by radial extension after choosing interior points. Across the two balls restore the identically labeled white projection faces. For a pure leaf restore the shaded pairing fixing O_i and sending C_{i1} to C_{i0} and C_{i2} to C_{i3}. For an odd leaf use the original half-disk rotation, which pairs the corresponding triangles across copies. Reversing these cuts gives exactly Y_E^{ε}.

This is a degree-one topological identification with all components and paths retained. It does not interpret an intermediate gluing quadrilateral as the physical four-punctured boundary of a tangle: that assertion is unnecessary and generally false.

4.3 Complete returns and a common horosphere decoration

Take two oppositely oriented copies of P_E. For a pure leaf, an edge e of one shaded triangle has a distinct partner e' on the other. Its four occurrences are

$$e^{(0)},\ e'^{(0)},\ e'^{(1)},\ e^{(1)}.$$

The labeled shaded maps are inverse maps and the white maps identify equal labels across copies, so the return is pointwise identity on the edge. The angle sum is $4(\pi/2) = 2\pi$. An odd pairing changes the copy permutation in this return. It still has four distinct occurrences, since the two shaded triangles have no common edge and the two cut copies are distinct. The local pure and odd return identities of [2, equations (5.7)–(5.8)], written again in (8.2) below, show that the full return is identity. Thus the quotient has neither an edge cone angle nor a residual translation along an edge.

At an ideal vertex all edges are right. Its small horosphere section is a finite Euclidean polygon with all angles $\pi/2$, hence a rectangle. Its side lengths are positive: the supporting planes are distinct, and coincident opposite sides would force the three-dimensional polyhedron into a plane. The colors alternate. Scale its horosphere until its two equal shaded side lengths are one. Each shaded ideal triangle now has its canonical three unit horocyclic segments, so every shaded isometry preserves these decorations, including the odd permutations. White maps preserve them because they identify the identical labeled copies. The rectangles therefore glue without dilation into smooth compact Euclidean tori, the original cusp tori of the cut. A common shrinking factor makes the horospheres disjoint; beyond them are complete horoball ends. The truncated core is compact. The resulting finite-volume metric is complete.

All cell attachments had disjoint interiors. There is one leaf pyramid and one prism per retained gate in each cut copy. Their volumes are πg and πm, respectively. Consequently $\mathrm{Vol}(Y_E^\varepsilon) = 2\,\mathrm{Vol}(P_E) = 2\pi F_0$ as an equality of ordinary metric volumes. This proves the metric part of Theorem 1.2, including the two-leaf strict case.

5 The primitive collar and its component information

The auxiliary apex retains its original leaf rectangle throughout the construction. In (2.5) its adjacent sides have lengths $2\sin\pi a$ and $2\cos\pi a$. The cusp has exactly two such rectangles. The original disk paths in [2, Section 7, equations (7.1)–(7.5)] are local to this unchanged collar. We recall the path and area checks so that no global completed-root marking is being imported.

Normalize the ordered leaf base to $(0, \infty, 1, R)$. Let W_j be the white transition at edge j between projection charts, and let S, A, B be the pure and the two odd shaded maps, as defined in Section 7. The traced paths are

$$\lambda = W_2^{-1}W_0, \qquad \mu_0 = S^{-1}, \qquad \mu_- = A^{-1}W_0, \qquad \mu_+ = W_0^{-1}B. \tag{5.1}$$

The longitude crosses white face 0 and returns through 2. The pure meridian crosses the shaded identification once in the reverse direction. The odd paths cross the respective half-disk pairing and white face 0. The retained negative crossing uses μ_- and the positive crossing uses μ_+ in the original disk frame. A physical turn transports the whole collar, not its sign.

Conjugation back to the original apex at infinity gives the translations

$$\mu_0 = E - 1, \qquad \lambda = -2(E+1), \qquad \mu_- = -2, \qquad \mu_+ = 2E. \tag{5.2}$$

The actual torus area is $8\sin\pi a\cos\pi a = 4\sin 2\pi a$, and $|\operatorname{Im}(\overline{\mu_0}\lambda)| = 4\sin 2\pi a$. Thus the closed paths span the full lattice, not a proper sublattice. Replacing μ_0 by either odd meridian

preserves this area. They too are primitive. As translation vectors on the corresponding half-twist partners,

$$\mu_\varepsilon = \mu_0 - \varepsilon\lambda/2.$$

This compares different marked tori; half a longitude is not being declared an integral curve on the pure torus. Since $(E+1)/(E-1) = -\mathrm{i}\cot\pi a$, the ratio λ/μ_ε is (1.6). This proves the remaining marking assertion of Theorem 1.2.

Proposition 5.1. *Every strict expression admits a choice of pure or odd leaves for which the main strands form one component. Either odd sign may be chosen, and subsequent full twists preserve the component count.*

Proof. Write the endpoint pairings as $I = (03)(12)$, $U = (01)(23)$ and $X = (02)(13)$. A leaf offers $\{U, X\}$, and a turn exchanges I, U and fixes X. Tracing the two actual product connections gives

	I	U	X
I	I	U	X
U	U	U and an internal circle	U
X	X	U	I

.

Discard only the product $U \cdot U$ in this existence argument. For sets $A = \{I, U\}$, $B = \{I, X\}$ and $C = \{U, X\}$, the output sets for AA, AB, AC, BB, BC, CC are respectively $A, \{I, U, X\}, C, B, \{I, U, X\}, A$, also in reversed orders. Thus every subexpression has at least two loop-free choices, starting with the leaf itself; the same is true if a child already offers all three. At least one choice at the root stack is U or X, both of which close to one component. Backtracking chooses the leaves. No identity of unmarked tangles or assertion about quantum connected-sum factors is used. □

6 A classical shear potential

Continue each occurrence of J holomorphically from its strict real argument. Its branch and additive constant are fixed by (1.2). For a leaf put

$$g(x,t) = J(2x) - 2J(x) - J(x-t) - J(x+t), \qquad v = g_t = \operatorname{Log}\frac{\sin\pi(x-t)}{\sin\pi(x+t)}. \tag{6.1}$$

At $t = 0$ this is $g(x)$ and $v = 0$. The local mixed pyramid has ordered base $(1, E/T, -1/T, -E)$, with $E = e^{2\pi\mathrm{i}x}$ and $T = e^{2\pi\mathrm{i}t}$. For small real t, its four half-arcs are $x-t, 1/2-x, x+t, 1/2-x$. The two positive tetrahedra across its first diagonal have angle triples $(x-t, 1/2-x, 1/2+t)$ and $(1/2-t, x+t, 1/2-x)$. The complementary Lobachevsky terms cancel, giving volume $\pi g(x,t)$. This fixes the local primitive without a quantum normalization argument.

For the retained sign ε, define the small hidden parameter $t = t(x,\chi)$ and its sheared potential by

$$\chi = t - \frac{\varepsilon v}{2\pi\mathrm{i}}, \qquad f_\varepsilon(x,\chi) = g(x,t) - \frac{\varepsilon v^2}{4\pi\mathrm{i}}. \tag{6.2}$$

At $(a, 0)$ the derivative of the first equation in t is $1 - \varepsilon\mathrm{i}\cot\pi a \neq 0$. The holomorphic implicit-function theorem applies. Differentiating both equations proves

$$\mathrm{d}f_\varepsilon = g_x\,\mathrm{d}x + v\,\mathrm{d}\chi. \tag{6.3}$$

The construction is separately even in each small χ because g is even in t and the shear equation is odd.

Choose independent real coordinates $\mathbf{y}$ on the original reduced wire space, keeping every turn and the complementary closing relation. The affine leaf and gate wires extend to its complexification. Set

$$\mathcal{A}_\varepsilon(\mathbf{y},\boldsymbol{\chi}) = \sum_i f_{\varepsilon_i}(a_i(\mathbf{y}),\chi_i) + \sum_{\text{gates}} m. \tag{6.4}$$

At zero color this is exactly $\mathcal{F}$. Its nondegenerate internal Hessian is the imported strict variational theorem. A second application of the holomorphic implicit-function theorem therefore gives a unique small solution $\mathbf{y}_*(\boldsymbol{\chi})$ of $\partial_{\mathbf{y}}\mathcal{A}_\varepsilon = 0$ through the complete optimizer. Define

$$\Phi_\varepsilon(\boldsymbol{\chi}) = \mathcal{A}_\varepsilon(\mathbf{y}_*(\boldsymbol{\chi}),\boldsymbol{\chi}). \tag{6.5}$$

It is separately even, and the envelope rule gives $\partial_i\Phi_\varepsilon = v_i$. All these statements concern an explicitly defined classical germ, independently of any finite-level source.

Lemma 6.1. *With τ_i the actual primitive ratio in (1.6),*

$$\partial_i\partial_j\Phi_\varepsilon(0) = \pi\mathrm{i}\tau_i\,\delta_{ij}. \tag{6.6}$$

Proof. At a leaf, $g_{tt}(a,0) = -2\pi c$ and $\chi_t = 1-\varepsilon\mathrm{i}c$. Hence $f_{\chi\chi} = -2\pi c/(1-\varepsilon\mathrm{i}c) = \pi\mathrm{i}\tau$. Also $g_{xt}(x,0) = 0$. Every mixed wire–auxiliary derivative vanishes at zero, so eliminating the stationary wires adds no Schur-complement term to this block. Distinct leaves have no direct mixed auxiliary derivative in (6.4). □

7 Actual and auxiliary white transitions

An intermediate mask bit distinguishes two different roles for a matrix. At a zero bit it is the actual map on the adjacent white piece. At a one bit there is a shaded face instead; a *completion map* records the white transition that must be supplied from the other side when the edge becomes internal. It is not an extra face or a pairing assigned to that shaded triangle.

7.1 Universal local formulas

Normalize the four ordered face flags to $(0,\infty,1,R)$. For the mixed leaf the relevant coordinates are

$$R = \frac{\cos^2\pi x}{\cos^2\pi t} = \frac{T(1+E)^2}{E(1+T)^2}, \qquad Z = \frac{1+E}{1+T}, \qquad Z^2-(1+E)Z+ER = 0. \tag{7.1}$$

Its ordered cross ratio is $R^{-1}-1 = e^{-g_x}$. The first apex is Z and the second-copy apex is Z/E: the latter is the reflected minus-t chart. Reflection here means inversion of the phase coefficients, not complex conjugation of free parameters. Every defining triple of face flags is distinct at completeness and remains so nearby. The two roots Z, TZ of the auxiliary fixed-point equation may coalesce at completeness; we require no distinctness of that pair there.

The four white or completion matrices are

$$\begin{aligned} W_0 &= \begin{pmatrix} 1 & 0 \\ 0 & E \end{pmatrix}, & W_1 &= \begin{pmatrix} 1 & E-1 \\ 0 & E \end{pmatrix}, \\ W_2 &= \begin{pmatrix} E-R & R(1-E) \\ E-1 & 1-ER \end{pmatrix}, & W_3 &= \begin{pmatrix} R & 0 \\ 1-E & ER \end{pmatrix}. \end{aligned} \tag{7.2}$$

All matrices are projective. At a leaf, W_0, W_2 are its actual white maps: they fix the endpoints of the appropriate base edge and send Z to Z/E. The apex equation verifies the statement for W_2. The other two matrices are completion maps at its shaded cap bases.

We use the rational prism identities of [2, equation (5.5)], equivalently [3, equation (3.7)]. To specify their charts, put $\alpha = e^{2\pi \mathrm{i}(z-x)}$, $\beta = e^{2\pi \mathrm{i}(x+y-z)}$, $\gamma = e^{2\pi \mathrm{i}(z-y)}$. The world prism is

$$\begin{aligned} (v_0, v_1, v_2, v_3) &= (1, \alpha, \alpha\beta, \alpha\beta\gamma), \\ P &= 1 + \alpha - \frac{\alpha[\alpha\beta(1+\gamma) - 1 - \alpha]}{\alpha^2\beta^2\gamma - \alpha}. \end{aligned} \tag{7.3}$$

For an ordered quadruple (a, b, c, d) let $M(z) = (z-a)(c-b)/((z-b)(c-a))$ and $R = M(d)$, with the usual limits at infinity. Its world transition is $(M^{\#})^{-1} W_j M$, where $\#$ reciprocates the three phase variables. Use the ordered tuples (2.7), with wire phases $\beta\gamma, \alpha\beta, \alpha\beta\gamma$. The resulting entire maps satisfy

$$\begin{aligned} W_{F,1} = W_{G,1} = W_{o,1} = W_{\mathrm{top}}, \qquad W_{F,3} = W_{G,3} = W_{o,3} = W_{\mathrm{bottom}}, \\ W_{F,2} = W_{G,0}, \quad W_{o,0} = W_{F,0}, \qquad W_{o,2} = W_{G,2}. \end{aligned} \tag{7.4}$$

The top and bottom maps are the actual white prism triangles. The cited proof is an identity in independent complex phase variables, so it applies also to a wire above $1/2$. No completed-child hypothesis enters that local calculation.

For turns and closure the corresponding identities are

$$\begin{aligned} T_Q(z) = \frac{R-1}{z-1}, \qquad C_Q(z) = \frac{z-R}{z-1}, \\ T_Q W_{j+1}(E, R) = W_j(E^{-1}, 1-R) T_Q, \\ C_Q W_j(E^{-1}, R) = W_{2-j}(E, R) C_Q. \end{aligned} \tag{7.5}$$

They are [2, equation (5.6)]. They retain all translations, not only derivatives at endpoints.

7.2 The mask induction for the entire maps

We prove the following strengthened statement simultaneously through the original expression: at each output, zero bits carry actual white maps and one bits carry completion maps; at every internal shaded edge the actual adjacent white piece already has the recorded completion map. White pieces joined flatly carry the same entire map.

At a child cap seam the prism is white. If the child bit is zero, (7.4) identifies two actual maps on a flat seam. If it is one, the shaded triangle remains and the actual prism map supplies exactly its completion. At the common free edge the possible bit pairs are $(0,0), (1,0), (0,1)$. In the first case the actual white maps merge by $W_{F,2} = W_{G,0}$. In the other cases the unmasked child's actual map supplies the masked child's completion. The excluded pair $(1,1)$ would require HH. Finally the inherited output free maps and the two actual prism cap maps give precisely (3.3).

A turn transports this distinction by (7.5). At the closing face the strict complementary pair excludes two shaded bits on the same seam, as already proved. Zero–zero joins actual white pieces; one–zero supplies the required completion from the actual white side. No output face remains, so every completion map that enters a leaf edge return is now its actual neighboring white transition.

Stationarity of (6.4) matches the same complete complex cross ratios, by (6.3). Three vertices determine the Möbius map and the cross ratio matches the fourth. This proves the induction for the deformed charts as well. A deformed white polygon need not be planar; all its retained triangular pieces have one common transition by the proved identities.

8 The original group and the retained positive chain

Proposition 8.1. *The stationary family* (6.5) *determines a holomorphic framed representation of the original* $\pi_1(Y_E^\varepsilon)$ *through its complete holonomy. It retains a compatible positive ideal fundamental chain. All main peripheral groups remain parabolic, and the original auxiliary basis has exactly the logarithms* (1.7)*.*

We prove the group, chain and peripheral assertions in that order. In a leaf frame the pure shaded map and the two odd maps are

$$S = \begin{pmatrix} 0 & -ER \\ 1 & -1-E \end{pmatrix}, \quad A = \begin{pmatrix} R & -R \\ 1 & ER-1-E \end{pmatrix}, \quad B = \begin{pmatrix} R/E-1-1/E & R \\ -1 & R \end{pmatrix}. \tag{8.1}$$

They send the original ordered triples as follows: $S : (Z, \infty, 1) \mapsto (Z, 0, R)$, $A : (Z, \infty, 1) \mapsto (Z/E, R, 0)$, and $B : (Z, R, 0) \mapsto (Z/E, \infty, 1)$. With $S^{\#} = S(E^{-1}, R)$, direct multiplication gives

$$\begin{gathered} W_0 S = S^{\#} W_0, \quad W_2 S = S^{\#} W_2, \quad W_3 S = S^{\#} W_1, \\ W_0^{-1} B W_2^{-1} A = I, \quad W_2^{-1} B W_0^{-1} A = I, \quad W_1^{-1} B W_3^{-1} A = I. \end{gathered} \tag{8.2}$$

Equalities are in the projective group. These local identities are [2, equations (5.7)–(5.8)], for independent E, R. The actual/ completion-map induction is what permits their use on the enlarged syntax.

The original cut balls have a dual cell decomposition: paired faces give generators and edge cycles give the relations of its two-skeleton. Truncating the cusps gives the same group presentation; higher-dimensional cells add no fundamental-group relation. Every white seam and every pure or odd shaded return has now been verified by whole Möbius maps. Hence these maps define an actual holomorphic framed representation ρ_χ of $\pi_1(Y_E^\varepsilon)$ through its oriented complete holonomy. This conclusion uses the closed topological cut, not a cross-ratio argument for a hypothetical open four-punctured boundary.

There is also a compatible positive chain on the original manifold. At completeness first assemble the actual polyhedron P_E, whose ideal vertices are distinct except for the prescribed terminal pairs. Order these cut vertices and restrict the order to every constituent cell. Pulling triangulations agree on each shared quadrilateral. The order selects diagonals; orient every tetrahedron by the ambient oriented cell, not by its increasing vertex list. Coning from the first vertex to faces not containing it gives positive nondegenerate tetrahedra. Give the second projection copy the same labeled subdivision. White pairings match it, and shaded pairings are already triangular, so no order on the quotient cusp vertices is required.

Transport these finitely many flags by the actual rational charts. Their positive complete shapes remain in the positive half-plane on one small parameter polydisk. The face maps match the subdivided faces, and edge products stay exactly one. The continuous edge-log branches have their original sums there. Thus the representation has the same oriented ideal fundamental chain and boundary triangulation as the complete metric. We neither require deformed half-balls to be convex nor triangulate the filled manifold by new positive cells.

8.1 All main cusps remain parabolic

At corner C_j the actual small main meridian is $W_{j-1}^{-1} W_j$ or its inverse. One of these maps may have been supplied as a completion, but both are now actual white maps. Their derivatives at the corner agree: they are E^{-1} at C_0, C_2 and E at C_1, C_3, using a finite coordinate at infinity. The meridian multiplier is identically one. The meridian is nonidentity at completeness and remains so nearby. It is therefore parabolic. Its commuting peripheral elements are translations in the same fixed-flag chart. This proves the statement for every main cusp, including any component closed inside an intermediate physical product.

8.2 The exact auxiliary logarithms

All the based curves (5.1) remain in the original leaf collar. Put

$$\rho = \frac{Z}{1+E-Z} = \frac{1}{T}, \qquad Q = \frac{E-Z}{Z-1} = \frac{ET-1}{E-T}. \tag{8.3}$$

In a frame sending the selected flag Z to infinity, the similarity multipliers of the four words are

$$\lambda : Q^2, \qquad \mu_0 : \rho, \qquad \mu_- : \rho Q, \qquad \mu_+ : \rho/Q. \tag{8.4}$$

At a finite fixed flag, this multiplier is the reciprocal of the Möbius derivative. Substitution of (7.1) into $\det M/(cZ+d)^2$ verifies these formulas. They are the local identities [3, equations (4.2)–(4.3)]. Since $\rho = e^{-2\pi \mathrm{i} t}$ and $Q = e^{-v}$, their small logarithms are

$$U = -2\pi \mathrm{i} t + \varepsilon v = -2\pi \mathrm{i}\chi, \qquad V = -2v. \tag{8.5}$$

These are similarity logs, twice selected eigenvalue logs. Their constants are zero at completeness. The envelope rule proves (1.7); physical turns only conjugate the words.

If $\chi_i = 0$, separate evenness gives $V_i = 0$ as well. The auxiliary meridian remains nonidentity nearby, so an auxiliary cusp left unfilled also remains a genuine parabolic translation group. All flags and defining face triples stay in a finite collection of bounded local frames. This proves Proposition 8.1.

9 Independent signed slopes and smooth completion

Fix the subset I and set all other χ_i equal to zero. For $i \in I$ write $r_i = 1/p_i$. Solve

$$V_i(\boldsymbol{\chi}) + r_i U_i(\boldsymbol{\chi}) = -2\pi \mathrm{i} r_i. \tag{9.1}$$

At $\mathbf{r} = 0$ the solution is $\boldsymbol{\chi} = 0$. Equations (1.7) and (6.6) give the invertible Jacobian

$$D_{\boldsymbol{\chi}} V(0) = -2\pi \mathrm{i}\, \mathrm{diag}(\tau_i)_{i\in I}. \tag{9.2}$$

The implicit-function theorem gives a unique solution on one reciprocal-slope polydisk. Shrink it until the representation and positive original chain are in their common fixed neighborhood. For real signed p_i this imposes a lower bound on each $|p_i|$, with no restriction on their ratios. For $p_i \neq 0$ the equations are

$$U_i + p_i V_i = -2\pi \mathrm{i}. \tag{9.3}$$

They are precisely the critical equations of (1.8).

The peripheral direction is actually loxodromic at these solutions. Separate evenness makes the apparent quotient $\widetilde{\tau}_i = V_i/U_i$ removable, holomorphic near zero, and equal to τ_i there. Shrink once more so that $\operatorname{Im} \widetilde{\tau}_i > 0$. Equation (9.1) yields

$$\chi_{i,*} = \frac{r_i}{\widetilde{\tau}_i + r_i}, \qquad \operatorname{Re} V_i = \frac{2\pi r_i^2 \operatorname{Im} \widetilde{\tau}_i}{|\widetilde{\tau}_i + r_i|^2} > 0. \tag{9.4}$$

This is valid for either sign of a nonzero real reciprocal slope. It is a classical consequence of the positive primitive cusp ratio.

We now apply the precise near-complete completion result [4, Proposition 6.1]. Its input is an oriented complete finite-volume manifold with marked primitive cusp bases, and an actual nearby framed representation with small similarity logs in bounded frames. At unfilled cusps both multipliers must be one. At each filled cusp a primitive integral combination of

the small logs must equal $2\pi\mathrm{i}$ or $-2\pi\mathrm{i}$. It then completes the original topological filling to a smooth complete hyperbolic metric.

Every hypothesis has been supplied on this enlarged syntax: Theorem 1.2 constructs the actual complete exterior; the unchanged two-rectangle collar proves its primitive basis; Sections 7–8 construct the actual framed representation and the original group relations; all unfilled cusps have translation holonomy; and (9.3) uses the primitive vector $(1, p_i)$. The fixed small parameter neighborhood is chosen before the slopes.

9.1 Why the integer equation gives one actual angular turn

We recall the part of the imported proof that prevents an exponential period equation from being mistaken for completion. On a complete cusp link use a translation lattice $\Lambda \subset \mathbb{C}$ and its fixed generator paths. A nearby peripheral action is affine, $A_\lambda(z) = a_\lambda z + b_\lambda$, in a bounded moving flag frame. A smooth partition η on the complete plane with $\sum_{\lambda\in\Lambda} \eta(w-\lambda) = 1$ gives

$$d_\rho(w) = \sum_{\lambda\in\Lambda} \eta(w-\lambda) A_\lambda(w-\lambda).$$

On a fundamental parallelogram this is C^1-close to the complete identity and has positive real Jacobian; equivariance propagates this fact. With the real peripheral logarithmic character in the height coordinate it gives a nonsingular developed cusp, matched over a compact collar to a nearby developing map on the original truncated core.

At a filled cusp a multiplier has nonunit modulus by (9.4). Commutativity gives a second common fixed point q besides infinity. Since the original complete translations are nonzero, $|q| \to \infty$ near completeness. The developed fixed generator paths stay bounded, with bounded length. They miss q, and their increments of $f = \mathrm{Log}(d_\rho - q)$ tend to zero. Quantitatively, a path of length at most L contained in $|d_\rho| \le R$ has $|\Delta f| \le L/(|q| - R)$. Thus those increments are the prescribed small logs U, V, without an added integral multiple of $2\pi\mathrm{i}$, *before* multiplication by the large integer p.

The metric $|df|^2$ descends to a compact nonsingular flat torus. Its complete lifted developing map is a diffeomorphism to $\mathbb{C}$, so U, V form its actual rank-two period lattice. The developed end contains a punctured metric tube with cylindrical metric

$$dR^2 + \cosh^2 R\, dX^2 + \sinh^2 R\, d\theta^2. \tag{9.5}$$


A complex period L acts by $(X, \theta) \mapsto (X + \mathrm{Re}\, L, \theta + \mathrm{Im}\, L)$. The original primitive slope has period $U + pV = -2\pi\mathrm{i}$, exactly one angular turn. Its complementary primitive period V has nonzero real part. Adding $R = 0$ produces the smooth solid torus with this original meridian. The slope fixes both q, ∞ and has multiplier one, so it is the identity in the group, not a nontrivial unipotent with the same eigenvalues. The remaining ends are full horoball quotients, and the middle region is compact. Completeness and finite volume follow.

This verifies the metric conclusion of Theorem 1.3 for the actual disk-filled exterior. In particular high-index estimates for a quantum coefficient, source positivity and whole character-variety rigidity have played no role.

10 Absolute volume on the original chain

Set $G = 2\pi\mathrm{i}\Phi_\varepsilon$. The exact peripheral identity gives

$$\mathrm{d}G = \frac{1}{2}\sum_i V_i\, \mathrm{d}U_i, \qquad \mathrm{Im}\, G(0) = 2\pi F_0 = \mathrm{Vol}(Y_E^\varepsilon). \tag{10.1}$$



Original primitive basis: U, V.

Filled meridian: $m = U + pV$.

The fixed small increments are established before forming m.

Figure 2: The developed logarithmic link of a filled original crossing circle. The old basis is U, V, while $m = U + pV = -2\pi\mathrm{i}$ is the primitive meridian of the added solid torus. The picture shows $p = 1$; the proof uses the small periods before multiplying by any integer p. In the tube metric, m makes one angular turn and V gives a nonzero axial translation.

This is the $V\,dU$ polarization. A primitive of $-\frac{1}{2}\sum U\,dV$ differs by the boundary term $\frac{1}{2}\sum UV$; its regulator correction cannot be reused unchanged here.

Let $\mathcal{V}(\rho)$ be the signed sum of the ideal tetrahedron volumes of the retained original chain. The local boundary differential [4, equation (7.3)] reads, for similarity logs,

$$\mathrm{d}\mathcal{V} = \frac{1}{2}\sum_i \bigl(\operatorname{Re} V_i\,\mathrm{d}\operatorname{Im} U_i - \operatorname{Re} U_i\,\mathrm{d}\operatorname{Im} V_i\bigr). \tag{10.2}$$

Its hypotheses hold on our positive nondegenerate chain: internal faces cancel by the actual maps, edge-log sums stay fixed, and the primitive cusp orientation is $\operatorname{Im}(\lambda/\mu) > 0$ by (1.6). All main logs vanish. The proof of that local identity differentiates Bloch–Wigner volume and cancels normal-face flows; it uses the positive ordered model $(0, \infty, 1, z)$ and the primitive cusp intersection pairing. It is not a theorem restricted to the complete packing of a particular diagram.

Taking the imaginary part of (10.1) and differentiating its boundary correction now gives

$$\mathcal{V}(\rho_{\boldsymbol{\chi}}) = 2\pi \operatorname{Re}\Phi_\varepsilon(\boldsymbol{\chi}) - \frac{1}{2}\sum_i \operatorname{Re} U_i\ \operatorname{Im} V_i. \tag{10.3}$$

The integration constant is fixed by the actual degree-one complete cell volumes, not just by a derivative computation. At a filling solution write $\chi_i = u_i + \mathrm{i}v_i$. Since $V_i = 2\pi\mathrm{i}(\chi_i - 1)/p_i$,

$$-\frac{1}{2}\operatorname{Re} U_i\ \operatorname{Im} V_i = -\frac{2\pi^2}{p_i}v_i(u_i - 1) = 2\pi \operatorname{Re}\left[\frac{\pi\mathrm{i}}{2p_i}(\chi_i - 1)^2\right]. \tag{10.4}$$

Unfilled auxiliaries have $U_i = V_i = 0$.

10.1 The same boundary triangulation after filling

It remains to identify $\mathcal{V}$ with the volume of the smooth metric just constructed. We verify the chain hypotheses of [4, Section 7.3]. At a filled cusp conjugate the common fixed points to $0, \infty$. On each triangle of the retained boundary triangulation the ideal conical development is $(a(w), Yb(w))$, with $b > 0$. All functions and their derivatives are bounded on a finite fundamental set. Choose an equivariant axis map $(0, Ye^{\eta(w)})$, where η has the real peripheral logarithmic character. On the *same* triangulation set

$$H_s(w, Y) = \bigl(sa(w), Y\exp\bigl((1-s)\eta(w) + s\log b(w)\bigr)\bigr), \qquad 0 \le s \le 1. \tag{10.5}$$

Both horizontal coordinates and heights have the required equivariance, and the homotopy agrees along every face. Its height is comparable to Y, horizontal derivatives are bounded, and height derivatives are $O(Y)$. The volume form $y^{-3}\,dx_1 \wedge dx_2 \wedge dy$ therefore bounds its three-dimensional boundary sweep by $O(Y^{-2})$.

Once the primitive slope is killed, extend that axis-valued boundary over the added solid torus. The remaining cyclic group acts along the axis, so all added vertices can lie on the axis and all added tetrahedra have zero signed volume. At unfilled cusps the same vanishing boundary estimate applies. Stokes on finite truncations, followed by $Y \to \infty$, identifies the retained ideal chain with the filled equivariant fundamental class. Comparing it with the complete metric developing map on a common subdivision gives

$$\mathcal{V}(\rho_{\boldsymbol{\chi}_*}) = \mathrm{Vol}(Y_E^{\varepsilon}(\mathbf{p}; I)).$$

An unrelated chain with the same peripheral differential would not suffice; (10.5) supplies the absolute comparison. Combining it with (10.3) and (10.4) proves (1.9).

11 Proof boundary and finite algebra

The strict variational existence, Hessian and boundary-profile identities are imported from [1], with their precise nonmetric content in Section 2. The local cells, entire rational prism maps and collar words are the local results of [2, 3]. The present paper proves the individual-mask induction, its enlarged closed topology and all actual versus completion-map status, before using those local formulas. Smooth metric completion and the local volume differential are the explicitly stated inputs from [4], with all hypotheses verified here.

No source asymptotic, full spectral bound or nonzero Fourier contribution is claimed. To identify the classical action with a quantum packet one must independently match each endpoint-normalized leaf action, including its color-dependent additive constant, and then eliminate these same wires. A further whole-source estimate and filling noncancellation theorem are needed for an invariant volume limit. The geometry alone does not supply them. Similarly no equivalence with a completed-root presentation is used, and no moving main-meridian or preferred-longitude deformation claim is made.

The accompanying `check.py` verifies the finite mask closure and its allowed edge pairings, the literal endpoint-connection table, projective turn/closing and pure/odd return identities, and the primitive Hessian and critical boundary normalization. It includes altered mask, pairing and factor-of-two controls. It writes a source-hashed JSON record with `-output`. These exact finite checks do not prove convexity, the variational existence theorem, the original manifold identification, implicit-function theory, smooth completeness or the chain comparison.

References

[1] *Stationary profiles of strict Newton trees*, companion paper, Lemma 2.2, Theorem 3.3, Lemma 4.1 and Proposition 4.2. Companion PDF.

[2] *Pyramids, prisms, and complete geometry of Conway closures*, companion paper, Section 2, equations (3.2)–(3.3), (5.5)–(5.8), and Section 7. Companion PDF.

[3] *Marked deformations and long fillings of Conway closures*, companion paper, Sections 2–4 and 7. Companion PDF.

[4] *Circle patterns and primitive fillings of augmented links*, companion paper, Proposition 6.1, equation (7.3) and Section 7.3. Companion PDF.

[5] J. S. Purcell, An introduction to fully augmented links, *Contemp. Math.* **541** (2011), 205–220. Author's manuscript.

[6] W. D. Neumann and D. Zagier, Volumes of hyperbolic three-manifolds, *Topology* **24** (1985), 307–332. doi:10.1016/0040-9383(85)90004-7.