# One entropy problem at every fixed bridge width

**Original positive endpoint source**

$$Z_{D,N} \quad = \quad [\prod_{f \text{ ordinary}} L_f^{N-1}] \prod_{e=1}^{R} P_e\big|_{L_u = L_v = 0}$$

Original faces, marked targets and level $N$ retained.

**Exact finite support**

Bounded integer incidence flows

One test level: $N_* = 10R + 2$

Eventually admissible patterns $\mathcal{A}_D$

**Uniform coefficient entropy**

$E(\mu, \nu, \xi)$ on the closed domain

All endpoint rows and outer degrees

Shared limiting flow polytope $\mathcal{P}_s$

**Proved for this positive source, through even levels**

$$\lim_{N \to \infty} \frac{\log Z_{D,N}}{N} \quad = \quad W_D \quad = \quad \max_{s \in \mathcal{A}_D} \max_{q \in \mathcal{P}_s} \sum_e E_{s_e}(q_e)$$

**Open geometric comparison**

$$W_D \quad \stackrel{?}{=} \quad \frac{\mathrm{Vol}(Y_D)}{2\pi}$$

Lower bound and sublinear-excess corridor proved under the labeled geometric certificate; macroscopic patterns remain.

The original Kashaev filling cycle is a further step.
The positive endpoint theorem is not the filled-knot volume conjecture.