One local twist transport for every integer power

$$\sigma = 2 + \frac{(b+z+A)^2}{P-bz}, \qquad b_{j+1} + \sigma b_j + b_{j-1} + 2A = 0$$

$A = x + y$, $P = xy$; the positive outer faces x, y stay fixed.

The advance follows the coefficient from its left output to its right input.

Generic spectral cover

$$\lambda^2 + \sigma\lambda + 1 = 0$$

$$\mathrm{Norm}(a) = 1$$

$$a \longmapsto \lambda^n a$$

The source retains its excluded stationary divisors.

Parabolic fiber $\sigma = -2$

$$t = b - z$$

$$t \longmapsto t + 2nA$$

$$(M - I)^3 = 0$$

Face coordinates depend quadratically on n.

The actual quantum operator keeps the same Hamiltonian

$$R^n = q^{nd^2/4} \exp\left[\frac{n}{2\hbar}\,\mathrm{arccosh}(-\Omega/2)^2\right]$$

$$d = N - 1,\ q = e^{\pi i/N},\ \hbar = 2\pi i/N.$$

Original scalar and repeated-node data retained.

The other singular fibers have their own stated schemes.
Local transport is explicit. A full period bound for arbitrary certified plats remains open in this route; the strict Conway-tree theorem is proved separately.