# One local twist transport for every integer power

$$\sigma \;=\; 2 \;+\; \frac{(b+z+A)^2}{P-bz}, \qquad b_{j+1} \;+\; \sigma b_j \;+\; b_{j-1} \;+\; 2A \;=\; 0$$

$A \;=\; x \;+\; y$, $P \;=\; xy$; the positive outer faces $x, y$ stay fixed.

The advance follows the coefficient from its left output to its right input.

## Generic spectral cover

$$\lambda^2 \;+\; \sigma\lambda \;+\; 1 \;=\; 0$$

$$\mathrm{Norm}(a) \;=\; 1$$

$$a \;\longmapsto\; \lambda^n a$$

The source retains its excluded stationary divisors.

## Parabolic fiber $\sigma \;=\; -2$

$$t \;=\; b \;-\; z$$

$$t \;\longmapsto\; t \;+\; 2nA$$

$$(M \;-\; I)^3 \;=\; 0$$

Face coordinates depend quadratically on $n$.

## The actual quantum operator keeps the same Hamiltonian

$$R^n \;=\; q^{\,nd^2/4} \exp\!\left[\frac{n}{2\hbar}\,\operatorname{arccosh}(-\Omega/2)^2\right]$$

$$d \;=\; N \;-\; 1,\; q \;=\; e^{\pi i/N},\; \hbar \;=\; 2\pi i/N.$$

Original scalar and repeated-node data retained.

The other singular fibers have their own stated schemes.
Local transport is explicit; the whole-source volume estimate remains open.