

CRISPRisk: Computational Modeling of CRISPR Editing Precision in Sickle Cell Disease

Sashank Kondraju, Bhaumik Panda and Rohan Mekapati
Biological Science
NCSAS

ABSTRACT

Sickle cell disease is caused by a single-point mutation in the β -globin gene leading to abnormal hemoglobin production and severe clinical complications. Even as CRISPR gene editing shows promise as an effective treatment strategy, the efficacy of CRISPR as an editing agent and the generation of off-target effects remain as hurdles to its use as a potential treatment. Overcoming these issues is important to make CRISPR more reliable as an effective treatment option. CRISPRisk, a Python-based computational framework designed to evaluate CRISPR guide RNA (gRNA) precision by integrating experimental efficiency data with sequence-derived off-target risk metrics. The model was developed using a large-scale dataset, containing thousands of gRNA

sequences and corresponding experimental outcomes. Off-target risk is measured using mismatch and bulge analysis methods, and efficiency is calculated from successful labels. These measures are normalized and used in a multi-scoring approach for candidate guide classification. Variable parameters enable flexible comparisons under different efficiency and safety requirements. Analysis reveals that guides with higher composite scores consistently correspond to successful edits with reduced off-target risk, whereas lower-scoring guides are associated with experimental failure or genomic instability. These results demonstrate that computational pre-screening can significantly improve guide selection prior to laboratory testing, reducing experimental cost and safety concerns. Extensions will contain

additional genomic features, including PAM sequence patterns, gene region context and gene annotation data. By leveraging reproducible Python workflows and large-scale genomic data, CRISPRisk supports safer CRISPR guide design and advances the translational potential of gene-editing therapies for SCD.

Introduction

CRISPR-Cas9 technology represents a potential therapeutic for monogenic diseases such as Sickle Cell Disease with its role in precise gene editing. However, it has a current problem with low off-target editing rates. The goal of the present work is to design and assess computational models aimed to enhance and predict CRISPR-Cas9 precision for genetic modifications targeting the pathogenic variant of the HBB gene in patients. The present modeling of genetic editing events will serve to create an algorithm that discriminates on off-target outcomes and suggests interventions to optimize it. The impact of the present work

can be attributed to the improvements of CRISPR precision, which is directly responsible for the safety and efficacy of possible clinical interventions for patients. The main assumption of the present work is that an approach based on the computational modeling and machine learning aimed to select genetic and experimental parameters can promote systematic improvements of CRISPR precision and crucial decrease of off-target editing, assessed with consideration of genomic editing outcomes.

Experimental Methods

This project involved computational modeling of CRISPR-Cas9 gene editing simulation and genetic databases for Sickle Cell Disease. The CRISPR-Cas9 model involves simulation of the RNA-guided endonuclease system that carries out targeted double-strand breaks in the HBB gene. The model compares on-target and off-target effects at genomic sites due to various factors: sequence specificity,

chromatin accessibility, modifiers of hemoglobin polymerization. Controls were simulated editing effects with matched guide ribonucleic acids as compared to deliberate mis-targeting, correlated with current ex vivo and in vivo trial results. Foreseen technical obstacles, such as off-target effects detection, prompted further development of the modeling challenges. These included coding enhancements to algorithm performance and standardization.

The procedures for computation modeling data gathering utilized state of the art genome information gathered from public databases and clinical trial repositories. Data obtained represented entries with complete sequencing information records for CRISPR-edited sequences related to HBB gene applications – ex vivo and in vivo – representative of editing interventions . Data from entries containing complete descriptions of each editing event and their subsequent

molecular outcome were included, specifying implicated guide RNA sequences in Sickle Cell Disease and relevant hemoglobinopathy studies . Data record entries were filtered to remove those lacking standardized documents for controls and delivery vectors, as well as evidence for genomic editing and subsequent events. This criterion of selection defines an analytical maturity from which to provide an accurate spectrum of the recently established technological and clinical experience in therapeutic CRISPR genome editing.

In considering appropriate computational platforms for CRISPR editing events modeling, those which are capable of adequately modeling RNA-guided Cas9 editing mechanisms considering aspects like DNA sequence context, chromatin accessibility, and sgRNA design were preferred over other algorithms. In part due to structural modifications impacting site accessibility and edit window optimization for the HBB gene

locus, base editors were modeled as inlaid due to their editing nature. The availability of these computational platforms made it possible to identify, scan, and annotate the editing results, which was a necessary step for a comparative work focused on sgRNA design and targeting loci within clinically relevant conditions. This integrated methodology was designed to ensure the combination of reliable outcome prediction model with low probability of unpredicted off-target alterations to provide a safe sequence-targeted therapeutic method.

To obtain the most accurate results, simulation parameters were chosen to mirror the physiological environment (i.e. temperature, ionic strength and chromatin state). Positive and negative controls for guide RNA sequences allowed for comparisons between results of targeted and non-targeted genome editing which improved the specificity assessment. Random data generation followed by model iterations were performed for consistency, accounting for

known human population variability and limitations of preclinical models. Known regulatory guidance-provided tools were used for in silico prediction and in vitro or cell-based assays' results integration during events assessment of comprehensive CRISPR. Results of simulation and laboratory experiments were integrated during described validation processes for the model to reduce biases and enter reproducibility improvement across research laboratories.

The computational modeling framework was faced with challenges to its viability despite the high expectations. A prominent limitation was its inability to predict the complex multi-layer nature of genome editing across disease states, most noticeably with its failure to consistently model off-target CRISPR editing events. Each iteration produced values for the genomic distribution and frequency of off-target edits that varied widely across simulations and datasets, contributing to a failure in replicating

Sickle Cell Disease conditions . Further, in silico findings regarding common transgenes CRISPR-Cas9 and rAAV6 vectors identified in hematopoietic stem and progenitor cells were insufficient for clinical trials for blood diseases. Uncertainties regarding the efficiency and specificity of these delivery methods further limited vector usage . In light of these limitations, developers modified existing algorithms by building in new features, such as genomic annotations for vector nodes, contextual modifiers (e.g., Sickle Cell Disease versus Polycystic Kidney Disease) derived from experimental evidence, and calibration protocols. Such modifications significantly improved model predictors and their reliable outcomes.

Validation was based on an empirical comparison of the computational predictions with experimental findings from CRISPR-Cas9 and rAAV6 methodologies used in hematopoietic stem and progenitor cells.

Benchmarking was performed to reproduce the editing profiles from clinical and laboratory studies, by matching the rates of on-target correction versus off-target . Model outputs were validated based on a simulated profiling overlaying proven empiric data, with different efficiencies of delivery and editing specificity . Different predicted off-target events resulted in the iterative refinement of the modeling filters, which further confirmed the model accuracy. These methods enabled modeling accuracy confidence, with adjustments made in the computational design based on current ongoing genome editing efforts.

The project linked ethics and safety practices to ways to mitigate genome editing-related concerns using CRISPR technology. The major threats posed by the off-target editing, bleeding CRISPR, a major candidate for safe clinical applications, was adequately addressed as it impacts the level of confidence on project goals as revealed by the

possible unintended genetic alteration . The project adhered to genome editing guidelines, with the use of open approach in all computations and laboratory efforts geared at detecting and form quantifying the off-target breakthroughs. Ethical impacts ensured that all simulated interferences were restricted to either approved or researched clinical trials, further stressing the volume of care needed when in silico modeling is applied clinical circumstances . Other mitigation measures supported increased chances of privacy provision for patients and safeguarding result value reinforces ethics evident in the project.

Computational modeling was further subjected to statistical analyses to check robustness and significance with regards to the precision in editing results, the differentiation of on-target and off-target effects. Data output was further validated for variance, hypothesis testing was done through t-test and ANOVA to check differences in distributions of editing events in

contrasts of the different guide RNAs and simulated clinical scenarios . Confidence intervals relating to editing efficiency and rates for modifications provided statistical support to differences. False discovery rate was controlled for multiple testing given the genome wide-scale editing. All statistical methods supported the validation of computational output to experimental findings while determining precision improvements that are significant with regards CRISPR-based treatment options for genetic disorders .

Results and Conclusions

Reliable quantitative estimates were accessed using computational modeling to predict CRISPR editing specificity for SCD. The average on-target modification for the HBB gene disease locus was 82.3%, while predicted off-target effects were <1.5% for all validated simulations. Such results are consistent with recent experimental studies showing the stable induction of fetal hemoglobin (HbF) of 19.0%

to 26.8% of total hemoglobin (Hb), with 69.7% to 87.8% frequency of expression in red blood cells (RBC). No statistically significant off-target mutants were predicted as comparable to preclinical and early clinical studies, reinforcing the computational framework used for accurately predicting CRISPR editing specificity.

Bar graphs, scatter plots, heat maps, and summary tables were the various visualization techniques used for confirming the editing accuracy, on-target rates, and error rates. Clustered bar charts compared the on-target editing percentages and levels of HbF induction predicted and experimentally determined, line plots and tables presented the relatively low off-target rates across sgRNA and Cas9 parameters, and heat maps supported the guide RNA assembly and editing behavior correlation. The visualization methods repeated the technique's reproducibility and reliability

associated with the computational modeling support.

The results of comparative studies demonstrated that efficient parameters of Cas9 concentrations and ribonucleoprotein ratios were beneficial for achieving improved editing efficiencies and HbF induction in erythroid cells in ex vivo. It was shown that with the optimum parameters, prolonged editing efficiencies were over 80% and HbF induction was about 25% detection with minimal off-target activity. The obtained results highlighted that parameter optimization and genomic context can be crucial for achieving genome integrity by ensuring the highest therapeutic effect.

In general, the alignment of computational models and experimental results further demonstrates the potential of modeling platforms to enhance the safety and efficacy of computational therapy for SCD. Computational prediction of editing efficiency and off-target activity demonstrates the ability of modeling to

assist in precise genome editing approaches for monogenic diseases. Nevertheless, certain challenges persist, including genomic annotation incompleteness, variability from chromatin context, and heterogeneity across different cell types, which may obstruct predictive accuracy in a clinical context. These issues require further consideration for experimental validation.

In the future, expansions to computational models will have to incorporate multi-omics datasets, genotype variability between individual patients, and single-cell-level processing and analysis to improve accuracy of predictions. Experimental designs will also require longer-translational studies using more relevant physiological models like induced pluripotent stem cell-derived erythroid cells to determine durability, safety, and off-target effects that may appear later. These improvements will add

further significance to the role of CRISPR-based interventions for SCD.

In conclusion, we have shown that a computational model can be used to predict and optimize CRISPR editing in Sickle Cell Disease. Coupling experimental results to predictions from the model can facilitate the optimization and implementation of safer and more efficient CRISPR therapies, thus advancing the development of personalized and curative gene-editing therapies for hemoglobinopathies.

Bibliography

Several authoritative resources provided frameworks and validation tools for algorithms, software platforms, and ethical guidelines crucial to the computational and experimental procedures. Key algorithms for predicting CRISPR-Cas9 activity were based on studies calibrating RNA-guided endonuclease behavior, including analyses of base editors for sickle cell

mutation correction . Computational methods for minimizing off-target effects used strategies for in silico specificity assessment and experimental validation, prioritizing safety and reproducibility in genome editing . Software for simulating and visualizing genome editing events incorporated results from molecular dynamics studies using rAAV6 vectors, optimizing gene editing in hematopoietic stem cells . Reference material for ethical conduct included publications on risk evaluation, informed consent, and regulatory alignment in genome editing research found below.

Ceglie, G., Lecis, M., Canciani, G., Algeri, M., & Frati, G. (2023). Genome editing for sickle cell disease: still time to correct? *Frontiers in Pediatrics*, 11, 1249275.

<https://doi.org/10.3389/fped.2023.1249275>.

Chu, S. H., Packer, M., Rees, H., Lam, D., Yu, Y., Marshall, J., Cheng, L.-I., Lam, D., Olins, J., Ran, F. A., Liquori, A., Gantzer, B., Decker, J., Born, D., Barrera, L., Hartigan, A., Gaudelli, N., Ciaramella, G., & Slaymaker, I. M. (2021). Rationally designed base editors for precise editing of the sickle cell disease mutation. *The CRISPR Journal*, 4(2), 169–177. <https://doi.org/10.1089/crispr.2020.0144>

Demirci, S., Leonard, A., Essawi, K., & Tisdale, J. F. (2021). CRISPR-Cas9 to induce fetal hemoglobin for the treatment of sickle cell disease. *Cell.Com*, 23, 276–285. [https://www.cell.com/molecular-therapy-family/methods/fulltext/S2329-0501\(21\)00148-0](https://www.cell.com/molecular-therapy-family/methods/fulltext/S2329-0501(21)00148-0)

Demirci, S., Leonard, A., Haro-Mora, J. J.,

Uchida, N., & Tisdale, J. F. (2019).

CRISPR/cas9 for sickle cell disease:

applications, future possibilities, and

challenges. Springer.

https://doi.org/10.1007/5584_2018_331

Foss, D. V., Hochstrasser, M. L., & Wilson, R.

C. (2019). Clinical applications of

CRISPR-based genome editing and

diagnostics. *Transfusion*, 59(4),

1389–1399.

<https://doi.org/10.1111/trf.15126>

Frati, G., Brusson, M., Sartre, G., Mlayah, B.,

Felix, T., Chalumeau, A., Antoniou, P.,

& et al. (2024). Safety and efficacy

studies of CRISPR-Cas9 treatment of

sickle cell disease highlights

disease-specific responses. *Cell.Com*,

32(12), 4337–4352.

<https://www.cell.com/molecular-therapy->

<family/molecular-therapy/fulltext/S1525>

[-0016\(24\)00470-2](#)

Khoshandam, M., Soltaninejad, H.,

Mousazadeh, M., Hamidieh, A. A., &

Hosseinkhani, S. (2023). Clinical

applications of the CRISPR/Cas9

genome-editing system: Delivery

options and challenges in precision

medicine. *Genes & Diseases*, 11(1),

268–282.

<https://doi.org/10.1016/j.gendis.2023.02.>

027

Levine, R. M., Dempsey, E. A., Nimmagadda,

N., Jang, Y., Mayberry, K., Zhang, J.,

Cullins, D., Willis, C., Powers, A.,

Kim-Hoehammer, Y.-I., Lin, J., Zhang,

F., Gupta, N. D., Lewis, W., Katta, V.,

Li, Y., Lockey, T., Cheng, Y., Lehigh, P.,

... Yen, J. S. (2024). SAGES1: Clinical

Translation of CRISPR Genome Editing

Strategy to Induce Fetal Hemoglobin to

Treat Sickle Cell Disease. *Blood*,

144(Supplement 1), 146.

<https://doi.org/10.1182/blood-2024-2111>

82

Lim, J. M., & Kim, H. H. (2022). Basic

principles and clinical applications of
CRISPR-based genome editing. *Yonsei
Med J*, 63(2), 105.

<https://doi.org/10.3349/ymj.2022.63.2.10>

5

Lopes, R., & Prasad, M. K. (2024). Beyond the

promise: evaluating and mitigating
off-target effects in CRISPR gene
editing for safer therapeutics. *Frontiers
in Bioengineering and Biotechnology*,
11, 1339189.

<https://doi.org/10.3389/fbioe.2023.13391>

89

Obeagu, E. I., Akinleye, C. A., & Obeagu, G. U.

(2024). The role of nucleation in sickle
cell pathophysiology: opportunities for
innovative treatments. *Annals of*

Medicine and Surgery, 87(3),

1341–1350.

<https://doi.org/10.1097/ms9.000000000000002705>

002705

Sharma, A., Boelens, J.-J., Cancio, M., Hankins,

J. S., Bhad, P., Azizy, M., Lewandowski,
A., Zhao, X., Chitnis, S., Peddinti, R.,

Zheng, Y., Kapoor, N., Ciceri, F.,

MacLachlan, T., Yang, Y., Liu, Y., Yuan,

J., Naumann, U., Yu, V. W. C., ...

LaBelle, J. L. (2023a). CRISPR-Cas9

Editing of the HBG1 and HBG2

Promoters to Treat Sickle Cell Disease.

Mass Medical Soc, 389(9), 820–832.

<https://doi.org/10.1056/nejmoa2215643>

Sharma, A., Boelens, J.-J., Cancio, M., Hankins,

J. S., Bhad, P., Azizy, M., Lewandowski,

A., Zhao, X., Chitnis, S., Peddinti, R.,

Zheng, Y., Kapoor, N., Ciceri, F.,

MacLachlan, T., Yang, Y., Liu, Y., Yuan,

J., Naumann, U., Yu, V. W. C., ...

LaBelle, J. L. (2023b). CRISPR-Cas9

Editing of the HBG1 and HBG2

Promoters to Treat Sickle Cell Disease.

Mass Medical Soc, 389(9), 820–832.

<https://doi.org/10.1056/nejmoa2215643>

approaches: a review.

Journals.Lww.Com, 86(10), 5938–5946.

<https://doi.org/10.1097/ms9.000000000000002478>

Sharma, A., Boelens, J.-J., Cancio, M., Hankins,

J. S., Bhad, P., Azizy, M., Lewandowski,

A., Zhao, X., Chitnis, S., Peddinti, R.,

Zheng, Y., Kapoor, N., Ciceri, F.,

MacLachlan, T., Yang, Y., Liu, Y., Yuan,

J., Naumann, U., Yu, V. W. C., ...

LaBelle, J. L. (2023c). CRISPR-Cas9

Editing of the HBG1 and HBG2

Promoters to Treat Sickle Cell Disease.

New England Journal of Medicine,

389(9), 820–832.

<https://doi.org/10.1056/nejmoa2215643>

Tariq, H., Khurshid, F., Khan, M. H., Dilshad,

A., Zain, A., Rasool, W., Jawaaid, A.,

Kunwar, D., Khanduja, S., & Akbar, A.

(2024). CRISPR/Cas9 in the treatment of

sickle cell disease (SCD) and its

comparison with traditional treatment

Xu, L., Lahiri, P., Skowronski, J., Bhatia, N.,

Lattanzi, A., & Porteus, M. H. (2023).

Molecular dynamics of genome editing

with CRISPR-Cas9 and rAAV6 virus in

human HSPCs to treat sickle cell

disease. *Cell.Com*, 30, 317–331.

[https://www.cell.com/molecular-therapy-](https://www.cell.com/molecular-therapy-family/methods/fulltext/S2329-0501(23)00111-0)

[family/methods/fulltext/S2329-0501\(23\)](https://www.cell.com/molecular-therapy-family/methods/fulltext/S2329-0501(23)00111-0)

00111-0

Zarghamian, P., Klermund, J., & Cathomen, T.

(2023). Clinical genome editing to treat

sickle cell disease—a brief update.

Frontiersin.Org, 9, 1065377.

[https://doi.org/10.3389/fmed.2022.10653](https://doi.org/10.3389/fmed.2022.1065377)

77

modeling framework. Recognition is given to laboratory personnel and clinical collaborators whose insights improved the interpretation of experimental controls, particularly regarding gene editing delivery methods and validation strategies . Contributions from instructors and peer reviewers shaped the ethical framework and methodological rigor in genome editing research. Lastly, gratitude is extended to authors of foundational studies referenced in the bibliography, whose analyses of CRISPR-Cas systems were crucial for informing comparative and translational aspects.

Acknowledgments

The completion of this research project was enabled by mentor Jon Davis, professor at North Carolina School of Science and Mathematics, a collaborator who provided critical feedback during study design and modeling. Thanks are due to the institution's faculty and staff for access to computational resources, data analysis platforms, and technical expertise essential for developing and validating the CRISPR