# Hospital Formulary Entry: Paced respiration, Sham comparator: fast shallow breathing

#### **Document Control Information**

Document ID:	MED-PROT-8385
Effective Date:	
Review Date:	2025-12-13
Therapeutic Class:	Clinical Trial
Risk Level:	Medium

# **Standard Dosing Protocol**

Current Standard Dose: Study protocol dosing

Monitoring Requirements: Standard study precautions

#### **Administration Guidelines**

Route of Administration: Subcutaneous

**Frequency:** Three times daily **Duration:** Until discontinued

Patient Population: Clinical trial participants (n=218)

#### **Contraindications and Precautions**

Contraindications: Per trial protocol

Adverse Events: As reported in trial monitoring

## **Detailed Clinical Considerations**

### **Patient Monitoring Protocol**

Regular monitoring is essential for safe administration of Paced respiration, Sham comparator: fast shallow breathing. Current protocols require Standard study precautions. Healthcare providers should assess patient response and adjust therapy accordingly. Baseline laboratory values should be obtained prior to initiation. The mechanism of action involves Under clinical investigation. Clinical efficacy has been demonstrated in Hot Flashes.

# **Special Populations**

Elderly Patients: Dose adjustment may be required based on renal function and overall health status.

Pediatric Patients: Safety and efficacy not established in patients under 18 years.

**Pregnancy:** Use only if potential benefit justifies potential risk to fetus. **Nursing Mothers:** Exercise caution when administering to nursing women.

Last revised: August 2024