

Hospital Formulary Entry: ACLASTA

Document Control Information

Document ID:	MED-PROT-2542
Effective Date:	
Review Date:	2025-10-14
Therapeutic Class:	Adverse Event Report
Risk Level:	Critical

Standard Dosing Protocol

Current Standard Dose: Standard dosing per label

Monitoring Requirements: Standard warnings

Administration Guidelines

Route of Administration: IM

Frequency: Twice daily

Duration: 14 days

Patient Population: Patients similar to case: 2, age 62

Contraindications and Precautions

Contraindications: Enhanced screening for risk factors

Adverse Events: Nasopharyngitis, Malaise, Pyrexia

Detailed Clinical Considerations

Patient Monitoring Protocol

Regular monitoring is essential for safe administration of ACLASTA. Current protocols require Standard warnings. Healthcare providers should assess patient response and adjust therapy accordingly. Baseline laboratory values should be obtained prior to initiation. The mechanism of action involves ACLASTA mechanism with identified safety concern. Clinical efficacy has been demonstrated in Nasopharyngitis, Malaise, Pyrexia.

Special Populations

Elderly Patients: Dose adjustment may be required based on renal function and overall health status.

Pediatric Patients: Safety and efficacy not established in patients under 18 years.

Pregnancy: Use only if potential benefit justifies potential risk to fetus.

Nursing Mothers: Exercise caution when administering to nursing women.

Last revised: May 2024