# **Hospital Formulary Entry: LETAIRIS**

## **Document Control Information**

| Document ID:       | MED-PROT-2252        |
|--------------------|----------------------|
| Effective Date:    |                      |
| Review Date:       | 2025-10-14           |
| Therapeutic Class: | Adverse Event Report |
| Risk Level:        | Medium               |

# **Standard Dosing Protocol**

**Current Standard Dose:** Standard dosing per label **Monitoring Requirements:** Standard warnings

## **Administration Guidelines**

Route of Administration: IV Frequency: Once daily Duration: Until discontinued

Patient Population: Patients similar to case: 2, age 35

#### **Contraindications and Precautions**

Contraindications: Enhanced screening for risk factors

Adverse Events: Syncope, Dizziness

## **Detailed Clinical Considerations**

# **Patient Monitoring Protocol**

Regular monitoring is essential for safe administration of LETAIRIS. Current protocols require Standard warnings. Healthcare providers should assess patient response and adjust therapy accordingly. Baseline laboratory values should be obtained prior to initiation. The mechanism of action involves LETAIRIS mechanism with identified safety concern. Clinical efficacy has been demonstrated in Syncope, Dizziness.

# **Special Populations**

Elderly Patients: Dose adjustment may be required based on renal function and overall health status.

Pediatric Patients: Safety and efficacy not established in patients under 18 years.

**Pregnancy:** Use only if potential benefit justifies potential risk to fetus. **Nursing Mothers:** Exercise caution when administering to nursing women.

Last revised: August 2024